

An integrated active case search for skin-NTDs in yaws endemic health districts in Cameroon, Côte d'Ivoire and Ghana

Serges Tchatchouang^{1*}, Laud A. Basing², Kouadio-Aboh Hugues^{3,4}, Becca L. Handley⁵, Camila G-Beiras^{6,7}, Ivy Amanor², Philippe Ndzomo¹, Mohammed Bakheit⁸, Lisa Becherer⁹, Sascha Knauf^{10,11}, Claudia Müller¹⁰, Earnest Njih-Tabah^{12,13}, Theophilus Njmanishi¹², Tania Crucitti¹⁴, Nadine Borst^{9,15}, Simone Lüert¹⁰, Sieghard Frischmann⁸, Helena Gmoser⁹, Emelie Landmann⁸, Aboubacar Sylla³, Mireille S. Kouamé-Sina³, Daniel Arhinful², Patrick Awondo¹, Gely Menguen¹, Emma-Michèle Harding-Esch¹⁶, Adingra Tano³, Mamadou Kaloga¹⁷, Paul Koffi-Aboa¹⁷, Nana Konama-Kotey¹⁸, Oriol Mitjà^{6,7}, Sara Eyangoh¹, Kennedy Kwasi-Addo², Solange Ngazoa-Kakou³, Michael Marks^{5, 16, 19}

Affiliation:

¹ Centre Pasteur du Cameroun, P.O. Box 1274; 451, Street 2005, Yaoundé 2; Yaoundé, Cameroon

² Noguchi Memorial Institute for Medical Research, University of Ghana, Ghana.

³ Institut Pasteur de Cote d'Ivoire, Abidjan, Lagunes, Côte d'Ivoire.

⁴ National Program of African Trypanosomiasis Elimination, Abidjan, Côte d'Ivoire.

⁵ Clinical Research Department, London School of Hygiene & Tropical Medicine, Keppel Street, London, UK

⁶ Skin Neglected Tropical Diseases and Sexually Transmitted Infections section, Hospital Universitari Germans Trias i Pujol; Fight Infectious Diseases Foundation, Badalona, Spain;

⁷ Universitat Autònoma de Barcelona, 08193 Bellaterra (Cerdanyola del Vallès), Spain

⁸ Mast Diagnostica GmbH, Reinfeld, Germany.

⁹ Laboratory for MEMS Applications, IMTEK - Department of Microsystems Engineering, University of Freiburg, Freiburg, Germany.

¹⁰ Institute of International Animal Health/One Health, Friedrich-Loeffler-Institut, Federal

¹¹ Professorship for One Health/International Animal Health, Faculty of Veterinary Medicine, Justus Liebig University, Giessen, Germany.

¹² National Buruli Ulcer, Leprosy, Yaws and Leishmaniasis Control Program, Ministry of Public Health, Yaounde, Centre Region, Cameroon.

¹³ Public Health & Epidemiology, University of Dschang, Dschang, West Region, Cameroon.

¹⁴ Experimental Bacteriology Unit, Institut Pasteur de Madagascar, Antananarivo, Madagascar

¹⁵ Hahn-Schickard, Freiburg, Germany.

34 ¹⁶ Clinical Research Department, London School of Hygiene & Tropical Medicine, Faculty of
35 Infectious and Tropical Diseases, London, UK.

36 ¹⁷ Programme National de Lutte contre l'Ulcère de Buruli, Abidjan, Côte d'Ivoire.

37 ¹⁸ National Yaws Eradication Program, Ghana Health Service, Accra, Ghana.

38 ¹⁹ Hospital for Tropical Diseases, London, UK.

39

40 * **Correspondence:** tchatchouang@pasteur-yaounde.org

41

42 **Short title: Overview of skin-NTDs in West Africa**

43

44 Abstract

45 Background: Integrated approaches to mapping skin Neglected Tropical Diseases (NTDs)
46 may be cost-effective way to guide decisions on resource mobilization. Pilot studies have
47 been carried out, but large-scale data covering multiple countries endemic for skin-NTDs are
48 lacking. Within the LAMP4YAWS project, we collected integrated data on the burden of
49 multiple skin-NTDs.

50 Methods: From March 2021 to March 2023, integrated case searches for yaws alongside other
51 skin conditions were performed in endemic health districts of yaws in Cameroon, Côte
52 d'Ivoire, and Ghana. Initial screening involved a brief clinical examination of participants to
53 determine if any skin conditions were suspected. Cases of skin-NTDs were then referred to a
54 health facility for appropriate management.

55 Results: Overall 61,080 individuals screened, 11,387 (18.6%) had skin lesions. The majority
56 of individuals (>90%) examined were children aged 15 years old and under. The proportion
57 of serologically confirmed yaws cases was 8.6% (18/210) in Cameroon, 6.8% (84/1232) in
58 Côte d'Ivoire, and 26.8% (440/1643) in Ghana. Other skin conditions based on clinical
59 examination included: scabies, Buruli ulcer, leprosy, lymphatic filariasis (lymphoedema and
60 hydrocele), tungiasis, and fungal infections. The most common conditions were scabies and
61 superficial fungal infections (scabies versus fungal infections) in Cameroon with 5.1%
62 (214/4204) versus 88.7% (3730/4204), Côte d'Ivoire with 25.2% (1285/5095) versus 50.4%
63 (2567/5095) and Ghana 20% (419/2090) versus 1.3% (28/2090). Other skin-NTDs were less
64 common across all three countries.

65 Conclusion: This study confirms that integrated screening allows simultaneous detection of
66 multiple skin-NTDs, maximising use of scarce resources.

67

68 **Keywords:** Yaws, integrated surveillance, skin-NTDs, Cameroon, Côte d'Ivoire, Ghana.

69

70 Plain English Summary

71 Many Neglected Tropical Diseases (NTDs) predominantly affect the skin and are referred to
72 as skin-NTDs. The World Health Organization (WHO) has developed a number of strategies
73 for the control, eradication and elimination of skin-NTDs and recognizes the importance of
74 integrated approaches to mapping skin-NTDs. We conducted a study adopting integrated
75 screening for multiple skin-NTDs and other skin conditions in Cameroon, Côte d'Ivoire, and
76 Ghana. This ran alongside a study focused on diagnostic tests for one specific skin-NTD –
77 yaws.

78 The results showed that integrated screening is a feasible and cost-effective way to detect
79 multiple skin-NTDs in a single intervention. Of more than 60,000 individuals screened almost
80 one in five had a skin lesion. Cases of yaws confirmed by blood tests were detected more
81 frequently in Ghana compared to Côte d'Ivoire and Cameroon. The most common skin
82 conditions were scabies and superficial fungal infections. Other skin-NTDs such as Buruli
83 ulcer, leprosy, lymphatic filariasis and tungiasis were less common. Integrated screening
84 allowed detection of skin conditions and co-endemicity of skin-NTDs and the data can guide
85 decisions on resource mobilization to manage skin-NTDs.

86

87

88 INTRODUCTION

89 Skin diseases affect over 900 million people worldwide annually and are the third most
90 frequent reason for seeking care (1-3). A subset of these conditions are also classified by the
91 World Health Organization (WHO) as Neglected Tropical Diseases (NTDs) (4). Of the twenty
92 NTDs recognised by WHO, more than half can present as skin diseases and are referred to as
93 skin-NTDs. If not diagnosed and treated promptly, these conditions can result in
94 disfigurement, disability, stigmatisation, impacts on mental health, and socioeconomic
95 difficulties (4-6).

96 In Cameroon, Côte d'Ivoire, and Ghana, several skin-NTDs are known to be endemic
97 including yaws (7-10), Buruli ulcer (11-13), leishmaniasis (14-16), leprosy (10, 17, 18),
98 lymphatic filariasis (19-21), mycetoma (22-24), onchocerciasis (25-27), and scabies (10, 28-
99 30). Alongside these conditions, a number of more common skin conditions including
100 superficial fungal and bacterial infections cause a considerable burden of disease (28, 31-33).

101 Most data on skin-NTDs rely on passive surveillance activities, which are likely to
102 underestimate the true burden of disease as they are dependent on patients who present to a
103 health centre, receive a diagnosis and are reported to the respective disease control
104 programme. The highest burden of skin-NTDs is believed to be in communities living in
105 difficult to reach areas with limited access to healthcare (34, 35). Consequently, routine data
106 are likely to underreport the true burden and epidemiology of these conditions.

107 To achieve the WHO goals for skin-NTDs, active case detection will be required. Therefore,
108 WHO has recently promoted integrated activities of skin-NTDs in order to accelerate progress
109 in their control, elimination and eradication (4, 36). Because NTDs often overlap in
110 geographic areas, integrated active case searching may offer a way to overcome some of the
111 challenges associated with stand-alone programmes and reduce the risk of late detection of
112 skin-NTDs. Integration can also result in a reduction of resources, thereby increasing cost-
113 effectiveness and allow for an expanded coverage of interventions (2, 36-38).

114 The LAMP4YAWS project in Cameroon, Côte d'Ivoire, and Ghana aimed to implement and
115 evaluate a loop-mediated isothermal amplification test for yaws. As part of the project, field
116 activities included active case searching of yaws using an integrated approach (39). Here we
117 report data on skin conditions from active case searches in yaws-endemic health districts from
118 all three countries.

119

120 **METHODS**

121 **Study design**

122 From March 2021 to March 2023, we undertook integrated case searches in rural
123 communities within suspected yaws-endemic health districts in Cameroon, Côte d’Ivoire, and
124 Ghana. Integrated approach for skin-NTDs offers possibility to address several diseases
125 simultaneously in the same communities in order to maximize the utilization of limited
126 resources (40). This approach was implemented by assessing burden of yaws in health
127 districts from notifications of NTD national programs, training of health workers and village
128 volunteers, conducting active case detection of skin conditions, clinical diagnosis and rapid
129 tests for yaws, treating yaws cases and referring other skin conditions to healthcare centres
130 (2). In each community visited, initial screening involved an examination to determine if
131 participants had skin lesions as described in the study protocol (39).

132 **Study sites**

133 The study was conducted in rural areas in 12 health districts in Cameroon, six in Côte
134 d’Ivoire, and seven in Ghana (39). The characteristics of the study sites are summarised in
135 Table 1. Study sites were selected based on clinical notifications of yaws by the national
136 programmes on NTDs.

137 **Recruitment of participants**

138 Case searches were conducted in collaboration with national NTD programmes. District
139 health staff were first trained to recognise skin-NTDs as well as other skin conditions.
140 Training materials included the WHO manual on recognition of skin-NTDs (41), as well as
141 other national NTD training materials.

142 Individuals presenting with yaws-like lesions underwent SD Bioline (Abbott, USA) test to
143 detect anti-treponemal antibodies. Those reactive underwent a Chembio Dual Path Platform
144 (DPP) test (Chembio Diagnostics, New York, USA) for simultaneous detection of both
145 treponemal and non-treponemal antibodies, interpreted as serologically confirmed yaws case.
146 Samples were then taken as part of the main study from participants as described elsewhere
147 (39). Other skin diseases were recorded and referred to the health district for management
148 following the national guidelines.

Several approaches were used for case finding, including community-based case detection, detection in healthcare centres, primary school-based screening programs, screening alongside mass drug administration (MDA) campaigns, church and house-to-house case searches (Table 1). The strategy was chosen based on other activities taking place in the selected district. The research team visited different study sites according to a pre-established schedule that was communicated to health workers, village chiefs, and school directors before the appointed day.

Table 1: Characteristic of study sites in Cameroon, Côte d'Ivoire and Ghana.

Country	Region	Health district	Sub-divisions	Population	Active case searching strategy
Cameroon	East	Doumé	Doumé, Doumaitang, Dimako	53,286	Community, House-to-house
		Bétaré-Oya	Betaré-Oya, Ngoura	141,405	Community, health centres
		Batouri	Batouri, Ndemnam	121,199	Community, health centres
		Messamena	Messamena, Somalomo	48,000	Community, health centres
		Abong-Mbang	Abong-Mbang, Atok, Angossas, Dja (Mindourou)	85,741	Community, health centres
		Lomié	Lomié, Messok, Ngoyla	47,695	Community, health centres
		Yokadouma	Gari-Gombo, Yokadouma	126,353	Community, health centres
		Mbang	Mbang	34,148	Community, House-to-house
	South	Djoum	Djoum, Oveng, Mintom	43,142	MDA
		Lolodorf	Bipindi, Lolodorf,	43,446	Community

			Mvengue, Akom 2		
		Sangmélina	Sangmelima , Meyomessi	110,385	School, community
	Adamawa	Bankim	Bankim	121,756	School, community, church
Côte d'Ivoire	Loh Djiboua	Divo	Goh- Djiboua	458,318	School, Community
	Agneby Tiassa	Tiassalé	Lagunes	270,412	
	Bélier	Toumodi	Lacs	189,794	
		Autonomou s district of Yamoussou kro	N/A	446,158	
	Haut sassandra	Daloa	Sassandra- Marahou	745,628	
		Vavoua	Sassandra- Marahou	451,492	
Ghana	Central	Asikuma Odoben Brakwa	N/A	129,580	School, Community
		Ajumako Enyan Essiam	N/A	138,046	
	Eastern	Upper West Akim	N/A	93,391	
		Ayensuanor	N/A	94,594	
		Akuapim North	N/A	105,315	
		Asene Manso Akroso	N/A	77,498	
	Western	Mpohor	N/A	52,473	

157 MDA: Mass drug administration; N/A: not applicable

158 **Case definition of other skin conditions**

159 In order to be considered a case of leprosy (Hansen's Disease), an individual must have one or
160 more of skin lesions that were hypopigmented or red and had definite loss of sensation.

Buruli ulcer case was based on clinical presentation consistent with a painless, non-healing ulcer with undermined edges and a necrotic base, typically located on the limbs; a painless nodule or plaque, typically located on the limbs; and oedema of a limb or other body part, with or without overlying ulceration.

Scabies were defined as people with rash consisting of papules, vesicles, and burrows, typically located on the hands, feet, and folds of the skin.

An individual suffering from lymphatic filariasis presented lymphedema characterised by the enlargement of body parts in the limbs and breasts; or hydrocele marked by swelling in the scrotal sack.

Case of tungiasis was defined as participant with a painful, pruritic nodule with a central black dot, mainly located on the feet, toes, or fingers.

Superficial fungal infections were defined as affection on the body commonly with a red, ring-shaped rash with a raised border for *Tinea corporis* (ringworm); scaly patches on the scalp sometime with hair loss for *Tinea capitis* (scalp ringworm); light or dark patches on the skin, most commonly on the chest and back for *Tinea versicolor* (pityriasis versicolor).

Data analyses

Study data were recorded using standardized data collection forms. We used basic descriptive statistics to summarise the characteristics of the individuals who were screened, the number who were diagnosed with each skin-NTD, and other skin conditions. Results are shown stratified by country. Analyses were performed in Microsoft Excel 2013.

Ethics approval

This study was approved in Cameroon by the National Ethics Committee on Research for Human Health (CNERSH) under decision N°2020/12/1327/CE/CNERSH/SP and administrative authorization of the Ministry of Public Health N°D30-308/L/MINSANTE/SG/DROS), in Côte d'Ivoire by the Comité National d'Ethique des Sciences de la Vie et de la Santé (IQR) through decision IQRG0075_16/09/2020, in Ghana by the Ghana Health Service of Ethical Review Committee (GHS-ERC) and Noguchi Memorial Institute for Medical Research-Institutional Review Board (NMIMR-IRB) under decisions GHS-ERC 005/12/20 and NMIMR-IRB CPS 019/20-21), and in United Kingdom through London School of Hygiene and Tropical Medicine (LSHTM) under decision N° 21633 (19 August 2021).

192 All participants and/or the parents/legal guardians of minors provided written informed
193 consent before enrolment. In addition, assent was obtained from child participants.

194

RESULTS

Demographic characteristics

Case searches were conducted in 12 districts in Cameroon, six in Cote d'Ivoire, and seven in Ghana. A total of 61,080 individuals were screened, including 20,414 in Cameroon, 16,530 in Côte d'Ivoire, and 24,136 in Ghana. The male to female ratio was approximately 1:1 in the three countries (10804/9610, 8430/8100, and 13241/10895 respectively for Cameroon, Côte d'Ivoire and Ghana), with more than 90% of participants aged up to 15 years old (56471/61080).

Skin lesions detected

Table 2 summarizes the skin diseases detected in Cameroon, Côte d'Ivoire, and Ghana. Skin lesion accounted for 18.6% (11389 /61080) and included 20.6% (4204/20414) in Cameroon, 30.8% (5095/16530) in Côte d'Ivoire and 8.7% (2090/24136) in Ghana.

The most frequent skin conditions encountered in Cameroon were *Tinea capitis*, scabies and yaws-like lesions. *T. capitis* accounted for 82.7% (3478/4204) of the skin conditions and was detected mainly in the health districts of Bankim (44.6%, 1552/3478) and Sangmélima (16.7%, 580/3478)). Scabies was the second most common skin condition detected in 5.1% (214/4204) cases of skin conditions seen. The majority of scabies cases were again detected in Bankim and Sangmélima with 37.9% (81/214) and 23.8% (51/214) of scabies among the detected skin lesions, respectively. Yaws-like lesions were the third most common skin condition (5%, 210/4204) and were identified in high numbers in Sangmélima (23.3%, 49/210), Djoum (22.4%, 47/210) and Mbang (15.7%, 33/210). However, only 18 individuals were serologically confirmed as yaws cases almost all among Baka population.

Pityriasis versicolor, scabies, and yaws-like lesions accounted for 99.8% (5084/5095) of skin conditions detected in Côte d'Ivoire. *P. versicolor* was the most common skin condition (50.4% (2567/5095)) across the six health districts visited. Scabies represented 25.2% (1285/5095) of the conditions and yaws-like lesions were reported among 24.2% of individuals presenting with a skin condition (1232/5095) and were serologically confirmed in 84 people.

In Ghana, yaws-like lesions accounted for 78.6% (1643/2090) of them, with serologically confirmed yaws among 440 individuals. Upper West Akim District and Asene Manso Akroso

225 recorded the highest number of yaws cases. Scabies and *T. capitis* accounted for 20%
226 (419/2090) and 1.3% (28/2090) of skin conditions, respectively.

227 **Table 2: Characteristics of skin diseases found in Cameroon, Côte d'Ivoire and Ghana**

Country	Health District	Population screened	Leprosy	Scabies	Lymphoedema	Hydrocele	<i>Pityriasis versicolor</i>	<i>Tinea capitis</i>	<i>Tinea corporis</i>	Suspected cases of BU	Suspected cases of yaws	Sd positive	DPP positive
Cameroon	Abong-Mbang	223	0	3	0	0	0	16	0	2	7	0	0
	Bankim	6911	2	81	0	0	112	1552	67	0	27	0	0
	Batouri	268	0	0	0	0	0	3	1	0	0	0	0
	Bétaré-Oya	1281	5	3	0	8	0	121	9	0	6	0	0
	Djoum	1344	0	7	1	0	0	241	1	0	47	1	1
	Doumé	1141	6	2	0	0	22	238	0	2	7	0	0
	Lolodorf	1102	0	10	0	0	4	102	1	1	5	0	0
	Lomié	1868	0	5	3	0	0	119	0	1	11	3	2
	Mbang	976	0	21	0	2	25	199	5	0	33	19	15
	Messamena	814	1	27	0	2	0	97	0	7	4	0	0
	Sangmélina	3150	0	51	2	0	0	580	0	0	49	0	0
	Yokadouma	1336	1	4	0	1	5	210	0	1	14	1	0
	Sub-total	20414	15	214	6	13	168	3478	84	14	210	24	18
Côte d'Ivoire	Divo	3080	1	180	0	0	134	187	39	2	112	12	11
	Tiassalé	4830	0	430	1	3	289	405	88	3	473	87	54
	Yamoussoukro	3700	0	221	0	0	144	201	42	0	259	16	12
	Toumodi	2160	0	132	0	0	115	161	34	0	151	5	4
	Daloa	2100	0	215	0	0	192	268	57	0	191	7	3
	Vavoua	660	1	107	0	0	86	120	25	0	46	0	0
	Sub-total	16530	2	1285	1	3	960	1342	285	5	1 232	127	84
Ghana	Ayensuanor	3119	0	11	0	0	0	0	0	0	194	29	23
	Asikuma	3496	0	0	0	0	0	0	0	0	201	56	43

	Odoben Brakwa												
	Asene Manso Akroso	3331	0	86	0	0	0	14	0	0	306	204	196
	Akuapim North	3817	0	14	0	0	0	3	0	0	201	27	16
	Mpohor	3323	0	86	0	0	0	1	0	0	206	80	74
	Ajumako Enyan Essiam	2864	0	98	0	0	0	2	0	0	211	15	8
	Upper West Akim	4186	0	124	0	0	0	8	0	0	324	89	80
	Sub-total	24136	0	419	0	0	0	28	0	0	1643	500	440

228 DPP+: Positive Dual Path Platform

229

230

DISCUSSION

In accordance with the NTD road map 2021-2030 (4), skin-NTDs are targeted for eradication (yaws), elimination (leprosy, onchocerciasis, lymphatic filariasis) and control (other skin-NTDs), and an integrated approach is the backbone to achieving these targets. The main study was focused on yaws while adopting an integrated approach in the screening of individuals in the selected health districts.

This project improved case detection of skin conditions such as yaws, scabies, lymphatic filariasis, Buruli ulcer, leprosy, tungiasis and superficial fungal infections in the three countries. Yaws and scabies were the predominant skin-NTDs.

The frequency of detection of serologically confirmed yaws among suspected cases in Cameroon, Côte d'Ivoire and Ghana was 8.6%, 6.8% and 26.8% respectively. These data show that yaws is more prevalent in endemic health districts in Ghana compared to Cameroon and Côte d'Ivoire, which is consistent with routine reporting data in Ghana with frequencies above 15% among suspected cases of yaws (34, 42, 43). Ghana has the second highest population density in West Africa and this can make it more difficult to control the spread of yaws. For instance, many MDA campaigns have been performed previously but the incidence of yaws is still high in many parts of the country (42, 44, 45). The data obtained in Cameroon overlap with previous findings, with less than 10% of yaws over suspected cases (46, 47) but lower to what has been reported during an outbreak (8). The decrease of yaws cases in Cameroon is linked to the surveillance impact by identifying and treating cases, and MDA implementation in the Congo Basin in December 2020 before the current study was conducted (48). The results of Côte d'Ivoire show that yaws remains prevalent in the study sites (1.6%; 81/5095) compared to findings in the Divo health district between 2016 and 2017 that identified 8 cases of yaws out of 1,302 individuals with skin conditions (10). Côte d'Ivoire is one of the countries in Africa that has been most affected by skin-NTDs, including yaws (49, 50). The government of Côte d'Ivoire has implemented a number of programmes to address these diseases. Areas of co-endemicity of skin-NTDs have undergone several investigations with treatment which have certainly reduced the incidence of yaws (10). The results of this study suggest that additional efforts are needed to control yaws in the three countries where the disease is endemic, in particular the number of rounds of MDA which can be reviewed to significantly reduce the incidence of the disease (51). More than 80% of yaws like ulcers were not yaws (542/3085), suggesting that proper identification and management of other cutaneous ulcers, such as *H. ducreyi*, is important to improve patient outcomes.

Scabies was the most common skin-NTD detected among participants with skin conditions. The frequency was 25.2% in Côte d'Ivoire, 19.8% in Ghana and 5.1% in Cameroon. These data are high compared to previous studies in the three countries where proportions of scabies in people with skin lesions ranged almost from 3 to 9% during active case search. Difference may be due to a number of factors, including the duration of the screening, the context of the study and the methods used to detect scabies. The field activities of the current study were carried out over almost three years unlike the previous ones whose duration was over a few months. Earlier data from the three countries highlight the variability in the proportions of scabies depending on the context and methods to detect in hospital, prison, boarding school or even during active research in the community and in schools. Thus, data from several regions in Côte d'Ivoire show the proportions of scabies during active case searches among people suffering from skin diseases varied between 3.7% and 8.6% (10, 49, 52). In Ghana, recent findings from 2019 show that the proportions of scabies are between 6 to 12% among patients suffering from skin infections reported to urban and rural study health facilities (30, 53), and prevalences of 11.2% to 71% during outbreaks (54, 55). In Cameroon, surveys in urban areas in boarding schools show the prevalence of scabies of nearly 18% (56), 32% in prison (57), and a proportion of 2.8% during active case search in rural areas among respondents with skin disease (28). These data suggest that scabies is a major public health problem in all three countries.

Other skin-NTDs such as leprosy, Buruli ulcer, lymphatic filariasis and tungiasis were less common. Leprosy cases were not found in Ghana while in Cameroon and Côte d'Ivoire, leprosy cases varied between 2 and 15 throughout the study period. Available data show that some districts in Cameroon and Côte d'Ivoire are still of high endemicity (17, 31, 52). No case from Ghana could be due to the significant progress in controlling leprosy or targeted study sites. Buruli ulcer and lymphatic filariasis case numbers followed the same trend as leprosy in Cameroon and Côte d'Ivoire, and tungiasis was only recorded in Cameroon. Buruli ulcer and lymphatic filariasis are known to be endemic in Cameroon, Côte d'Ivoire and Ghana (11-13, 19-21) suggesting that the diseases may have similar risk factors, and that tungiasis may be more common in Cameroon due to specific environmental factors.

The study also found that superficial fungal infections were the most common form of skin disease in two of the three countries. In Cameroon, these conditions accounted for almost 90% with the majority due to *Tinea capitis* while *Pityriasis versicolor* constituted the bulk of these infections in Côte d'Ivoire (50.4%). The findings are concordant with field data in

Cameroon, Côte d'Ivoire and Togo highlighting fungal infections as the main skin conditions (10, 28, 31, 58). In previous investigation, proportion of *Pityriasis versicolor* and *Tinea capitis* among diagnosed skin diseases in Côte d'Ivoire was approximately 45% of each (31); while in Cameroon proportion of *Pityriasis versicolor* in a rural setting was around 21% (28). In Ghana, superficial fungal infections were likely to be less common. Previous data indicated that fungal infections accounted for between 20 and 30% of patients with skin diseases (59, 60). In this study, superficial fungal infections were not automatically recorded in Ghana, thus, only *Tinea capitis* was detected in 1.3% of individuals. In previous studies in Ghana, *Tinea capitis* was one of the main fungal agents with proportions above 20% (59).

We found that the proportion of people with skin lesions was not the same from one country to another and according to the diseases. This is likely due to a variety of factors, such as ecological zones (climate, vegetation and culture), population density and standard of life. However, early diagnosis and treatment are important to prevent the spread and to reduce the risk of complications. Reducing the burden of skin conditions may require improved access to clean water and sanitation facilities, education of people about the diseases and their prevention measures, availability and affordability of treatment, and reinforcement of public health measures to control the spread of diseases such as mass drug administration programmes.

All data were recorded at the health district level and subsequently transmitted to the respective NTD programmes. Through these data, we can see the diversity that can exist in terms of the burden of different diseases in the communities studied, hence the importance of pooling efforts to overcome NTDs. The studies that have been conducted in Côte d'Ivoire (10, 31), Benin (61) and Burkina Faso(62), are examples of integrated approaches to NTD control, elimination and eradication. These studies have shown that integrated approaches can be effective in reducing the prevalence of NTDs and improving the health of people in these countries. Moreover, since many skin-NTDs are zoonotic or have a (possible) non-human reservoir, a One Health guided approach should be used to handle integrated case search activities (63). Synergies between skin-NTD programs and, for example, neglected zoonotic disease programs (such as rabies elimination) could optimize the use of resources (logistics) and stimulate cross-sectoral collaboration that is key for the sustainable control of NTDs.

This study has some limitations. The skin-NTDs reported apart from yaws were only suspected cases because the diagnoses were made based on clinical examinations. Unfortunately, the additional diagnostic tools required for confirmation of the clinical

diagnosis were not available for this investigation. However, the data collected are still useful for guiding where skin-NTD eradication, elimination or control work should be directed in the future, i.e., data collection, training, diagnostics and the needs for medical care in these remote areas. We also acknowledge that the field visits were focused on areas where suspected yaws cases were reported which constitutes a bias for the estimation of other skin-NTDs.

This study highlights how the use of existing disease-specific resources can be maximized to determine the burden of other NTDs. Integration, technical and scientific progress are all pillars of the road map to end the suffering caused by NTDs by 2030.

CONCLUSION

We demonstrate improved data reporting of skin-NTDs when an integrated approach is used. This study contributes to the strengthening of the yaws surveillance system into a guided integrated system that is being put in place throughout each country. The detection of yaws cases in areas that have undergone MDA is remarkable and highlights the importance of rigorous active surveillance ideally with test confirmation of clinically suspected cases and optimal coverage of MDA campaigns. The main challenges remain the difficulty of accessing some communities where yaws is endemic, the follow-up of patients, and the inter-community movements which are sources of reinfection, and resources to manage the affected populations.

ACKNOWLEDGEMENTS

This study is part of the Lamp4Yaws EDCTP funded project (grant number RIA2018D-2495). Our gratitude goes towards all participants and individuals involved in the field.

355 REFERENCES

- 356 1. Hay RJ, Johns NE, Williams HC, Bolliger IW, Dellavalle RP, Margolis DJ, et al. The global
357 burden of skin disease in 2010: an analysis of the prevalence and impact of skin conditions. *J*
358 *Invest Dermatol.* 2014;134(6):1527-34.
- 359 2. Mitjà O, Marks M, Bertran L, Kollie K, Argaw D, Fahal AH, et al. Integrated Control and
360 Management of Neglected Tropical Skin Diseases. *PLoS Negl Trop Dis.*
361 2017;11(1):e0005136.
- 362 3. Ryan TJ. A brief history of tropical dermatology. *Clin Dermatol.* 2009;27(3):248-51.
- 363 4. WHO. Ending the neglect to attain the Sustainable Development Goals: A road map for
364 neglected tropical diseases 2021–2030. World Health Organization; 2020. Licence: CC BY-
365 NC-SA 3.0 IGO.
- 366 5. Asiedu K, Marks M, Hay R. Podoconiosis, skin-NTDs and global health. *Transactions of the*
367 *Royal Society of Tropical Medicine and Hygiene.* 2020;114(12):887-8.
- 368 6. Tora A, Mengiste A, Davey G, Semrau M. Community Involvement in the Care of Persons
369 Affected by Podoconiosis-A Lesson for Other Skin NTDs. *Tropical medicine and infectious*
370 *disease.* 2018;3(3).
- 371 7. Boock AU, Awah PK, Mou F, Nichter M. Yaws resurgence in Bankim, Cameroon: The
372 relative effectiveness of different means of detection in rural communities. *PLoS Negl Trop*
373 *Dis.* 2017;11(5):e0005557.
- 374 8. Ndzomo Ngono JP, Tchatchouang S, Noah Tsanga MV, Njih Tabah E, Tchualeu A, Asiedu
375 K, et al. Ulcerative skin lesions among children in Cameroon: It is not always Yaws.
376 2021;15(2):e0009180.
- 377 9. Ghinai R, El-Duah P, Chi KH, Pillay A, Solomon AW, Bailey RL, et al. A cross-sectional
378 study of 'yaws' in districts of Ghana which have previously undertaken azithromycin mass
379 drug administration for trachoma control. *PLoS Negl Trop Dis.* 2015;9(1):e0003496.
- 380 10. Koffi AP, Yao TAK, Barogui YT, Diez G, Djakeaux S, Zahir MH, et al. Integrated approach
381 in the control and management of skin neglected tropical diseases in three health districts of
382 Côte d'Ivoire. *BMC public health.* 2020;20(1):517.
- 383 11. Marion E, Landier J, Boisier P, Marsollier L, Fontanet A, Le Gall P, et al. Geographic
384 expansion of Buruli ulcer disease, Cameroon. *Emerging infectious diseases.* 2011;17(3):551-
385 3.
- 386 12. Porten K, Sailor K, Comte E, Njikap A, Sobry A, Sihom F, et al. Prevalence of Buruli ulcer
387 in Akonolinga health district, Cameroon: results of a cross sectional survey. *PLoS Negl Trop*
388 *Dis.* 2009;3(6):e466.

- 389 13. Gyamfi E, Dogbe MA, Quaye C, Affouda AA, Kyei-Baffour E, Awuku-Asante D, et al.
390 Variable Number Tandem Repeat Profiling of *Mycobacterium ulcerans* Reveals New
391 Genotypes in Buruli Ulcer Endemic Communities in Ghana and Côte d'Ivoire. *Frontiers in*
392 *microbiology*. 2022;13:872579.
- 393 14. Ngouateu OB, Dondji B. Leishmaniasis in Cameroon and neighboring countries: An
394 overview of current status and control challenges. *Current Research in Parasitology & Vector-*
395 *Borne Diseases*. 2022;2:100077.
- 396 15. Akuffo R, Sanchez C, Chicharro C, Carrillo E, Attram N, Mosore M-T, et al. Detection of
397 cutaneous leishmaniasis in three communities of Oti Region, Ghana. *PLoS Negl Trop Dis*.
398 2021;15(5):e0009416.
- 399 16. Eholié SP, Tanon AK, Folquet-Amorissani M, Doukouré B, Adoubryn KD, Yattara A, et al.
400 [Three new cases of visceral leishmaniasis in Côte d'Ivoire]. *Bull Soc Pathol Exot*.
401 2008;101(1):60-1.
- 402 17. Tabah EN, Nsagha DS, Bissek AZ, Bratschi MW, Njamnshi TN, Plushke G, et al. The
403 Burden of Leprosy in Cameroon: Fifteen Years into the Post-elimination Era. *PLoS Negl*
404 *Trop Dis*. 2016;10(10):e0005012.
- 405 18. Ofosu AA, Bonsu G. Case control study to determine the factors associated with leprosy in
406 the sene district, brong ahafo region of ghana. *Ghana Med J*. 2010;44(3):93-7.
- 407 19. Deribe K, Bakajika DK, Zoure HM, Gyapong JO, Molyneux DH, Rebollo MP. African
408 regional progress and status of the programme to eliminate lymphatic filariasis: 2000-2020.
409 *International health*. 2020;13(Suppl 1):S22-s7.
- 410 20. de Souza DK, Osei-Poku J, Blum J, Baidoo H, Brown CA, Lawson BW, et al. The
411 epidemiology of lymphatic filariasis in Ghana, explained by the possible existence of two
412 strains of *Wuchereria bancrofti*. *The Pan African medical journal*. 2014;17:133.
- 413 21. Simpson H, Konan DO, Brahima K, Koffi JdA, Kashindi S, Edmiston M, et al. Effectiveness
414 of community-based burden estimation to achieve elimination of lymphatic filariasis: A
415 comparative cross-sectional investigation in Côte d'Ivoire. *PLOS Global Public Health*.
416 2022;2(8):e0000760.
- 417 22. Mendouga Menye CRB, Kouotou EA, Atangana PJA. [Contribution of histopathology in the
418 diagnosis of mycetoma in a Cameroonian trader and possibility of an urban contamination].
419 *Journal de mycologie medicale*. 2017;27(3):417-20.
- 420 23. Adoubryn KD, Koffi KE, Troh E, Doukoure B, Kouadio-Yapo CG, Ouhon J, et al. Les
421 mycétomes autochtones de Côte d'Ivoire : caractères épidémiologiques et étiologiques des cas
422 confirmés. *Journal de mycologie medicale*. 2009;19(2):71-6.

423 24. Oladele RO, Ly F, Sow D, Akinkugbe AO, Ocansey BK, Fahal AH, et al. Mycetoma in West
424 Africa. Transactions of the Royal Society of Tropical Medicine and Hygiene.
425 2021;115(4):328-36.

426 25. Siewe JFN, Ngarka L, Tatah G, Mengnjo MK, Nfor LN, Chokote ES, et al. Clinical
427 presentations of onchocerciasis-associated epilepsy (OAE) in Cameroon. Epilepsy & behavior
428 : E&B. 2019;90:70-8.

429 26. Biritwum NK, de Souza DK, Asiedu O, Marfo B, Amazigo UV, Gyapong JO.
430 Onchocerciasis control in Ghana (1974-2016). Parasites & vectors. 2021;14(1):3.

431 27. Koudou BG, Kouakou M-M, Ouattara AF, Yeo S, Brika P, Meite A, et al. Update on the
432 current status of onchocerciasis in Côte d'Ivoire following 40 years of intervention: Progress
433 and challenges. PLoS Negl Trop Dis. 2018;12(10):e0006897.

434 28. Bissek A-CZ-K, Tabah EN, Kouotou E, Sini V, Yepnjio FN, Nditanchou R, et al. The
435 spectrum of skin diseases in a rural setting in Cameroon (sub-Saharan Africa). BMC
436 Dermatology. 2012;12(1):7.

437 29. Kouotou EA, Nansseu JR, Sieleunou I, Defo D, Bissek AC, Ndam EC. Features of human
438 scabies in resource-limited settings: the Cameroon case. BMC Dermatol. 2015;15:12.

439 30. Boateng LA. Healthcare-seeking behaviour in reporting of scabies and skin infections in
440 Ghana: A review of reported cases. Transactions of the Royal Society of Tropical Medicine
441 and Hygiene. 2020;114(11):830-7.

442 31. Yotsu RR, Kouadio K, Vagamon B, N'guessan K, Akpa AJ, Yao A, et al. Skin disease
443 prevalence study in schoolchildren in rural Côte d'Ivoire: Implications for integration of
444 neglected skin diseases (skin NTDs). PLoS Negl Trop Dis. 2018;12(5):e0006489.

445 32. Bourée P, Simeni Njonou R, Takougang I, Kaptue L. [Tungiasis in Bangou (West
446 Cameroon)]. Medecine et sante tropicales. 2012;22(4):440-3.

447 33. Deka MA. Mapping the Geographic Distribution of Tungiasis in Sub-Saharan Africa.
448 Tropical medicine and infectious disease. 2020;5(3).

449 34. Kazadi WM, Asiedu KB, Agana N, Mitjà O. Epidemiology of yaws: an update. Clin
450 Epidemiol. 2014;6:119-28.

451 35. Yotsu RR, Fuller LC, Murdoch ME, van Brakel WH, Revankar C, Barogui MYT, et al. A
452 global call for action to tackle skin-related neglected tropical diseases (skin NTDs) through
453 integration: An ambitious step change. PLoS Negl Trop Dis. 2023;17(6):e0011357.

454 36. WHO. Ending the neglect to attain the sustainable development goals: a strategic framework
455 for integrated control and management of skin-related neglected tropical diseases. World
456 Health Organization; 2022. Licence: CC BY-NC-SA 3.0 IGO..

- 457 37. Molyneux DH, Hotez PJ, Fenwick A. "Rapid-impact interventions": how a policy of
458 integrated control for Africa's neglected tropical diseases could benefit the poor. *PLoS Med.*
459 2005;2(11):e336.
- 460 38. Engelman D, Fuller LC, Solomon AW, McCarthy JS, Hay RJ, Lammie PJ, et al.
461 Opportunities for Integrated Control of Neglected Tropical Diseases That Affect the Skin.
462 *Trends in parasitology.* 2016;32(11):843-54.
- 463 39. Handley BL, González-Beiras C, Tchatchouang S, Basing LA, Hugues KA, Bakheit M, et al.
464 LAMP4yaws: *Treponema pallidum*, *Haemophilus ducreyi* loop mediated isothermal
465 amplification - protocol for a cross-sectional, observational, diagnostic accuracy study. *BMJ*
466 *open.* 2022;12(3):e058605.
- 467 40. Chandler DJ, Fuller LC. The Skin-A Common Pathway for Integrating Diagnosis and
468 Management of NTDs. *Tropical medicine and infectious disease.* 2018;3(3).
- 469 41. WHO. Recognizing neglected tropical diseases through changes on the skin. Geneva: World
470 Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO.
- 471 42. Abdulai AA, Agana-Nsiire P, Biney F, Kwakye-Maclean C, Kyei-Faried S, Amponsa-
472 Achiano K, et al. Community-based mass treatment with azithromycin for the elimination of
473 yaws in Ghana—Results of a pilot study. *PLoS Negl Trop Dis.* 2018;12(3):e0006303.
- 474 43. Basing LA, Djan M, Simpson SV, Adu-Sarkodie Y. Mapping of yaws endemicity in Ghana;
475 Lessons to strengthen the planning and implementation of yaws eradication. *medRxiv.* 2020.
- 476 44. Marks M, Mitjà O, Bottomley C, Kwakye C, Houinei W, Bauri M, et al. Comparative
477 efficacy of low-dose versus standard-dose azithromycin for patients with yaws: a randomised
478 non-inferiority trial in Ghana and Papua New Guinea. *The Lancet Global health.*
479 2018;6(4):e401-e10.
- 480 45. Ghinai R, El-Duah P, Chi K-H, Pillay A, Solomon AW, Bailey RL, et al. A Cross-Sectional
481 Study of 'Yaws' in Districts of Ghana Which Have Previously Undertaken Azithromycin
482 Mass Drug Administration for Trachoma Control. *PLoS Negl Trop Dis.* 2015;9(1):e0003496.
- 483 46. Mbonda Noula Aimé Gilbert, Bissek Anne C., Kamgno Joseph, Tabah Ernest N., Nneck Jan
484 René, K NA. Re-emergence of Yaws in the Centre Region, Cameroon: data from four health's
485 districts. *SSRN.* 2018.
- 486 47. Pampiglione S, Wilkinson AE. A study of yaws among pygmies in Cameroon and Zaire. *The*
487 *British Journal of Venereal Diseases.* 1975;51(3):165.
- 488 48. Tchatchouang S. Eradicating yaws in Africa: challenges and progress. *The Lancet Microbe.*
489 2022;3(4):e250.

490 49. Yotsu RR, Comoé CC, Ainyakou GT, Konan Ng, Akpa A, Yao A, et al. Impact of common
491 skin diseases on children in rural Côte d'Ivoire with leprosy and Buruli ulcer co-endemicity:
492 A mixed methods study. *PLoS Negl Trop Dis*. 2020;14(5):e0008291.

493 50. Konan DJ, Aka J, Yao KJ, Kouassi-Gohou V, Yao KE, Faye-Kette H. [Update on a
494 neglected tropical disease from the routine health information system in Côte d'Ivoire: Yaws,
495 2001 to 2011]. *Medecine et sante tropicales*. 2013;23(4):433-8.

496 51. John LN, Beiras CG, Houinei W, Medappa M, Sabok M, Kolmau R, et al. Trial of Three
497 Rounds of Mass Azithromycin Administration for Yaws Eradication. *New England Journal of*
498 *Medicine*. 2022;386(1):47-56.

499 52. Yotsu RR, Comoé CC, Ainyakou GT, Konan N, Akpa A, Yao A, et al. Impact of common
500 skin diseases on children in rural Côte d'Ivoire with leprosy and Buruli ulcer co-endemicity: A
501 mixed methods study. *PLoS Negl Trop Dis*. 2020;14(5):e0008291.

502 53. Rosenbaum BE, Klein R, Hagan PG, Seadey MY, Quarcoo NL, Hoffmann R, et al.
503 Dermatology in Ghana: a retrospective review of skin disease at the Korle Bu Teaching
504 Hospital Dermatology Clinic. *The Pan African medical journal*. 2017;26:125.

505 54. Kaburi BB, Ameme DK, Adu-Asumah G, Dadzie D, Tender EK, Addeh SV, et al. Outbreak
506 of scabies among preschool children, Accra, Ghana, 2017. 2019;19(1):746.

507 55. Amoako YA, Phillips RO, Arthur J, Abugri MA, Akowuah E, Amoako KO, et al. A scabies
508 outbreak in the North East Region of Ghana: The necessity for prompt intervention. *PLoS*
509 *Negl Trop Dis*. 2020;14(12):e0008902.

510 56. Kouotou EA, Nansseu JR, Kouawa MK, Zoung-Kanyi Bissek AC. Prevalence and drivers of
511 human scabies among children and adolescents living and studying in Cameroonian boarding
512 schools. *Parasites & vectors*. 2016;9(1):400.

513 57. Kouotou EA, Nansseu JRN, Sangare A, Moguieu Bogne LL, Sieleunou I, Adegbiidi H, et al.
514 Burden of human scabies in sub-Saharan African prisons: Evidence from the west region of
515 Cameroon. *The Australasian journal of dermatology*. 2018;59(1):e6-e10.

516 58. Saka B, Kassang P, Gnossike P, Head MG, Akakpo AS, Teclessou JN, et al. Prevalence of
517 skin Neglected Tropical Diseases and superficial fungal infections in two peri-urban schools
518 and one rural community setting in Togo. *PLoS Negl Trop Dis*. 2022;16(12):e0010697.

519 59. Hogewoning A, Amoah A, Bavinck JN, Boakye D, Yazdanbakhsh M, Adegnika A, et al.
520 Skin diseases among schoolchildren in Ghana, Gabon, and Rwanda. *International journal of*
521 *dermatology*. 2013;52(5):589-600.

- 522 60. Lopez DM, Cervantes BYH, Emmanuel D, Agordoh PD, Almaguer FM, Lambert RG, et al.
523 Infections of the Skin among Children in Ho Teaching Hospital of the Volta Region, Ghana.
524 OALib. 2020;07(03):1-9.
- 525 61. Barogui YT, Diez G, Anagonou E, Johnson RC, Gomido IC, Amoukpo H, et al. Integrated
526 approach in the control and management of skin neglected tropical diseases in Lalo, Benin.
527 PLoS Negl Trop Dis. 2018;12(6):e0006584.
- 528 62. Lompo SZ, Barogui Y, Compaore J, Ouedraogo Y, Kafando C, Kambire-Diarra MC.
529 Résultats de l'évaluation intégrée de la situation de la lèpre, de l'ulcère de Buruli et du pian
530 autour des périmètres irrigués de Bagré et Kompienga au Burkina Faso. Medecine et sante
531 tropicales. 2019;29(3):327-32.
- 532 63. WHO. Ending the neglect to attain the sustainable development goals. One Health: Approach
533 for action against neglected tropical diseases 2021-2030. Geneva: World Health Organization;
534 2022. Licence: CC BY-NC-SA 3.0 IGO..
- 535