

Trends in Efficacy, Safety, and Tolerability of Approved Multiple Sclerosis Disease Modifying Treatments

Kyle Valke, BS¹, Kangho Suh, PharmD, PhD¹

¹School of Pharmacy, University of Pittsburgh, Pittsburgh, PA

Corresponding Author:

Kangho Suh, PharmD, PhD

Assistant Professor

School of Pharmacy

University of Pittsburgh

Pittsburgh, PA 15206

kas551@pitt.edu

Telephone: 412-383-3579

Credit Author Statement: **Kyle Valke:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Visualization **Kangho Suh:** Conceptualization, Methodology, Formal analysis, Writing - Review & Editing, Visualization, Supervision

Conflict of Interests: The authors declare no relevant conflicts of interest or financial relationships.

Funding Support: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Key Words: Multiple sclerosis, relapsing-remitting multiple sclerosis, disease-modifying therapy, treatment, trends, safety, efficacy, tolerability, review

Abstract

Background: Disease-modifying therapies (DMTs) have become the mainstay of treatment for relapsing forms of multiple sclerosis (MS), reducing relapse rates and slowing disease progression.

Objectives: To determine whether or not available MS DMTs have demonstrated an increase in safety, efficacy, and tolerability over time.

Methods: Results from pivotal phase III trials of approved MS DMTs were used to create a dataset of relevant outcomes. Common endpoints analyzed include annualized relapse rates (ARR), rates of serious adverse events (SAE), and rates of discontinuation due to adverse events. Trial comparator, active or placebo, was also documented. Descriptive statistics and Fisher exact tests were performed on outcomes stratified by recency of pivotal trials.

Results: On visual inspection, there was a trend of decrease in ARR. A significant relationship was seen between recent approvals and trial design with an active comparator ($p=0.004$), as well as between recent approvals and ARR ($p=0.020$). No significance was found between recent approvals and SAE ($p=0.138$), formulation and discontinuation ($p=0.559$), or recent approvals and formulation ($p=0.352$).

Conclusion: DMTs for relapsing forms of MS increased in efficacy over time. Oral therapies offered similar tolerability to other routes of administration. Further research is warranted to identify if these clinical trial findings translate to real world evidence.

What was already known:

- The number of FDA approved disease-modifying therapies for multiple sclerosis has been steadily increasing. Available routes of administration include injectable, oral, and infusions.
- These medications are proven to be effective in reducing MS relapse rates and slowing overall disease progression, with varying degrees of safety and tolerability.
- The comparative efficacy of these therapies varies, with certain medications often deemed high efficacy.

What this study adds:

- We used published phase III trial results for each medication to provide a direct comparison between each medication's efficacy, safety, and tolerability at time of approval.
- Our analysis demonstrates a trend in increasing efficacy of available therapies along with the use of active comparators for controls in disease modifying treatments for multiple sclerosis.

Background:

Multiple sclerosis is a chronic autoimmune central nervous system (CNS) disorder characterized by neuronal demyelination. Prevalence of the disease varies by region, seen significantly more commonly in North America and Europe.¹ Disease course varies by individual, often broken down into four subtypes: the majority of patients present with a relapsing remitting course (RRMS) at disease onset, characterized by acute symptomatic exacerbations with a full or nearly complete return to function afterwards.² The most widely used diagnostic criteria for the disease is the McDonald criteria³ which combines clinical presentation and history, lesions found on MRI results, and most recently the presence of CSF-specific oligoclonal bands. Clinical presentation of the disease is complex, with a wide range of potentially debilitating symptoms manifesting particularly during acute relapses, requiring a multifaceted approach to treatment.

The mainstay of multiple sclerosis therapy has become disease-modifying therapies (DMTs) since their emergence in the 90s. These therapies exert immunomodulatory effects and have demonstrated reductions in the risk of relapses and delaying disease progression.⁴ Approved indications vary by drug however every therapy currently available is approved for treatment of relapsing-remitting multiple sclerosis. DMTs also vary by route of administration: injection, oral, or infusion. Newer therapies, particularly monoclonal antibodies which exert more direct immunomodulatory effects, are generally thought to be the higher efficacy approach to treatment.⁵

Objective:

The primary objective of this study was to summarize trends in reported efficacy and safety data for every currently approved DMT, examining whether or not newer agents have demonstrated equal or greater efficacy and improved safety and tolerability outcomes.

Methods:

PubMed was utilized to collect pivotal phase III trials for all currently FDA approved DMTs, with trials dating between 1993 through 2022.⁶⁻²⁹ Generic equivalents, novel formulations of existing medications, and monomethyl fumarate, an active metabolite of dimethyl fumarate approved based on bioavailability data, were excluded. Safety and efficacy data common amongst each trial was pulled to create a dataset. Therapies were also categorized based on route of administration: injectable, oral, or infusion. The primary efficacy endpoint that was identified and collected from every trial was annualized relapse rate (ARR). Other endpoints included in the dataset when available were percentages of patients with serious adverse events as a measure of safety and percentages of patients with adverse events leading to discontinuation as a measure of tolerability. Trial comparator, active or placebo, was also collected.

ARR was grouped by route of administration and plotted against time to visualize potential trends. Based on our objective to assess efficacy and safety in recently approved DMTs compared to older DMTs, we chose *a priori* to run several Fisher's exact tests for recently approved DMTs in the past 10 years (1993-2012 or 2013-2022) by 1) pivotal trial-reported ARR (<0.2 or >0.2), 2) annual serious adverse effect (SAE) probability (< 5% or ≥5%), 3) formulation

(oral vs injection/infusion), and 4) trial comparator, active or placebo. Additionally, we assessed whether there was a relationship between tolerability (i.e., annual discontinuation probability <5% vs ≥5% due to adverse event) and formulation. Level of significance was set at $p = 0.05$ for all tests. ARR was deemed the primary measure of overall efficacy as it was reported in all trials gathered and the primary efficacy endpoint in the vast majority. A cutoff of 0.2 was chosen based on the mean and median of reported ARRs since a point of clinical relevance was unclear. Annual probability of SAE was used as the primary measure of safety. Annual probabilities of SAE and discontinuation due to AE were calculated from respective percentages of patients from each trial. Due to limitations in available data, safety and tolerability outcomes were assessed over the last 12 years of the dataset (Table 1), resulting in cutoffs of 2009-2016 or 2017-2022.

Results:

A total of 25 pivotal phase III trials were included in the dataset. All were randomized controlled trials with the exception of the single-arm EVOLVE-MS-1 trial.²⁶ DMTs, regardless of treatment modality, showed a decline in ARR over the 30-year period demonstrating a general trend in increased efficacy (Figure 1). This was supported by a significant association between ARR less than 0.2 and more recent DMT approvals ($p=0.020$). However, for SAE categorized at above and below 5%, we did not find a significant relationship with recently approved drugs ($p=0.138$).

The other characteristics that we decided to investigate were related to mode of administration (oral vs. injection/infusion). Tolerability was assessed at a cutoff of 5% annual discontinuation probability, and there was no significant association with oral vs. injection/infusion ($p=0.559$). Additionally, while there seemed to be a shift in developing more oral formulations as a higher proportion of recently approved DMTs were oral ($n=6/11$) compared to older approvals ($n=2/8$), this was not statistically significant ($p=0.352$). We did find recently approved DMTs had a positive significant association in the use of an active comparator vs. placebo ($p=0.004$).

Discussion:

To our knowledge, this is the first study examining trends in efficacy, safety, and tolerability of MS DMTs inclusive of all currently approved therapies. Our findings support the notion that newer therapies are generally more effective. Should this trend continue MS will potentially become increasingly more manageable. Although no currently approved therapy is truly curative, the increasing potential to reduce probability of acute relapses, and by extension overall disease progression, offers hope for a future where patients with MS may experience a far lower average disease burden than historically.³⁰ The lack of a trend towards increasing safety is noteworthy; these therapies still undoubtedly have a positive risk/benefit profile for the vast majority of patients, however a greater focus on safety outcomes could be an area of focus in developing future therapies. Notably the trials referenced for our database focused specifically on RRMS, and indeed the majority of DMTs available are approved for management of RRMS and SPMS. Currently only ocrelizumab is FDA approved for PPMS and only mitoxantrone for PRMS. Other therapies may be used off-label for PPMS and PRMS, however their comparative efficacy and safety profiles are less clear in these contexts.³¹ These subtypes

warrant further study to identify optimal medication therapy management and potential novel treatments.

Our findings on tolerability by route of administration were interesting given the widespread reputations of certain therapies included. Interferons, for instance, are notorious for significant dose limiting flu-like symptoms.³² Injections and infusions also tend to have very high rates of injection site/infusion associated reactions. Notably the use of probability of discontinuation due to AEs may not fully elucidate tolerability of each drug as factors outside of AEs also weigh in. For example, injection anxiety³³ is an otherwise noted barrier to access for many individuals when choosing a prospective DMT; many patients may find frequent injections to be intolerable and avoid such medications altogether. With this in mind, the trend towards development and approval of comparably safe and effective oral therapies is altogether sensible, offering a more suitable option for many patients. Unfortunately many currently approved oral therapies also have a number of potentially dose-limiting AEs³⁴, such as gastrointestinal upset, which is likely why our results indicated no significant difference in tolerability between oral therapies and alternatives when examining impact of AEs in isolation. Managing adverse effects of DMTs and increasing tolerability overall going forwards is an important consideration given the impact adherence to DMT regimens is noted to have on improving patient outcomes and reducing costs.³⁵

The clear trend towards trial designs featuring active comparators in our dataset was to be expected given ethical considerations of withholding treatments that have increasingly become widely available, affordable, and proven effective in reducing burden of disease. Notably other studies have also acknowledged a shift towards implementation of active comparators in clinical trials.³⁶

This study has several limitations. Generally, earlier trials were more limited in terms of reported outcomes, particularly in regard to MRI data and specific safety outcomes, limiting what could be included. Selected outcomes for this study were chosen on the basis that they were reported amongst all trials included over the respective date ranges chosen. However, we selected what we believed to be most RRMS such as ARR. Trial duration also varied, ranging from 48 to 104 weeks. We adjusted for this by using annualized rates. Another limitation was that earlier trials, particularly those conducted before widespread implementation of the McDonald Criteria as a standard for diagnosis, likely enrolled participants with a higher baseline disease burden when compared to contemporary trials. In addition, certain outcome measures such as measures of disability or severity of adverse effects have an inherent degree of subjectivity by nature, further complicated once again by differences in protocols between trials in defining such outcomes. The consolidation of IM injectables and IV infusions into one subgroup for analysis may also have implications as outcomes of interest may vary between these different routes of administration. However, for the purpose of comparing oral therapies to currently available alternatives, this grouping does provide clear insight.

Conclusion:

The number of FDA approved MS DMTs has gradually increased over the past 30 years. Data from phase III trials pivotal to each drug's approval supports trends in increasing efficacy but not safety over the period. Despite increasing adoption, there was no demonstrable

1 advantage of oral therapies over injections/infusions in terms of tolerability, and there has not
2 been a significant shift towards oral therapy approvals. Studies have shifted towards inclusion of
3 an active comparator rather than a placebo over time. Reported results are limited to patients
4 with a diagnosis of RRMS; other subtypes of MS warrant further study.


References:

1. Leray E, Moreau T, Fromont A, et al. Epidemiology of multiple sclerosis. *Revue Neurologique* 2016;172(1):3-13. doi: <https://doi.org/10.1016/j.neurol.2015.10.006>
2. Rejdak K, Jackson S, Giovannoni G. Multiple sclerosis: a practical overview for clinicians. *British Medical Bulletin* 2010;95(1):79-104. doi: 10.1093/bmb/ldq017
3. Thompson AJ, Banwell BL, Barkhof F, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *The Lancet Neurology* 2018;17(2):162-73. doi: 10.1016/s1474-4422(17)30470-2
4. De Angelis F, John NA, Brownlee WJ. Disease-modifying therapies for multiple sclerosis. *Bmj* 2018;363:k4674. doi: 10.1136/bmj.k4674 [published Online First: 20181127]
5. Samjoo Imtiaz A, Worthington E, Drudge C, et al. Efficacy classification of modern therapies in multiple sclerosis. *Journal of Comparative Effectiveness Research* 2021;10(6):495-507. doi: 10.2217/ce-2020-0267
6. Calabresi PA, Kieseier BC, Arnold DL, et al. Pegylated interferon beta-1a for relapsing-remitting multiple sclerosis (ADVANCE): a randomised, phase 3, double-blind study. *The Lancet Neurology* 2014;13(7):657-65. doi: [https://doi.org/10.1016/S1474-4422\(14\)70068-7](https://doi.org/10.1016/S1474-4422(14)70068-7)
7. Calabresi PA, Radue EW, Goodin D, et al. Safety and efficacy of fingolimod in patients with relapsing-remitting multiple sclerosis (FREEDOMS II): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Neurol* 2014;13(6):545-56. doi: 10.1016/s1474-4422(14)70049-3 [published Online First: 20140328]
8. Cohen JA, Barkhof F, Comi G, et al. Oral fingolimod or intramuscular interferon for relapsing multiple sclerosis. *N Engl J Med* 2010;362(5):402-15. doi: 10.1056/NEJMoa0907839 [published Online First: 20100120]
9. Cohen JA, Coles AJ, Arnold DL, et al. Alemtuzumab versus interferon beta 1a as first-line treatment for patients with relapsing-remitting multiple sclerosis: a randomised controlled phase 3 trial. *Lancet* 2012;380(9856):1819-28. doi: 10.1016/s0140-6736(12)61769-3 [published Online First: 20121101]
10. Cohen JA, Comi G, Selmaj KW, et al. Safety and efficacy of ozanimod versus interferon beta-1a in relapsing multiple sclerosis (RADIANCE): a multicentre, randomised, 24-month, phase 3 trial. *The Lancet Neurology* 2019;18(11):1021-33. doi: 10.1016/S1474-4422(19)30238-8
11. Coles AJ, Twyman CL, Arnold DL, et al. Alemtuzumab for patients with relapsing multiple sclerosis after disease-modifying therapy: a randomised controlled phase 3 trial. *The Lancet* 2012;380(9856):1829-39. doi: [https://doi.org/10.1016/S0140-6736\(12\)61768-1](https://doi.org/10.1016/S0140-6736(12)61768-1)
12. Comi G, Kappos L, Selmaj KW, et al. Safety and efficacy of ozanimod versus interferon beta-1a in relapsing multiple sclerosis (SUNBEAM): a multicentre, randomised, minimum 12-month, phase 3 trial. *Lancet Neurol* 2019;18(11):1009-20. doi: 10.1016/s1474-4422(19)30239-x [published Online First: 20190903]
13. Ebers GC. Randomised double-blind placebo-controlled study of interferon β -1a in relapsing/remitting multiple sclerosis. *The Lancet* 1998;352(9139):1498-504. doi: 10.1016/S0140-6736(98)03334-0
14. Fox RJ, Miller DH, Phillips JT, et al. Placebo-controlled phase 3 study of oral BG-12 or glatiramer in multiple sclerosis. *N Engl J Med* 2012;367(12):1087-97. doi: 10.1056/NEJMoa1206328
15. Giovannoni G, Comi G, Cook S, et al. A Placebo-Controlled Trial of Oral Cladribine for Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2010;362(5):416-26. doi: 10.1056/nejmoa0902533

16. Gold R, Kappos L, Arnold DL, et al. Placebo-Controlled Phase 3 Study of Oral BG-12 for Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2012;367(12):1098-107. doi: 10.1056/NEJMoa1114287
17. Group TIMSS. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. *Clinical results of a multicenter, randomized, double-blind, placebo-controlled trial* 1993;43(4):655-55. doi: 10.1212/wnl.43.4.655
18. Hartung H-P, Gonsette R, König N, et al. Mitoxantrone in progressive multiple sclerosis: a placebo-controlled, double-blind, randomised, multicentre trial. *The Lancet* 2002;360(9350):2018-25. doi: [https://doi.org/10.1016/S0140-6736\(02\)12023-X](https://doi.org/10.1016/S0140-6736(02)12023-X)
19. Hauser SL, Bar-Or A, Cohen JA, et al. Ofatumumab versus Teriflunomide in Multiple Sclerosis. *N Engl J Med* 2020;383(6):546-57. doi: 10.1056/NEJMoa1917246
20. Hauser SL, Bar-Or A, Comi G, et al. Ocrelizumab versus Interferon Beta-1a in Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2017;376(3):221-34. doi: 10.1056/nejmoa1601277
21. Jacobs LD, Cookfair DL, Rudick RA, et al. Intramuscular interferon beta-1a for disease progression in relapsing multiple sclerosis. The Multiple Sclerosis Collaborative Research Group (MSCRG). *Ann Neurol* 1996;39(3):285-94. doi: 10.1002/ana.410390304
22. Johnson KP, Brooks BR, Cohen JA, et al. Extended use of glatiramer acetate (Copaxone) is well tolerated and maintains its clinical effect on multiple sclerosis relapse rate and degree of disability. Copolymer 1 Multiple Sclerosis Study Group. *Neurology* 1998;50(3):701-8. doi: 10.1212/wnl.50.3.701
23. Kappos L, Bar-Or A, Cree BAC, et al. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *The Lancet* 2018;391(10127):1263-73. doi: [https://doi.org/10.1016/S0140-6736\(18\)30475-6](https://doi.org/10.1016/S0140-6736(18)30475-6)
24. Kappos L, Fox RJ, Burcklen M, et al. Ponesimod Compared With Teriflunomide in Patients With Relapsing Multiple Sclerosis in the Active-Comparator Phase 3 OPTIMUM Study. *JAMA Neurology* 2021;78(5):558. doi: 10.1001/jamaneurol.2021.0405
25. Kappos L, Radue E-W, O'Connor P, et al. A Placebo-Controlled Trial of Oral Fingolimod in Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2010;362(5):387-401. doi: 10.1056/nejmoa0909494
26. Naismith RT, Wolinsky JS, Wundes A, et al. Diroximel fumarate (DRF) in patients with relapsing-remitting multiple sclerosis: Interim safety and efficacy results from the phase 3 EVOLVE-MS-1 study. *Multiple Sclerosis Journal* 2020;26(13):1729-39. doi: 10.1177/1352458519881761
27. O'Connor P, Wolinsky JS, Confavreux C, et al. Randomized trial of oral teriflunomide for relapsing multiple sclerosis. *N Engl J Med* 2011;365(14):1293-303. doi: 10.1056/NEJMoa1014656
28. Polman CH, O'Connor PW, Havrdova E, et al. A Randomized, Placebo-Controlled Trial of Natalizumab for Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2006;354(9):899-910. doi: 10.1056/nejmoa044397
29. Steinman L, Fox E, Hartung H-P, et al. Ublituximab versus Teriflunomide in Relapsing Multiple Sclerosis. *New England Journal of Medicine* 2022;387(8):704-14. doi: 10.1056/nejmoa2201904
30. Torkildsen Ø, Myhr K-M, Bø L. Disease-modifying treatments for multiple sclerosis – a review of approved medications. *European Journal of Neurology* 2016;23(S1):18-27. doi: <https://doi.org/10.1111/ene.12883>
31. Chedid T, Moisset X, Clavelou P. Rationale for off-label treatments use in primary progressive multiple sclerosis: A review of the literature. *Revue Neurologique* 2022;178(9):932-38. doi: <https://doi.org/10.1016/j.neurol.2022.02.461>

32. Munschauer FE, Kinkel RP. Managing side effects of interferon-beta in patients with relapsing-remitting multiple sclerosis. *Clinical Therapeutics* 1997;19(5):883-93. doi: [https://doi.org/10.1016/S0149-2918\(97\)80042-2](https://doi.org/10.1016/S0149-2918(97)80042-2)
33. Turner AP, Williams RM, Sloan AP, et al. Injection anxiety remains a long-term barrier to medication adherence in multiple sclerosis. *Rehabil Psychol* 2009;54(1):116-21. doi: 10.1037/a0014460
34. Jivraj F, Kang S, Reddie S, et al. The Patient and Clinician Assessment of Gastrointestinal (GI) Related Adverse Events Associated with Oral Disease-Modifying Therapies in Multiple Sclerosis: A Qualitative Study. *Advances in Therapy* 2022;39(11):5072-86. doi: 10.1007/s12325-022-02250-x
35. Burks J, Marshall TS, Ye X. Adherence to disease-modifying therapies and its impact on relapse, health resource utilization, and costs among patients with multiple sclerosis. *ClinicoEconomics and Outcomes Research* 2017;9:251-60. doi: 10.2147/CEOR.S130334
36. Zhang Y, Salter A, Wallström E, et al. Evolution of clinical trials in multiple sclerosis. *Therapeutic Advances in Neurological Disorders* 2019;12:1756286419826547. doi: 10.1177/1756286419826547

1 **Figure 1. Trends of annualize relapse rate by formulation type**



2
3
4
5

Table 1. Characteristics of multiple sclerosis disease modifying treatments

Brand Name	Generic Name	Approval Date	Route	Comparator(s)	ARR [†]	DC ^{†‡}	SAE ^{†‡}
Betaseron	interferon beta-1b	July 23, 1993	injectable	placebo	0.84	-	-
Copaxone	glatiramer acetate	April 20, 1996	injectable	placebo	0.65	-	-
Avonex	interferon beta-1a	May 17, 1996	injectable	placebo	0.61	-	-
Novantrone	mitoxantrone	October 29, 2000	infusion	placebo	0.35	0.041	-
Rebif	interferon beta-1a	March 29, 2002	injectable	placebo	0.88	-	-
Tysabri	natalizumab	November 23, 2004	infusion	placebo	0.23	-	-
Gilenya	fingolimod	September 21, 2010	oral	both [§]	0.18	0.061	0.066
Aubagio	teriflunomide	September 12, 2012	oral	placebo	0.37	0.051	0.074
Tecfidera	dimethyl fumarate	March 27, 2013	oral	both [¶]	0.19	0.073	0.091
Plegridy	peginterferon beta-1a	August 15, 2014	injectable	placebo	0.25	0.054	0.118
Lemtrada	alemtuzumab	November 14, 2014	infusion	active*	0.22	0.010	0.100
Ocrevus	ocrelizumab	March 28, 2017	infusion	active	0.16	0.019	0.038
Mayzent	siponimod	March 26, 2019	oral	placebo	0.07	0.047	0.123
Mavenclad	cladribine	March 29, 2019	oral	placebo	0.14	-	0.046
Vumerity	diroximel fumarate	October 30, 2019	oral	single-arm**	0.16	0.034	0.041
Zeposia	ozanimod	March 25, 2020	oral	active*	0.17	0.015	0.033
Kesimpta	ofatumumab	August 20, 2020	injectable	active [¶]	0.10	0.036	0.057
Ponvory	ponesimod	March 18, 2021	oral	active	0.20	0.042	0.042
Briumvi	ublituximab-xiiy	December 28, 2022	infusion	active [¶]	0.08	0.022	0.060

[†] Weighted averages reported for drugs with multiple trials included

[‡] Annual probabilities of serious adverse events and discontinuation due to adverse events as calculated from percentages reported in trials

[§] Three trials included in dataset, two with placebo and one with active comparator

[¶] Two trials included in dataset, one with placebo and one with active comparator

* Two trials included in dataset, both with active comparator

** Single-arm trial design, omitted from analysis

[¶] Single publication included in dataset reporting results of two parallel studies, both included in dataset and both with active comparator