

1 **Title:** Exploring athlete pain assessment experiences and priorities; A two-part qualitative series of
2 athlete and physiotherapist interactions.
3 Part Two. “Forging Our Future” - Athlete and physiotherapists’ priorities for pain assessment and
4 beyond.

5
6 **Authors**

7 Ciarán Purcell^{1,2,3,4,5,6} Caoimhe Barry Walsh³ Garrett Van Oirschot^{1,2} Brona M Fullen¹ Tomás Ward⁶
8 Brian M Caulfield^{1,2}

9 ¹*School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland*

10 ²*Insight SFI Research Centre for Data Analytics, Dublin, Ireland*

11 ³*School of Allied Health, University of Limerick, Limerick, Ireland*

12 ⁴*Physical Activity for Health Research Cluster, University of Limerick, Limerick, Ireland*

13 ⁵*Sports and Human Performance Centre, University of Limerick, Limerick, Ireland*

14 ⁶*Ageing Research Centre, University of Limerick, Limerick, Ireland*

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 **Abstract**

38 *Objectives:* To explore the priorities and directions of athlete upper and lower limb pain assessment
 39 by facilitating shared understandings of athletes and sports physiotherapists.
 40 *Design:* Qualitative Research using a hermeneutic phenomenological approach.
 41 *Methods:* We carried out focus groups using a deliberate criterion sample and a constructivist
 42 perspective. At the end of each focus group, we used the nominal group technique method to generate
 43 a list of consensus-based priorities for future pain assessment. Our paper follows the consolidated
 44 criteria for reporting qualitative research (COREQ) guidelines.
 45 *Results:* We completed five focus groups, comprising twelve athletes (female, n=5, male n=7) and
 46 four sports physiotherapists (male, n=4) Two final themes (and six subthemes) were developed; (i)
 47 Enhanced Communication and Pain Descriptions (describing and representing pain, better
 48 communication, the role of technology, providing direction and setting the pace), (ii) Integrating Sport
 49 Specific and Multidimensional Assessments (broadening the pain assessment toolkit, the role of
 50 technology). We developed a set of thirteen practical priorities for pain assessment that span the
 51 subjective, objective, and general aspects of the athlete pain assessment.
 52 *Conclusion:* We have presented stakeholder-generated perspectives, directions, and priorities for
 53 athlete pain assessment. Athletes and Physiotherapists must continue to work together to achieve a
 54 comprehensive sport-specific multidimensional pain assessment experience alongside their wider
 55 support networks to ensure optimal representation and communication. We have highlighted some
 56 available pain assessment tools and strategies and outlined how novel tools may address certain gaps
 57 in the assessment process. Researchers, clinicians, and athletes can consider the practical guidance we
 58 have provided to address these priorities.

60 **Keywords**

61 ‘Pain Measurement’, ‘Musculoskeletal Pain’, ‘Athletic Injuries’ ‘Athletic Performance’ ‘Focus
 62 Groups’, ‘Pain Management’.

64 **Introduction**

65

66 In this series, we acknowledge the value of qualitative research in sport, and we are pursuing a
 67 comprehensive understanding of athlete pain experience and assessment through qualitative methods.¹
 68 The International Association for the Study of Pain (IASP) outlines the importance of appreciating
 69 and validating each individual's experience, including both the sensory and emotional aspects of pain
 70 in their definition of pain.² The contemporary predictive processing model of pain describes how pain
 71 is not merely something we experience but something that influences how we act and interact in the
 72 world, shaping our behaviours.³ Pain neuroscience has been applied to athlete cohorts with guidelines
 73 recommending a multidimensional biopsychosocial approach to facilitate optimal pain assessment.⁴⁶
 74 Contextual pain assessment tools (affective, cognitive and socioenvironmental) have been used
 75 substantially less frequently than the more traditional tools (neurophysiological and biomechanical
 76 pain assessment) in both research and practice settings over the past fifty years and, somewhat
 77 concerningly, the gap between the use of these wider aspects and the more commonly used traditional
 78 tools is widening.⁷ In Part One of this series, athletes and physiotherapists shared their experiences of
 79 the content and qualities of pain assessments. Athletes and physiotherapists discussed the commonly
 80 used tools, measures and scales highlighting the strengths and limitations of current practice in the
 81 context of best available guidance. The quality of the pain assessment was closely linked with the pain
 82 interview, an opportunity for athletes to tell their stories, share information related to the affective,
 83 cognitive, and socioenvironmental aspects of pain and develop a strong therapeutic alliance. These
 84 findings align with recent literature emphasising the power of narrative methods to express the
 85 complexity of feelings, emotions and experiences influenced by sport-specific sociocultural aspects
 86 that athletes negotiate daily.⁸⁻¹⁰ In this paper, the second in this two-part series we present the
 87 priorities for future practice as part of an integrated overview and culmination of our overall findings.

88

89

90 **Methods**

91 *Focus Groups*

We carried out focus groups with a deliberate criterion sample of athletes and sports physiotherapists based in Ireland from diverse sporting backgrounds. We developed a topic guide which guided discussion from broad pain experience-related questions to more focused questions on priorities for pain assessment. A moderator and neutral observer helped to ensure equity of participation and full exploration of the topic guide. We used reflexive thematic analysis and developed codes, candidate themes and final themes in an iterative fashion.¹¹ A critical friend (CBW) independently reviewed the data and added additional perspectives. We have described the methodology for the focus groups component of this paper in detail in Part One of this series.¹² In the current paper, Part Two of the series, we use data from focus group questions that address priorities and directions for athlete pain assessment. The full published data set can be accessed at <https://data.mendeley.com/datasets/t47tw94mzd/2>.).

Nominal Group Technique

Nominal Group Technique (NGT) is a method of problem-solving and idea generation where all participants have an equal opportunity to contribute their priorities, effectively minimising power imbalances and facilitating the ranking of priorities suggested by participants.¹³ We used the NGT at the end of each focus group session to facilitate brainstorming of priorities and generate consensus for items to be included in the next phase of an athlete upper and lower pain assessment framework development. We chose this method to give equal weight to the opinions of all participants and to focus the ideas and topics discussed in the focus groups on priorities for pain assessment practice. NGT is a method of consensus generation where all participants contribute multiple ideas which are then voted on and prioritised equitably.¹⁴ Members rank their top five with their highest priority idea receiving five votes down to their fifth highest priority idea receiving one vote. NGT consists of four stages: Silent generation, round robin, clarification, and voting. We used five sequential steps (state the subject, reflection and writing, polling, discussion, and prioritisation) to operationalise this technique which was first developed in 1975 and has been used across domains and settings including healthcare and sport.^{15 16} We recorded and discussed all ideas generated by participants during the polling stage of the NGT. Participants voted on initial ideas which then became priorities. We

discarded all ideas that did not receive a minimum of one vote during this voting stage from our analysis. See Appendix A for a detailed description of how we applied each of the five steps in this study. We carried out the five NGTs independently and pooled the results during the data analysis stage.

Data Analysis

During the Focus Group data analysis process, we assigned athletes alphanumeric participant IDs beginning with the letter “A” and physiotherapists alphanumeric IDs beginning with “P”. Athletes and physiotherapists were talking about and experiencing the same concepts so we merged the data for coding and analysis. Participant IDs were known only to CP, the lead researcher. Once participant IDs were allocated and analysis was complete all records of participant information were deleted ensuring anonymity.

After we completed the NGTs, we reviewed all priorities that received a minimum of one vote, and I (CP) applied the initial codes. We (CP, BC, TW and GvO) reviewed the initial coding framework. We updated the codes and renamed them where necessary, grouping similar codes to generate candidate themes of assessment items and aspects. CBW reviewed the codes and candidate themes independently in her role as a critical friend and we discussed the additional perspectives. We then updated our candidate themes and developed a set of finalised themes.

Figure 1 Themes (Enhanced Communication and Pain Descriptions; Integrating Sport Specific and Multidimensional Assessments) and codes inserted here.

Figure 2 The Pain Assessment Priority Pyramid is inserted here.

144

145 **Results & Discussion**

146 The results are derived from five focus groups which gathered the experiences and interactions of
 147 sixteen participants (athletes, n=12, physiotherapists, n=4) from a broad range of sports and
 148 competition levels, a full description of which is provided in Part One of this series.¹² Athletes and
 149 physiotherapists voiced their opinions on what the future of an athlete pain assessment could and
 150 should look like. From the focus group codes, we developed two clear themes highlighting priority
 151 areas for pain assessment.

152 In Figure 1 we present these two themes and their associated codes. The first theme centres around
 153 better strategies to describe and represent pain, including tools that go beyond the current methods for
 154 capturing pain intensity and enhanced pain communication strategies. Participants highlighted aspects
 155 such as the context, timing and frequency of pain assessments and proposed technology-based
 156 solutions as one option to address current shortcomings. The challenge and opportunity of providing
 157 athletes with clear direction in the next steps of their pain assessment and management process are
 158 highlighted. The second theme explores prioritising assessment strategies for the different aspects of
 159 pain experience. Future pain assessments should be multimodal, considering a wide range of
 160 biopsychosocial and sports-specific factors to get a more comprehensive picture of the pain the athlete
 161 is experiencing. Both strategies promise to enhance current assessments offering a deeper and more
 162 nuanced understanding to better guide decision-making and management.

163 In Figure 2 we present the finalised thematic map of the athlete pain assessment priority pyramid
 164 which demonstrates how the themes and subthemes explored in this paper build on Part One of the
 165 series.

166

167

168 **Theme 1 Enhanced communication and pain descriptions.**

Theme 1 includes three sub themes; 1.1 – describing and representing pain, 1.2 – better communication and 1.3 – the role of technology.

1.1 Describing and representing pain

Participants discussed the struggle to describe and represent pain, and how it may be improved by using more comprehensive descriptions. This reflects the contemporary drive for valuing the patient’s language and selecting appropriate metaphors to help represent and explain the pain experience.¹⁷ Additionally athletes felt providing contextual information such as their pain and rehabilitation history helped explain their current pain experience. Athletes and physiotherapists highlighted the development of criteria and categories for the classification of pain intensity that go beyond the traditional numerical pain rating scale (NPRS) to include more detailed descriptions of pain and the impact of pain on aspects of daily life as a priority for pain assessment.

“I think more kinds of adjectives describing pain it definitely would be beneficial.” - A09

“Your rehab experience. Because I think they all play into how you experience pain and maybe your history of pain or your even perception of pain because everyone is an individual and they experience lots of things differently.” - A02

“It would be great to see something where it was like how does this pain influence and then there’s like several categories so maybe one would be sleep for me anyway, one would be like work you know or your sport...” – A10

There may be value in revisiting long-established measures such as the Brief Pain Inventory Scale (which includes intensity, location, body chart, interference with everyday life and pain within a certain timeframe) and the McGill Pain Questionnaire (which includes a comprehensive list to describe pain sensation, physical and emotional aspects, interaction with treatment options and wider lifestyle factors and measures of intensity that provide context to this pain experience) and developing contemporary multidimensional tools tailored to an athlete population.^{18 19}

1.2 Better communication

The timing, setting and communication of pain were earmarked by participants as areas to improve athlete pain assessments and enable some of the proposed enhanced descriptions and categorisation. Athletes and physiotherapists felt including more time for reflection, both before and after the athlete-physiotherapist assessment session, would facilitate athletes to recall their pain experiences in greater detail. Additionally, varying the time of assessments to capture pain during competition and activities of daily living that provoke pain was suggested as an opportunity to enhance pain assessment practice, particularly when athlete-physiotherapist interactions are infrequent, and athletes may struggle to accurately recall the entirety of their pain experience since their previous assessment.

“I like the idea of writing stuff down before you go in and see someone because you get a chance to actually think of it.” – P04

“I know that helped for me, because.. before I go into a physio, I do write down what I’m going to say, because more often than not I’d leave forgetting to mention things and you’re like, oh god, well it’s gone now and I’m never going to say anything.” - A11

Ecological momentary assessment (EMA) involves the repeated sampling of behaviours and experiences in real-time and is carried out in the person’s natural environment.²⁰ EMA has been applied in chronic pain and sports psychology research and our results suggest it is something which may provide value when considering improved communication in future pain assessments for athletes.^{21 22} Participants felt applying enhanced communication methods would help with difficult decision-making when it comes to athlete pain. Athletes described the challenge of not knowing when to play or compete through pain and how to effectively gauge progress whilst physiotherapists noted the opportunity for additional communication to enhance this process.

“I think as an athlete .. we want to know .. when is that pain too much to keep doing what I’m doing, when should I stop? That type of thing like what’s the recovery process?” – A06

“Sometimes as physios that’s where we could maybe have that missing link..., not necessarily to be so individualised and so 24/7 care but there has to be some level of a way of you know

220 *being able to kind of link in with them I think a bit more that I think would help sometimes.” -*

221 P02

222 **1.3 The role of technology**

223 The effective use of technology to facilitate enhanced communication and contextual assessment was
 224 proposed as a future opportunity. Technology could be used to integrate novel assessment methods
 225 such as EMA. Athletes based in high-performance environments discussed how they used
 226 smartphones to facilitate more frequent communication about pain with their sports medicine teams
 227 which they found helpful. The value of insights gleaned from regular subjective wellness reporting
 228 with athletes has been established and has been proposed to provide more value than objective
 229 measures for certain training load, health, and well-being markers.²² However, further consideration is
 230 needed to smoothly integrate technology into the contemporary pain assessment process and wider
 231 athlete ecosystem. Aspects such as validity, data protection and integration into current workflows
 232 and health system records all warrant consideration. The clinical utility of a single self-report item has
 233 been challenged in line with the need for interpretation and reasoning when it comes to
 234 musculoskeletal clinical practice standards.^{23 24} Furthermore, the future integration of technology must
 235 meet the needs of both athletes and physiotherapists, enhancing rather than complicating the pain
 236 assessment clinical encounter.

237 *“An app we use for training, and she (coach) would put in each day like how many hours did*
 238 *you sleep last night? What’s your perceived rate of recovery from the previous day’s training?*
 239 *Perceived rate of like pain that day.” – A08*

240 *“Could you fill in maybe an assessment after each training session, describing your pain and*
 241 *send that back to the physio or have regular updates throughout that process as well?” – A05*

242 *“A pain app where every morning it was what’s your pain and then a journaling section and*
 243 *it’s like how did you sleep last night? Did you take painkillers? How was your training? All*

244 *these different sections so and then say next time you see your physio potentially they would*
 245 *have access to your app and it's like okay here's with your rating over the last week" – A10*

246

247 **1.4 Providing direction and setting the pace**

248 An effective assessment sets the pace and expectations for the management process. Athletes found
 249 that receiving an accurate prognosis and timeline to build goals as a pivotal part of pain management.
 250 Athletes and Physiotherapists highlighted that the role of the physiotherapist is to provide an objective
 251 perspective to aid in the decision-making process. The optimal level of objectivity and guidance
 252 varied amongst athletes and sports environments. Regarding priorities for pain assessment practice,
 253 physiotherapists acknowledged the importance of providing clear direction at the end of the
 254 assessment.

255

256 *"I think clarity is probably the most important thing, just having a timeline said to you that*
 257 *you can believe and actually see is realistic, I think that's very important in terms of progressing" –*

258 *A05*

259 *"As much as you're a fan of the sport and a runner as well you kind of have to then be able to*
 260 *stand aside and go hold on what's the best for the athlete" – P01*

261

262 *"You're having issues with these three areas, this is what we're going to work on.." – P04*

263

264 However, both athletes and physiotherapists emphasised how diagnoses, timelines and decision-
 265 making are not always straightforward. Physiotherapists found making a definitive diagnosis and
 266 providing direction was particularly challenging in persistent pain presentations, where the signs and
 267 symptoms that frequently accompany acute pain and injury presentations were not present and
 268 recovery and improvement were often protracted requiring a greater level of pain knowledge and

clinical reasoning skills. These challenges align with gaps and future priorities in athlete pain assessment and management identified by the International Olympic Committee (IOC).²⁵

“I agree completely with what P03 was saying as regards how cloudy things become over time and I think that that’s a difficulty like for ... with a patient... I would always say to them .. look the longer we go away from your injury to where we are now, the more likely is we won’t be able to give you a definite diagnosis of what’s going on” – P02

Acknowledging these challenges in an area that does not always have a clear answer, physiotherapists prioritised understanding their limitations as an important aspect of the role of the sports physiotherapist. Additionally, they highlighted the importance of honesty and focusing on aspects within their remit to facilitate the development of the trust required for an effective athlete-physiotherapist assessment and relationship.²⁶

“Lads come in with pain and ask what is it?..I don’t know exactly what it is, it could be this, this and this, but all I know is that you’re having issues with these three areas, this is what we’re going to work on..we’ll see what it’s like.” – P04

“It’s okay to not know and I think sometimes it’s better to tell someone you don’t know but definitely have a direction you feel like they need to go in” – P04

Theme 2 Integrating sport-specific and multidimensional assessments.

Theme 2 includes two subthemes: 2.1 – broadening the pain assessment toolkit and 2.2 – the role of technology. In parallel to enhanced communication and descriptions, athletes felt multimodal and sport-specific pain assessment tools would give a more comprehensive assessment and understanding of the athlete's pain experience. This theme was developed from codes gleaned from athlete perspectives only and so it is a truly athlete-focused theme.

2.1 Broadening the pain assessment toolkit.

Psychosocial, lifestyle and environmental pain assessment tools were recommended by athletes to give a more encompassing and holistic overview of athlete pain. Athletes highlighted sport-specific pain assessment tools such as assessing movement, strength and motor control patterns relevant to their sport, asking about training load and the impact of pain on sports activities and measuring pain during, before or after sports activities to represent the unique athlete pain experience and provide enhanced contextual information in line with the best available athlete pain guidance.^{4 27} Athletes discussed how clinic-based pain assessments could be enhanced by completing more assessments in the athlete's sports environment such as on the pitch, court or track, a finding that could facilitate the practical implementation of contemporary IOC guidance.^{4 27}

“ The other environmental or the psychological factors that could have led to it because sometimes it's a very obvious thing, that reason it happened. But asking about how you feel around it and a little bit of support...getting a good grasp of someone's life and all the other stressors and factors I think definitely would help in diagnosing the thing rather than you know just going for more scans and what have you to try to pinpoint it.” – A09

“The questions that they use for pain, there's no differentiation between some guy who hurt his leg working on a building site versus somebody who's like an amateur athlete who's you know playing sport five nights a week and the questions need to be more specific to that person, that kind of activity maybe.” - A02

“More assessments directed at for the readiness to play in terms of I think even out on a pitch is completely different in the physio room .. when you're out on the pitch afterwards and you feel different types of pain .. could the physio be out and involved more?” - A05

2.2 The role of technology

Technology was once again suggested by athletes as a potential method to add multimodal and sport-specific measures into an integrative pain assessment. Additionally, the role of the physiotherapist and their expertise in pain assessment and management was stressed, and future pain assessment strategies should augment rather than replace this vital relationship. Additionally, a recent systematic review

found that there is moderate to high certainty evidence supporting the integration of technology into a physiotherapy assessment for musculoskeletal disorders in the general population.²⁸ The study found substantial to excellent validity and excellent inter and intra-rater reliability for pain (97-98% agreement) and patient-reported outcome measures (ICC 0.99-1.00, 95% CI 0.97 to 1.00). However, patients reported a superior user experience and confidence in the examination from a face-to-face assessment, supporting our findings. Additionally conducting and interpreting pain assessments with athletes requires sports and context-specific information (such as sport-specific physical demands, and coach and team expectations for return to play) that may be understood better if conducted in the athlete's sporting environment.

“So, if I could see changes in the industry its App based for sure it's probably interactive ..but there's some clear definitive I want him (the physiotherapist) to have a relationship with me, know what I'm doing and know when I can get back to my sport so there's something in that. I don't know what but...” – A06

From theory to practice: practical priorities for pain assessment.

In Table 1 we present the pooled results of the nominal group techniques we carried out. Each theme is a practical example of an aspect of assessment that can be used to operationalise the pain assessment priorities we identified through the focus group findings above. We present the codes that comprise each theme alongside a qualitative description of how clinicians can apply the assessment item in practice. We have indicated whether each item can be incorporated as part of the subjective assessment (initial interview), objective assessment (physical testing) or should be considered as a general aspect of the wider assessment process. The proportion of total votes each theme received when codes were pooled is also presented. The majority of the practical pain assessment priorities we present align closely with the published literature including 1) subjective components such as; establishing the history, context, characteristics, severity and impact of the athlete's pain and exploring the athlete's stressors, psychological aspects and lifestyle factors as well as their training

load and 2) objective components such as; assessing and identifying pain through movement, completing sports specific objective measures both at the site of pain and throughout the kinetic chain. ^{2 4 25} Developing a specific, clear, and time-appropriate assessment is a novel finding. Although the IOC guidelines recommend tailoring the pain assessment based on the presentation of the athlete and the stage in the pain management process our findings add that the assessment should align closely with the athlete's goals and the plan should be articulated clearly to the athlete so that they understand each aspect of the pain assessment and why it is being completed. Another novel finding is the need for alternative pain severity scales that go beyond the traditional numerical pain rating scale. Scales such as the traffic light pain scale which offers athletes guidance on decision-making regarding playing through pain have clinical utility and are used widely in practice.²⁹ Additionally there is scope for the inclusion of a pain severity scale that helps to overcome the limitations of a numerical scale to describe and represent the pain experience more appropriately although what that might look like is yet unclear. The International Association for the Study of Pain (IASP) identifies how a verbal description is just one method of representing pain.² Additional options include athletes using a body chart to indicate their pain or the demonstration of their pain through a specific movement or action to have their experience of pain understood and validated which is another novel finding.

Conclusion

In this paper we have presented themes focused on future pain assessment priorities and directions that build on our “athlete pain assessment” findings presented in Part One of this series. Moving beyond unidimensional, point-in-time clinical pain measures to encompass better communication and direction for pain management will involve the use of more comprehensive and descriptive pain scales and tools that athletes and physiotherapists can relate to. We have outlined examples of pain assessment tools that are currently available and should be considered by clinicians. Conversely, we have highlighted how novel tools may encompass capturing pain at critical time points in sports environments to help tell the full story of athlete pain. Whilst Physiotherapists may be justifiably reticent to introduce additional measures to avoid complicating the assessment process for athletes,

athletes are embracing existing and emerging technology as part of evolving sports science. Applying a carefully developed and well-implemented solution that incorporates available technology and keeps the athlete-physiotherapist interaction and relationship at its core is one potential future solution that must be considered.

Although we collected experiences from a diverse range of athletes and physiotherapists and variety was achieved in sport, competition level and practice setting the experiences and priorities gathered from these focus groups and nominal group techniques may not apply to all athlete pain assessment settings. Notably, participants were all recruited from Ireland and whilst some of the female athletes also had a physiotherapy background, no female sports physiotherapists were available to participate. The focus of our research was to highlight the athlete and physiotherapist voices and explore their experiences of pain assessment through open discussion. In addition, to augment these findings, we selected the Nominal Group Technique to complement the exploratory nature of the focus group methodology. We have presented a comprehensive series of pain assessment priorities for physiotherapists to consider. There is scope for future research to consider these findings in light of contemporary research and practice and develop an expert sports physiotherapist consensus-based athlete pain assessment framework that can be implemented in research and practice settings.

Practical Implications

- The pain assessment must include assessment tools and measures that allow athletes to, represent and therefore validate their specific pain experience.
- Physiotherapists should use pain assessment tools that capture the context of an athlete's pain as well as the intensity and impact on sports and activities of everyday life at key points in an athlete's day.
- Physiotherapists must consider a variety of pain assessment tools that address the multidimensional nature of pain and integrate psychological, social, environmental and sport-specific aspects of the athlete's pain experience in the pursuit of providing direction for athletes to manage their pain.

- Physiotherapists should consider the careful and effective use of technology to augment the athlete-physiotherapist pain assessment interaction.

References

1. Bekker S, Bolling C, H Ahmed O, et al. Athlete health protection: Why qualitative research matters. *Journal of Science and Medicine in Sport* 2020;23(10):898-901. doi: 10.1016/j.jsams.2020.06.020
2. Raja SN, Carr DB, Cohen M, et al. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. *Pain* 2020;161(9):1976-82. doi: 10.1097/j.pain.0000000000001939 [published Online First: 2020/07/23]
3. Kiverstein J, Kirchhoff MD, Thacker M. An Embodied Predictive Processing Theory of Pain Experience. *Review of Philosophy and Psychology* 2022 doi: 10.1007/s13164-022-00616-2
4. Hainline B, Derman W, Vernec A, et al. International Olympic Committee consensus statement on pain management in elite athletes. *Br J Sports Med* 2017;51(17):1245-58. doi: 10.1136/bjsports-2017-097884 [published Online First: 2017/08/23]
5. Caneiro JP, Alaiti RK, Fukusawa L, et al. There is more to pain than tissue damage: eight principles to guide care of acute non-traumatic pain in sport. *British Journal of Sports Medicine* 2021;55(2):75-77. doi: 10.1136/bjsports-2019-101705
6. Hoegh M, Stanton T, George S, et al. Infographic. Pain or injury? Why differentiation matters in exercise and sports medicine. *British Journal of Sports Medicine* 2022;56(5):299-300. doi: 10.1136/bjsports-2021-104633
7. Purcell C, Duignan C, Fullen BM, et al. Comprehensive assessment and classification of upper and lower limb pain in athletes: a scoping review. *British Journal of Sports Medicine* 2023;bjsports-2022-106380. doi: 10.1136/bjsports-2022-106380

- 429 8. Everard C, Wadey R, Howells K. Storying sports injury experiences of elite track athletes: A
430 narrative analysis. *Psychology of Sport and Exercise* 2021;56:102007. doi:
431 <https://doi.org/10.1016/j.psychsport.2021.102007>
- 432 9. Tabben M, Verhagen E, Warsen M, et al. Obstacles and opportunities for injury prevention in
433 professional football in Qatar: exploring the implementation reality. *BMJ Open Sport &*
434 *Exercise Medicine* 2023;9(1):e001370. doi: 10.1136/bmjsem-2022-001370
- 435 10. Costa N, Butler P, Dillon M, et al. “I felt uncertain about my whole future”—a qualitative
436 investigation of people's experiences of navigating uncertainty when seeking care for their
437 low back pain. *PAIN* 2023
- 438 11. Braun V, Clarke V. Reflecting on reflexive thematic analysis. *Qualitative Research in*
439 *Sport, Exercise and Health* 2019;11(4):589-97. doi:
440 10.1080/2159676x.2019.1628806
- 441 12. Purcell C, Barry Walsh C, Van Oirschot G, et al. Exploring athlete pain assessment experiences
442 and priorities; A two-part qualitative series of athlete and physiotherapist interactions.
443 Part One. “Gauging and discerning” - Athlete & physiotherapist pain assessment experiences
444 and interactions. *medRxiv* 2023:2023.12.28.23300487. doi: 10.1101/2023.12.28.23300487
- 445 13. McMillan SS, King M, Tully MP. How to use the nominal group and Delphi techniques.
446 *International Journal of Clinical Pharmacy* 2016 doi: 10.1007/s11096-016-0257-x
- 447 14. Delbecq ALG, David H & Van de Ven, Andrew H. Group techniques for program planning : a
448 guide to nominal group and Delphi processes / Andre L. Delbecq, Andrew H. Van de Ven,
449 David H. Gustafson. III ed. Foresman Glenview: Scott 1975.
- 450 15. US Dept of Health & Human Services, . Gaining Consensus Among Stakeholders Through the
451 Nominal Group Technique. In: Prevention DoHHSCfDCa, ed. Evaluation Briefs Centre for
452 Disease Control & Prevention 2018.
- 453 16. Dallinga J, Janssen M, van der Werf J, et al. Analysis of the Features Important for the
454 Effectiveness of Physical Activity–Related Apps for Recreational Sports: Expert Panel
455 Approach. *JMIR mHealth and uHealth* 2018;6:e143. doi: 10.2196/mhealth.9459

456 17. Louw A, Puentedura EJ, Diener I, et al. Pain neuroscience education: Which pain neuroscience
457 education metaphor worked best? *South African Journal of Physiotherapy* 2019;75(1) doi:
458 10.4102/sajp.v75i1.1329

459 18. Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain*
460 1975;1(3):277-99. doi: 10.1016/0304-3959(75)90044-5 [published Online First: 1975/09/01]

461 19. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med*
462 *Singap* 1994;23(2):129-38. [published Online First: 1994/03/01]

463 20. Pagé MG, Gauvin L, Sylvestre M-P, et al. An Ecological Momentary Assessment Study of Pain
464 Intensity Variability: Ascertaining Extent, Predictors, and Associations With Quality of Life,
465 Interference and Health Care Utilization Among Individuals Living With Chronic Low Back
466 Pain. *The Journal of Pain* 2022;23(7):1151-66. doi: 10.1016/j.jpain.2022.01.001

467 21. Reifsteck EJ, Anderson SN, Newton JD, et al. A practical guide and empirical example for
468 implementing ecological momentary assessment in sport psychology research with athletes.
469 *Sport, Exercise, and Performance Psychology* 2021;10(3):408-22. doi: 10.1037/spy0000252

470 22. Saw AE, Main LC, Gastin PB. Monitoring the athlete training response: subjective self-reported
471 measures trump commonly used objective measures: a systematic review. *Br J Sports Med*
472 2016;50(5):281-91. doi: 10.1136/bjsports-2015-094758 [published Online First: 2015/10/02]

473 23. Duignan C, Doherty C, Caulfield B, et al. Single-Item Self-Report Measures of Team-Sport
474 Athlete Wellbeing and Their Relationship With Training Load: A Systematic Review. *J Athl*
475 *Train* 2020;55(9):944-53. doi: 10.4085/1062-6050-0528.19 [published Online First:
476 2020/09/30]

477 24. Bishop A, Blackburn, J., Hallam, F., McComiskie, E., Rankin, G. Musculoskeletal physiotherapy
478 service standards; The delivery of musculoskeletal (MSK) physiotherapy services in the UK
479 for adults of 16 years and over. In: *Physiotherapy TCSO*, ed. London 2021.

480 25. Zideman DA, Derman W, Hainline B, et al. Management of Pain in Elite Athletes: Identified Gaps
481 in Knowledge and Future Research Directions. *Clin J Sport Med* 2018;28(5):485-89. doi:
482 10.1097/JSM.0000000000000618 [published Online First: 2018/06/29]

26. David S, Hitchcock J. Understanding Patient Trust in the Athletic Setting through Interviews.
Internet Journal of Allied Health Sciences and Practice 2018 doi: 10.46743/1540-580x/2018.1683
27. Hainline B, Turner JA, Caneiro JP, et al. Pain in elite athletes-neurophysiological, biomechanical and psychosocial considerations: a narrative review. *Br J Sports Med* 2017;51(17):1259-64. doi: 10.1136/bjsports-2017-097890 [published Online First: 2017/08/23]
28. Bernhardsson S, Larsson A, Bergenheim A, et al. Digital physiotherapy assessment vs conventional face-to-face physiotherapy assessment of patients with musculoskeletal disorders: A systematic review. *PLoS One* 2023;18(3):e0283013. doi: 10.1371/journal.pone.0283013 [published Online First: 2023/03/22]
29. Silbernagel KG, Thomeé R, Eriksson BI, et al. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med* 2007;35(6):897-906. doi: 10.1177/0363546506298279 [published Online First: 2007/02/20]

511

512

513

514

515

516

517

518

519

520

Theme	Codes	Description/Theme	Subjective/ Objective or General Aspect	Votes Nr	Votes %
Pain History & Context	Pain History: "What does this pain mean to you?", Pain context: subjective questions Pre-Assessment Model - Written Is pain threatening or not? Pain medication	Ask the athlete about the history of this pain presentation and prior pain experiences. Previous pain related to injuries or trauma will help establish the context. Questions around the time of the season and current home and sports environment will help to establish the "full picture". Gauge the athlete's understanding (internal context) by asking "what does this pain mean to you?" or "Do you feel this pain is threatening or not?". A pre-assessment survey could help gather this information.	Subjective	44	18.53%
Movement & Pain ID	Active ROM & pain/deficits Assessing Functional ADLs Pain provocation= movement; Palpation.	Identify the athlete's specific pain through a movement assessment which may include a full range of motion tests (noting how the athlete describes any pain to help establish the nature of the pain presentation), and assessing functional movements of daily living that provoke pain. Palpation can also be considered.	Objective	27	11.25%
Objective Assessments (Sport Specific)	Objective Sport-Specific Measures Sport-Specific Functional Assessment	Measure movement, functional patterns, and strength assessments specific to the athlete's sport. This may include movement patterns such as running, jumping, squatting and throwing as well as bilateral/unilateral strength measures.	Objective	22	9.17%
Specific clear and time appropriate assessment	Structured, tailored assessment; Prompt diagnostic imaging; Daily Monitoring/ checking Explanation of assessment process & timeline	Carry out a well-structured, goal-oriented pain assessment that is specific to the phase of the athlete's pain assessment and management process. The plan should be explained clearly to the athlete. Diagnostic imaging should be used judiciously but promptly when it is required, daily monitoring/ checkpoints may help to establish an athlete's pain recovery and wellness overtime.	General Aspect of Assessment	19	7.90%
Stressors & Lifestyle Factors	Sleep, Nutrition & Lifestyle Factors; Work & Family related stress	Fully explore the athlete's lifestyle factors. Ask the athlete about their diet, hydration and nutritional status. Ask about sleep duration, quality and the relationship between the athlete's sleep and pain. Factors such as smoking and alcohol and work/family life stressors should be explored as appropriate. Use the assessment to educate athletes regarding the impact of lifestyle factors on pain perception and experience.	Subjective	19	7.92%
Pain Severity Scales & Measures	Pain Severity & Irritability Measures & Scales Painful & Functional Pain Scale	Measure the severity and irritability of the athlete's pain. Consider the use of traditional scales such as the Numerical Pain Rating Scale, alongside more practical and descriptive scales incorporating the athlete's pain context and previous pain experiences.	Subjective/Objective	17	7.08%
Pain Characteristics	Aggravating & Easing Factors Pain Onset/Mechanism & Duration Nature/ Type of Pain	Comprehensively explore the factors that aggravate and ease the athlete's pain experience. Document the pain timeline from initial onset (including mechanism of onset where available) to the current pain presentation. Encourage athletes to describe the nature or quality of the pain with different adjectives to help determine the underlying pain mechanism.	Subjective	16	6.67%
Knowing & Understanding the athlete	Sports priorities/goals; Clinician openness re pain; Understanding individual & background; Personality Assessment	Get to know and understand the athlete as an individual, their unique background and circumstances and their sporting priorities and goals. Take an open-minded approach to pain assessment, acknowledging individual differences and communication preferences.	General aspect of assessment	15	6.25%
Athlete identifying their pain	Body Chart for ID of painful areas; show, demonstrate or describe your pain	Prioritise the accurate representation of the athlete's pain. Consider various measures to facilitate athletes to identify their pain including body charts, demonstrating and describing their pain. An athlete's ability to pinpoint their pain may indicate whether pain is localised or diffuse which can aid in establishing underlying pain mechanisms.	Objective	15	6.25%
Objective Measures (General)	Objective strength tests and measures; Kinetic Chain & Fitness Assessments Balance/Proprioception	Use a comprehensive set of objective measures to establish positive goals and identify areas of improvement to develop the athlete's robustness and performance alongside addressing their pain. These include strength (handheld dynamometry, upper and lower limb repetitions max) fitness, and balance/proprioception throughout the kinetic chain (and not just the site of pain/injury).	Objective	15	6.25%
Pain Impact	Effect of pain on QoL, ADLs & Sports performance	Explore the impact of pain on an athlete's performance within their sport as well as the influence of pain on their everyday activities. Over time the experience of pain can have an effect on an athlete's quality of life which should be discussed.	Subjective	15	6.25%
Current Training & Rehab load	Current Rehab; Training schedule & load	Determine the athlete's rehabilitation history, what worked well previously and what did not, particularly for recurrent pain episodes. Document current rehabilitation or prehabilitation activities. Establish the (acute) training load leading up to the athlete's pain presentation alongside the chronic training load preceding this episode. Record the impact of this pain episode on training load capacity.	Subjective	9	3.75%
Psychological/Emotional aspects	Post Session Psych Assessment; Discussion & support; Fear/anxiety	Explore the emotional and psychological aspects of pain. Ask the athlete about the fears, worries or concerns surrounding their pain. Take some time to explain the psychological impact of pain and injury and offer the athlete appropriate support.	Subjective	7	2.90%

521 Table 1 – Nominal Group Technique Results inserted here

522 *Votes Nr – is the number of votes each theme received. Each participant received a total of 15 votes which they*
523 *used to rank their top five priorities/codes (5 votes, 4 votes, 3 votes, 2 votes and 1 vote.) Votes % displays the*
524 *overall percentage of votes each theme received. Pain ID - Pain identification, ROM - range of motion, ADLs –*
525 *activities of daily living, QoL – Quality of life.*

526

527

528

529

530

531 **Figure Captions**

532 **Figure 1. Themes (Enhanced Communication and Pain Descriptions; Integrating Sport-specific**
 533 **and Multidimensional Assessments) and codes.** Dark shading – indicates codes that were present in
 534 athletes and physiotherapists. Light shading – indicates codes that were present in athletes only. No
 535 shading – indicates codes that were present in physiotherapists only

536

537

538 **Figure 2. Athlete pain assessment priorities themes and subthemes.** The themes for each part of
 539 this series represent a row in the priorities for pain assessment pyramid, this paper presents the top
 540 row of the pyramid, the themes and subthemes that address the priorities for athlete pain assessment.
 541 These themes build on the bottom row or foundation of the pyramid presented in Part One. Source:
 542 The cartoon element at the top of this image was designed by Freepik www.freepik.com

543

544

545

546

547

548

549

550

551

552

553


554

555


556 **Appendix A – The Five Steps of the Nominal Group Technique**

- 557 1) **State the subject.** The lead researcher wrote the title of the subject at the top of a whiteboard
558 (virtual whiteboard for Zoom session) “List all the items you think should be
559 included/prioritised in an athlete pain assessment framework.”
- 560 2) **Reflection & writing.** Ten minutes were given for silent reflection and consideration and
561 participants were asked to write down all of their ideas on separate sticky note pages to keep
562 their ideas private until the polling stage.
- 563 3) **Polling.** Each group member revealed one idea at a time, taking turns until every idea was
564 recorded by placing the sticky notes on the whiteboard. Minimal discussion took place at this
565 stage with equal contribution from each member.
- 566 4) **Discussion.** Each idea was then clarified and explained by the participant who proposed it,
567 followed by discussion and queries by the group. The specific wording of ideas was changed
568 in some circumstances following discussion upon approval from the person who came up
569 with the idea. Similar ideas were merged or grouped together at this stage.
- 570 5) **Prioritisation** Following discussion each participant ranked their top five assessment priority
571 ideas from all of the ideas generated and approved in earlier stages. A score of five was
572 allocated to the idea the participant ranked highest, a score of 4 was allocated to the idea the
573 participant ranked next highest and so on continuing to one. The ideas were then ranked based
574 on the highest to lowest scoring. Participants were then invited to share any element they felt
575 was missing or needed to be adjusted before the final ranking was approved. All ideas that
576 received a minimum score of one vote were kept.

Enhanced Communication & Pain Descriptions



Integrating Sport Specific & Multidimensional Assessments





Describing and
representing pain

Better
communication

Providing direction
and setting the
pace

Enhanced
Communication &
Pain Descriptions

Integrating Sport
Specific &
Multidimensional
Assessments

Broadening the
pain assessment
toolkit

The role of
technology

Measures, Scales &
Dimensions

Connect, Listen &
Learn

Lighthouse in the
storm

Theme	Codes	Description of Theme	Subjective, Objective or General Aspect	Votes Nr	Votes %
Pain History & Context	Pain History; What does this pain mean to you?; Pain context subjective questions; Pre Assessment Model - Written; Is pain Threatening or not?; Pain medication	Ask the athlete about the history of this pain presentation and prior pain experiences. Previous pain related to injuries or trauma will help establish the context. Questions around the time of the season and current home and sports environment will help to establish the "full picture". Gauge the athlete's understanding (internal context) by asking "What does this pain mean to you?" or "Do you feel this pain is threatening or not?" A pre-assessment survey could help gather this information.	Subjective	44	18.33%
Movement & Pain ID	Active ROM & pain/deficits; Assessing Functional ADLs; Pain provocation - movement; Palpation.	Identify the athlete's specific pain through a movement assessment which may include a full range of motion test (noting how the athlete describes any pain to help establish the nature of the pain presentation) and assessing functional movements of daily living that provoke pain. Palpation can also be considered.	Objective	27	11.25%
Objective Assessments (Sport Specific)	Objective Sport Specific Measures; Sport Specific Functional Assessment	Measure movement, functional patterns, and strength assessments specific to the athlete's sport. This may include movement patterns such as running, jumping, squatting and throwing as well as bilateral/unilateral strength measures.	Objective	22	9.17%
Specific clear and time appropriate assessment	Structured tailored assessment; Prompt diagnostic imaging; Daily Monitoring/check ins; Explanation of assessment process & timeline	Carry out a well-structured, goal-oriented pain assessment that is specific to the phase of the athlete's pain assessment and management process. The plan should be explained clearly to the athlete. Diagnostic imaging should be used judiciously but promptly when it is required, daily monitoring or check-ins may help to establish an athlete's pain, recovery and wellness over time.	General Aspect of Assessment	19	7.92%
Stressors & Lifestyle Factors	Sleep, Nutrition & Lifestyle Factors; Work & Family related stress	Fully explore the athlete's lifestyle factors. Ask the athlete about their diet, hydration and nutritional status. Ask about sleep duration, quality and the relationship between the athlete's sleep and pain. Factors such as smoking and alcohol and work/family life stressors should be explored as appropriate. Use the assessment to educate athletes regarding the impact of lifestyle factors on pain perception and experience.	Subjective	19	7.92%
Pain Severity Scales & Measures	Pain Severity & Irritability Measures & Scales; Practical & Functional Pain Scale	Measure the severity and irritability of the athlete's pain. Consider the use of traditional scales such as the Numerical Pain Rating Scale, alongside more practical and descriptive scales incorporating the athlete's pain context and previous pain experiences.	Subjective/Objective	17	7.08%
Pain Characteristics	Aggravating & Easing Factors; Pain Onset/Mechanism & Duration; Nature/ Type of Pain	Comprehensively explore the factors that aggravate and ease the athlete's pain experience. Document the pain timeline from initial onset (including mechanism of onset where available) to the current pain presentation. Encourage athletes to describe the nature or quality of the pain with different adjectives to help determine the underlying pain mechanism.	Subjective	16	6.67%
Knowing & Understanding the athlete	Sports priorities/goals; Clinician openness re pain; Understanding individual & background; Personality Assessment	Get to know and understand the athlete as an individual, their unique background and circumstances and their sporting priorities and goals. Take an open-minded approach to pain assessment acknowledging individual differences and communication preferences.	General aspect of assessment	15	6.25%
Athlete identifying their pain	Body Chart for I.D of painful areas; Show, demonstrate or describe your pain	Prioritise the accurate representation of the athlete's pain. Consider various measures to facilitate athletes to identify their pain including body charts, demonstrating and describing their pain. An athlete's ability to pinpoint their pain may indicate whether pain is localised or diffuse which can aid in establishing underlying pain mechanisms.	Objective	15	6.25%
Objective Measures (General)	Objective strength tests and measures; Kinetic Chain & Fitness Assessments; Balance/Proprioception	Use a comprehensive set of objective measures to establish positive goals and identify areas of improvement to develop the athlete's robustness and performance alongside addressing their pain. These include strength (handheld dynamometry, upper and lower limb repetitions max) fitness, and balance/proprioception throughout the kinetic chain (and not just the site of pain/injury).	Objective	15	6.25%
Pain Impact	Effect of pain on QoL, ADLs & Sports performance	Explore the impact of pain on an athlete's performance within their sport as well as the influence of pain on their everyday activities. Overtime the experience of pain can have an effect on an athlete's quality of life which should be discussed.	Subjective	15	6.25%
Current Training & Rehab Load	Current Rehab; Training schedule & load	Determine the athlete's rehabilitation history, what worked well previously and what did not, particularly for recurrent pain episodes. Document current rehabilitation or prehabilitation activities. Establish the (acute) training load leading up to the athlete's pain presentation alongside the chronic training load preceding this episode. Record the impact of this pain episode on training load capacity	Subjective	9	3.75%
Psychological/Emotional aspects	Post Session Psych Assessment, Discussion & supports; Fear or concern	Explore the emotional and psychological aspects of pain. Ask the athlete about their fear, worries or concerns surrounding their pain. Take some time to explain the psychological impact of pain and injury and offer the athlete appropriate support.	Subjective	7	2.92%

