

Microbiota-Short Chain Fatty Acid Relationships Underlie Clinical Heterogeneity and Identify Key Microbial Targets in Irritable Bowel Syndrome (IBS)

¹Andrea Shin,* ²Yue Xing, ³Mohammed Rayyan Waseem, ³Robert Siwec, ³Toyia James-Stevenson, ³Nicholas Rogers, ³Matthew Bohm, ³John Wo, ³Carolyn Lockett, ³Anita Gupta, ³Jhalka Kadariya, ⁴Evelyn Toh, ³Rachel Anderson, ⁵Huiping Xu, ²Xiang Gao*

(*co-corresponding authors)

from the ¹Vatche and Tamar Manoukian Division of Digestive Diseases at the University of California Los Angeles, Los Angeles, CA; the ²Department of Medicine at Stritch School of Medicine, Loyola University Chicago, Chicago, IL; the ³Division of Gastroenterology and Hepatology, Indiana University School of Medicine, Indianapolis, IN; the Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis, IN; and the ⁵Department of Biostatistics and Health Data Science, Indiana University School of Medicine, Indianapolis, Indiana

Short Title: Microbiota-SCFA Associations in IBS

Acknowledgements: The authors would like to thank the Mayo Clinic Department of Laboratory Medicine and Pathology and Purdue University Metabolite Profiling Facility for their collaboration and assistance with quantification of stool bile acids and short chain fatty acids.

Word Count: 5076 excluding abstract, references, table/figure legends).

Funding: AS has received funding from NIDDK K23DK122015 and R03DK132446.

Abbreviations: ALR=additive log-ratio; BAM=bile acid malabsorption; CTT=colonic transit time; gDNA=genomic DNA; GI=gastrointestinal; GLM=generalized linear model; GMM=gut

metabolic module; IBS=irritable bowel syndrome; IBS-D=IBS with diarrhea; IBS-C IBS with constipation; KEGG=Kyoto Encyclopedia of Genes and Genomics; KO=KEGG Orthogroup; LC-MS=liquid chromatography-mass spectrometry; SCFA=short chain fatty acids

Disclosures: AS serves on Ardelyx Scientific Communications Advisory Board for irritable bowel syndrome with constipation.

Author Contributions: Developing study concept: AS. Planning study design: AS, HX, XG. Participant recruitment: AS, MRW, RS, TJS, NR, MB JW, CL, AG, JK, RA. Data collection and study procedures: AS, MRW, CL, AG, JK, ET. Data Management: AS, YX, MRW, CL, JK, HX, XG. Data Analysis and Interpretation: YX, XG. Drafting the manuscript: AS, XG. Critically revising the manuscript: MRW, RS, TJS, NR, MB, JW, ET, HX.

Data Transparency Statement: Microbiome sequencing reads are deposited into NCBI with accession number SUB13882354. The data that support the findings of this study are available upon reasonable request from the corresponding author.

Correspondence:

Andrea S. Shin, M.D., M.Sc.; G. Oppenheimer Center for Neurobiology of Stress and Resilience; David Geffen School of Medicine at UCLA; 10833 Le Conte Avenue, CHS 42-210; Los Angeles, CA 90095-7378; Tel: (310) 206-0192; Fax: (310) 825-1919

Xiang Gao, PhD.; Department of Medicine; Stritch School of Medicine at Loyola University Chicago; 2160 S. Frist Avenue; Maywood, IL 60153; Tel: (708) 327-9021

ABSTRACT

Background: Identifying microbial targets in irritable bowel syndrome (IBS) and other disorders of gut-brain interaction (DGBI) is challenging due to the dynamic nature of microbiota-metabolite-host interactions. SCFA are key microbial metabolites that modulate intestinal homeostasis and may influence IBS pathophysiology. We aimed to assess microbial features associated with short chain fatty acids (SCFA) and determine if features varied across IBS subtypes and endophenotypes. Among 96 participants who were screened, 71 completed the study. We conducted in-depth investigations of stool microbial metagenomes, stool SCFA, and measurable IBS traits (stool bile acids, colonic transit, stool form) in 41 patients with IBS (IBS with constipation [IBS-C] IBS with diarrhea [IBS-D]) and 17 healthy controls. We used partial canonical correspondence analyses (pCCA), conditioned on transit, to quantify microbe-SCFA associations across clinical groups. To explore relationships between microbially-derived SCFA and IBS traits, we compared gut microbiome-encoded potential for substrate utilization across groups and within a subset of participants selected by their stool characteristics as well as stool microbiomes of patients with and without clinical bile acid malabsorption.

Results: Overall stool microbiome composition and individual taxa abundances differed between clinical groups. Microbes-SCFA associations differed across groups and revealed key taxa including *Dorea* sp. CAG:317 and *Bifidobacterium pseudocatenulatum* in IBS-D and *Akkermansia muciniphila* and *Prevotella copri* in IBS-C that that may drive subtype-specific microbially-mediated mechanisms. Strongest microbe-SCFA associations were observed in IBS-D and several SCFA-producing species surprisingly demonstrated inverse correlations with SCFA. Fewer bacterial taxa were associated with acetate to butyrate ratios in IBS compared to health. In participants selected by stool form, we demonstrated differential abundances of

microbial genes/pathways for SCFA metabolism and degradation of carbohydrates and mucin across groups. SCFA-producing taxa were reduced in IBS-D patients with BAM.

Conclusion Keystone taxa responsible for SCFA production differ according to IBS subtype and traits and the IBS microbiome is characterized by reduced functional redundancy. Differences in microbial substrate preferences are also linked to bowel functions. Focusing on taxa that drive SCFA profiles and stool form may be a rational strategy for identifying relevant microbial targets in IBS and other DGBI.

Key words: colonic microflora, bile acid metabolism, gut-brain interaction, functional gastrointestinal disorder, microbiome

INTRODUCTION

Irritable bowel syndrome (IBS) is a burdensome disorder of gut-brain interaction with an estimated global prevalence rate of 5-10%.¹ Pathophysiological mechanisms of IBS include disturbances in motility or transit, altered intestinal secretion, impaired intestinal permeability, immune cell reactivity, visceral hypersensitivity, and dysregulated neural signaling and/or central processing.² Despite advancements in the understanding of IBS pathogenesis, diagnostic and therapeutic IBS biomarkers are limited.

Accumulating evidence suggests that the gastrointestinal (GI) microbiome is associated with risk of IBS and may also mediate many of the mechanisms that underlie symptoms including altered motility,^{3,4} barrier dysfunction,⁵ immune activation,^{6,7} signaling along the brain-gut axis,⁸ and visceral sensation.⁹ Characterization of microbial composition in IBS has suggested decreased microbial diversity, reduced temporal stability, or changes in the relative abundance of specific bacteria in patients with IBS.^{10,11} However, these findings have not been sufficiently consistent across studies to establish a clear microbial profile in IBS and gaps in our understanding of the functional microbiome persist. Integrating complementary strategies in the investigation of microbial metabolites will be crucial for gathering actionable insights into the impact of the microbiome in IBS.

Atypical profiles of microbial metabolites including luminal bile acids^{12,13} and short chain fatty acids (SCFA) have been described in some patients with IBS.^{14,15} Bile acid malabsorption (BAM) is recognized as a mechanistic IBS subtype that can be assessed through several diagnostic methods including measurement of total or primary stool bile acids.¹³ Studies have demonstrated BAM to be associated with physiological traits, symptoms, and quality of life.¹⁶⁻¹⁹ Recently, researchers have examined microbial contributions to BAM in IBS to report

alterations such as enrichment of *Clostridia* bacteria including *C. scindens*,²⁰ lower microbial alpha diversity, higher Firmicutes to Bacteroidetes ratio,¹⁹ and presence of endoscopically visible biofilms correlating with overgrowth of *Escherichia coli* and *Ruminococcus gnavus*.²¹

SCFA are produced by anaerobic fermentation of dietary fibers and resistant starch that enter the colon and regulate intestinal homeostasis and physiology.²² Compared to bile acids, the role of SCFA in IBS is less well understood. Studies of stool SCFA in IBS have yielded variable results, which may be related to the heterogeneity of the IBS patient populations and multiple pathways through which SCFA may modulate intestinal physiology. Therefore, while stool SCFA are unlikely to serve as categorical IBS biomarkers, they may provide critical insights into the pathophysiological mechanisms that underlie IBS symptoms or serve as a tool for identifying metabolically relevant microbial targets. Studies^{15, 23} that have assessed stool SCFA in distinct IBS subgroups have reported more consistent associations of stool SCFA with IBS subtypes as well as correlations of stool SCFA with measurable IBS traits such as colonic transit, bowel functions, and bile acid excretion.²³⁻²⁵ Despite these reports, the intercorrelation between SCFA and transit time complicates the assessment of whether SCFA profiles represent the metabolic capacity of the resident microbiome. It remains unclear if studying the relationships between the microbiome and excreted SCFA is clinically informative. Recent work²⁶ has suggested that individual or keystone taxa, rather than complex ecological communities, could drive changes in SCFA output in response to dietary fiber. Therefore, strategies that isolate major microbial features (or keystone taxa) that shape luminal SCFA may serve as a rational method for selecting functionally relevant microbial targets in patients with IBS. To address these questions, we conducted an in-depth investigation of GI microbiome composition and function, stool SCFA,

and IBS endophenotypes defined according to quantitative traits (transit, bile acids, bowel functions) in adults with and without IBS.

MATERIALS AND METHODS

Participant Recruitment and Study Design: The study was approved by the Indiana University Institutional Review Board and the protocol registered within ClinicalTrials.gov (NCT02981888). The study was designed as an observational investigation of stool SCFA, stool bile acids, colonic transit, and stool microbiota in adults with and without IBS. We enrolled adults ages 18-65 years of age through the Indiana University Gastroenterology Clinics, the Indiana Clinical and Translational Research Institute Research Registry, and from the local community. We included individuals with IBS with diarrhea (IBS-D) or IBS with constipation (IBS-C) according to Rome IV criteria²⁷ and healthy controls with no prior history of GI diseases or symptoms. Detailed eligibility criteria are available in the Supplemental Methods.

Data Collection: Study eligibility, medications, medical history, and baseline diet using a food frequency questionnaire²⁸ were assessed during a screening visit with a study physician. Over a two-week period, data were collected on bowel functions using a standardized bowel pattern diary including the Bristol stool form scale.²⁹ All participants submitted a 48-hour stool collection collected during the last 2 days of a 4-day 100 g fat diet, consistent with clinically validated methods for identifying BAM. Specimens were refrigerated during the collection period, returned to the research team on ice, and stored at -80°C.

Colonic Transit by Radiopaque Markers: Participants underwent assessment of colonic transit time with a previously validated and optimized method using radiopaque markers.³⁰

Stool SCFA and Bile Acids: Frozen aliquots of stool were shipped to the Metabolite Profiling Facility at Purdue University to measure total and individual SCFA concentrations per mg of dry weight by liquid chromatography-mass spectrometry (LC-MS) using published methods³¹ and to the Mayo Clinic Department of Laboratory Medicine and Pathology to measure total and primary stool bile acid levels by high-performance LC-MS through a commercially available, CLIA-approved assay.^{16, 32, 33} Individual SCFA of interest included acetate, butyrate, and propionate as these represent the predominant SCFA produced in humans.³⁴

DNA Extraction, Purification, and Sequencing: Genomic DNA was isolated from stool using the QIAmp® PowerFecal® DNA kit (QIAGEN Inc., Germantown, MD, USA). DNA quality and concentration were on a Qubit fluorometer. Purified DNA underwent library preparation (Nextera XT, Illumina) and paired-end (2x150 bp) sequencing using the NovaSeq v1.5 SP (Illumina, San Diego, CA, USA) to target a sequencing depth of 40M sequences per sample.

Metagenomic Data Analysis: Metagenomic sequencing reads were quality filtered and processed for taxonomic profiling using MetaPhlAn3. Additive log-ratio (ALR) transformation was applied to analyze differential taxonomic abundances. Based on the identified microbial taxa from each sample, β -diversity indices (e.g., Bray-Curtis dissimilarity) were calculated using the R packages phyloseq and vegan.^{35, 36} Functional profiling was conducted using the HUMAnN3 pipeline annotated by KEGG (Kyoto Encyclopedia of Genes and Genomes) orthogroups (KO).

Statistical Considerations: We summarized major endpoints of interest: (1) stool microbiome composition, (2) stool SCFA concentrations (total, acetate, propionate, butyrate), total and percent primary stool bile acids, and (3) colonic transit time (overall and segmental). Our primary objective was to quantify the association between microbiome composition and SCFA

(microbe-SCFA associations) across clinical groups after controlling for transit and to account for the mechanistic heterogeneity that underlies clinical IBS populations.

We compared microbial taxa abundances between groups and analyzed associations of taxa abundances with IBS endophenotypes/traits (stool SCFA, stool bile acids, and transit) for the collective cohort and within groups. Associations of microbiome composition with group were assessed using general linear regression models (GLM) adjusted for covariates including age, sex, and BMI. Only high abundant species (relative abundance $\geq 0.1\%$) that were prevalent in ≥ 2 specimens within ≥ 1 clinical group were considered. Partial Canonical Correspondence Analysis (pCCA) conditioned on transit time was employed to quantify the relationship between the microbiome and stool SCFA in all participants and within clinical groups, using the vegan package in R. The significance of the model was assessed through a permutation test with 999 iterations. To identify bacteria whose abundances were strongly associated with the stool SCFA concentrations, we focused on those bacteria scattered along the direction of specific SCFA axis in the biplot, within a 60-degree angle centered around the SCFA axis in both positive and negative directions. The strength of association between the bacteria and the SCFA was ranked by projecting of the bacteria species scores onto each SCFA axis in the biplot. Bacteria with projection scores less than 0.5 were excluded from the subsequent analysis.

For exploratory analyses, we analyzed multivariable associations of microbiome composition with clinical BAM among individuals with IBS-D. Relative abundances were used to examine the associations of gene family/pathway abundances across clinical groups. A manually curated gut-metabolic analysis framework³⁷ was applied to examine KO identifiers associated with carbohydrate degradation, SCFA production or metabolism, and mucin degradation using GLM adjusting for covariates. As stool form³⁸ has been identified as a strong

source of human microbiota variation and closely linked to transit,^{39, 40} we further explored associations of functional potential across groups in a subset of participants who were selected based on stool form features (i.e. those individuals best representing the stool types within their respective clinical group according to bowel diary data) using the Kruskal-Wallis test. For example, we chose the IBS-D and IBS-C participants with the loosest and firmest stool types, respectively, and controls with consistently normal stool types. For all endpoints, missing values were excluded from the analysis for that endpoint.

RESULTS

Participant Characteristics: Among 96 volunteers who underwent screening evaluation, 71 completed the study, and 58 participants (Figure 1) with a mean [$\pm$ SD] age = 35.5 ($\pm$ 13.8) years and mean [$\pm$ SD] BMI = 26.2 [$\pm$ 7.5] kg/m²) were included in the final analysis after excluding those who ineligible (n=16), lost to follow-up (n=9), or and/or did not provide SCFA data (n=13). Baseline clinical characteristics (Table 1) were not significantly different across groups (all p=ns). Comparisons of quantitative traits demonstrated differences in total stool bile acids and transit between IBS-D and control participants and in total stool SCFA and stool acetate) between IBS and controls Table 2, Figures 2-3.

Stool microbiome composition differs between IBS and health and between IBS subtypes:

Shotgun metagenomic sequencing of stool samples was undertaken to obtain total of 3.1 Gb of sequence data after removal of contaminants with an average of 37.6 million paired-end reads per sample (deposited into NCBI with accession number SUB13882354). Taxonomic classification identified 461 taxa at species level classification. Comparisons of microbial metagenomes based on Bray-Curtis Dissimilarity revealed significant divergence (Supplemental Figure 1) of community distance between groups (p=0.003, R² =0.06) after adjusting for age,

BMI, and diet. We compared abundances of individual taxa across groups using the Kruskal-Wallis test to identify 18 unique and differentially abundant taxa. Highest mean relative abundances of 14 taxa (Supplemental Table 1) including *Dorea* sp. CAG:317, *Blautia* sp. CAG:257, *Ruminococcus gnavus*, and *Proteobacteria* bacterium CAG:139 were observed in IBS-D, which have previously been linked to mechanisms such as BAM^{20, 21} and serotonin biosynthesis in patients with IBS-D.⁴¹ *Lawsonibacter asaccharolyticus* abundance was highest in IBS-C and *Firmicutes* bacterium CAG:83 abundance was highest in controls.

We followed unadjusted analyses with pairwise comparisons of ALR-transformed taxa abundances using covariate-adjusted GLM, focusing on high abundant species, to demonstrate significant differences in 17 pairwise comparisons including 12 unique species (Figure 4) of which 11 exhibited differential abundances of ≥ 3 -fold. Among these taxa, we observed significantly higher abundances of *Dorea* sp. CAG:317 and *Bifidobacterium pseudocatenulatum* in IBS-D compared to IBS-C or controls and higher abundances of *Blautia* sp. CAG:257, and *Proteobacteria* bacterium CAG:139 in IBS-D compared to controls. We found significantly higher abundances of *Clostridium* sp. CAG:58 and lower abundances of *Firmicutes* bacterium CAG:83 in both IBS-D and IBS-C relative to controls, respectively. *Akkermansia muciniphila* and *Prevotella copri* were increased in IBS-C compared to controls. Differences of ≥ 3 -fold in relative abundances among high abundant bacteria were also observed in 151 pairwise comparisons (Supplemental Table 2) but were not statistically significant after adjustment for covariates.

Microbe-SCFA associations differ between IBS subtypes and controls: We conducted pCCA on microbiome and SCFA data, conditioned on transit time, to quantify relationships between taxa abundances and stool SCFA in the overall cohort and within clinical groups. In the overall

cohort, we observed a significant correlation between microbiome abundance matrix and SCFA concentration matrix ($p=0.033$, $R^2 = 0.095$). Relationships between microbiome abundances and SCFA concentrations within clinical groups (Supplemental Figure 2) were more pronounced within IBS-D ($p=0.015$) and less pronounced in HV and IBS-C ($p=ns$). In the overall cohort, strongest positive associations with total SCFA were observed with *Bacteroides plebeius*, *Prevotella* sp. CAG:1031, and *Bifidobacterium pseudocatenulatum*.

Distinct taxa correlate with individual SCFA across clinical subgroups: Keeping in mind that SCFA effects may depend on SCFA type, we ranked bacterial taxa according to their projection scores for individual SCFAs to assess the pattern of associations between bacterial species and SCFA profiles. Bacteria with higher projection scores for individual SCFAs were considered to have stronger associations with those specific SCFAs. Bacteria with projection scores less than 0.5 were excluded from the subsequent analysis. In general, the strength and direction of correlations of individual taxa with acetate, butyrate, and propionate differed according to clinical group (Figure 5). No microbes demonstrated consistent patterns of strong correlations with individual SCFA across all three groups. The greatest number of bacteria with strong associations with SCFA were observed in IBS-D, as evidenced by the larger number of bacteria remaining in the ranked list after filtering. Most microbe-SCFA patterns differed between IBS-D and IBS-C and only a few bacterial species including *Ruminococcus torques*, *Coprococcus comes*, *Clostridium* sp. CAG:299, *Bacteroides eggerthii*, and *Adlercreutzia equolifaciens* demonstrated consistent associations across both IBS subtypes with acetate and butyrate.

In IBS-D, both positive and negative correlations with stool acetate were observed with multiple taxa. Largest negative projections were observed with several bacteria that have previously linked to SCFA production (*B. plebeius*, *Roseburia hominis*),^{42, 43} dietary

polysaccharides and protein utilizers (*Firmicutes* bacterium CAG:110),⁴⁴ starch or fructooligosaccharide degradation (*Bifidobacterium adolescentis*), fiber fermentation (e.g. *Clostridium leptum*⁴⁵), β -fructan utilizers (e.g. *Roseburia inulinivorans*).⁴⁵ Positive projections were observed with *Dialister* sp. CAG 357, *B. pseudocatenulatum*, *Alistipes* species, *Prevotella* sp. CAG:1031, *R. toques*, and *R. gnavus*. In IBS-C, largest negative projections for acetate were observed between abundances of *Dorea* sp. CAG:317, *Lachnospira pectinoschiza*, and *Prevotella* sp. CAG:1031 while the largest positive projections were observed with *Streptococcus salivarius*, *Bacteroides fragilis*, and *R. torques*. In controls, largest negative projections for acetate were observed with *B. plebeius*, *Dialister* sp. CAG:357, *Bacteroides dorei*, and *A. muciniphila*. The largest positive projection for acetate among controls was observed with *Ruminococcus lactaris*.

Analysis of microbe-butyrate associations in IBS-D demonstrated negative projection scores for butyrate with several bacteria including *Ruminococcaceae* bacterium D5, *Firmicutes* bacterium CAG:110, *C. leptum*, *R. hominis*, *R. inulinivorans*, and *L. asaccharolyticus*. The largest positive projection scores for butyrate were observed with *Alistipes* species, *Prevotella* sp. CAG:1031, and *R. torques*. In IBS-C and controls, patterns of microbe-butyrate associations largely mirrored the patterns of microbe-acetate associations observed within the respective clinical groups. In addition, a positive relationship between butyrate and *Bacteroides thetaiotaomicron*, a well-known human commensal^{46, 47} that exhibits the capacity to digest a broad array of polysaccharides and host glycans, was also observed in IBS-C.

For propionate, most microbes including *Ruminococcaceae* bacterium D5, *Firmicutes* bacterium CAG:110, *C. leptum*, *L. asaccharolyticus*, *Ruthenibacterium lactatiformans* and *Bacteroides massiliensis* exhibited negative associations in IBS-D with only a few exceptions

including *S. salivarius*, for which the largest positive correlation was observed. In contrast, both positive and negative projections for stool propionate were observed in IBS-C; the largest positive scores were observed for *Paraprevotella xylaniphila*, *Phascolarctobacterium succinatutens*, and *P. copri*. Among controls, both positive and negative associations with propionate were observed for multiple bacterial taxa.

Lower number of bacterial species associated with acetate to butyrate ratios in IBS: Since interconversion of SCFA from acetate to butyrate may represent a major route of butyrate formation within the gastrointestinal tract,⁴⁸ we examined bacterial relationships with acetate to butyrate ratios to identify top taxa associated the metabolic processes that drive SCFA output. The largest positive associations were observed with *Prevotella* sp. 885, *Firmicutes* bacterium CAG:83, *P. xylaniphila* in IBS-D, suggesting that these species were associated with shift toward decreased butyrate production. Unlike the patterns observed with individual SCFA, analysis acetate to butyrate ratios demonstrated that the number of ranked bacteria were substantially lower in IBS-D or IBS-C compared to controls (Figure 5), suggesting the possibility of reduced functional redundancy for butyrate formation in IBS.

Reduced SCFA-producing microbes and bile acid malabsorption (BAM): Stool microbial metagenomes were analyzed in 22 IBS-D patients with (n=8) and without (n=14) clinical BAM according to previously validated diagnostic cutoff values.^{12, 13} Comparisons of beta diversity based on Euclidian dissimilarity of ALR transformed abundance data showed significant dissimilarity ($p = 0.039$; $R^2 = 0.061$) between patients with and without BAM (Supplemental Figure 3). Significant differences in six high abundant species were observed between groups, including *L. asaccharolyticus* *R. inulinivorans*, *L. pectinoschiza*, and *F. saccharivorans* which

were also associated with stool SCFA, but not bile acids, in IBS-D. These four species with known SCFA-producing capacity were negatively correlated with BAM.

Functional potential for carbohydrate degradation, SCFA metabolism, and mucin degradation:

Examination of microbial gene families and pathways yielded 949,223 total gene families within the metagenome data set including 4,042 named KO terms. We applied a gut metabolic module (GMM) framework to analyze modules related to carbohydrate degradation, SCFA production or metabolism including bacterial cross-feeding pathways, mucin degradation, and one module related to serine degradation based on significant species-based predictors. Abundances of KO identifiers within modules for lactose degradation, ($p=0.0026$), serine degradation ($p=0.019$), and propionate production ($p=0.026$) were significantly enriched in IBS-C compared to controls. Relative abundance of a KO identifier within a galactose degradation module was significantly reduced in IBS-D ($p=0.033$) compared to controls. In the collective cohort, we applied the GMM framework to analyze the relationships between metagenomically-encoded functions and stool SCFA. We identified 23, 23, 15, and 16 KO identifiers from relevant GMMs that were significantly associated with total SCFA, acetate, butyrate, and propionate, respectively. Within a representative subset ($n=6$ controls, $n=5$ IBS-D, $n=5$ IBS-C) selected by stool form characteristics, functional metagenomic analyses demonstrated that microbial genes/pathways associated with SCFA production/metabolism and degradation of carbohydrates/ mucin were differentially associated with clinical group (Figure 6).

DISCUSSION

It has been proposed that changes in intestinal SCFAs in some patients with IBS are caused by shifts in microbial composition and function to drive and maintain symptoms. We hypothesized that specific features of the GI microbiome may correlate with SCFA output and

that these relationships may distinguish IBS subtypes and endophenotypes. We further hypothesized that functional features of the GI microbiome defined by substrate preferences and SCFA metabolism may be linked to bowel dysfunction. Using a dual-omics approach, our findings demonstrate microbiota-SCFA relationships differ between IBS subtypes and endophenotypes, and between patients with IBS and healthy controls. We elected to focus on microbe-SCFA relationship due to the established biological relevance of SCFA in GI physiology and to use a focused strategy for identifying the microbes involved in shaping the microbial metabolome. Our findings suggest that key taxa may predict IBS mechanisms and that the functional repertoire of the intestinal microbiome influences bowel functions. By capturing quantifiable traits, we also gather evidence suggesting the microbiome has direct effects on excreted SCFA and were able to identify microbe-SCFA relationships that are not explained by transit, to potentially isolate major microbial features (or keystone taxa) that shape SCFA output.

While numerous studies have reported compositional changes¹¹ in IBS such as decreased microbial diversity and altered abundances of individual taxa, consistent patterns are difficult to pinpoint and limit our ability to draw actionable insights. Recent data suggest that putative microbial biomarkers are related to mechanistic and host-specific features in IBS rather than the overall clinical syndrome. Consistent with these prior studies, we observed differences in overall microbial community composition across groups. Comparisons of taxa abundances demonstrated significant changes were most common in IBS-D and predominantly characterized by higher abundances of several bacterial species including *Blautia* sp. CAG:257, *R. gnavus*, *Dorea* sp. CAG:317, and *Proteobacteria* bacterium CAG:139. *R. gnavus*, a Lachnospiraceae, has previously been identified to play a pathogenic role in IBS-D through serotonin biosynthesis,⁴¹ biofilm formation and increased stool bile acid excretion,²¹ production of proinflammatory

polysaccharides,⁴⁹ and mucin degradation.⁵⁰ Interestingly, the relationship between *R. gnavus* and IBS lost its significance in adjusted analyses due to a positive association between *R. gnavus* and BMI which may suggest a role *R. gnavus* in explaining the connection between IBS-D or chronic diarrhea with metabolic syndrome and obesity-related disorders although further validation of this hypothesis will be required.⁵¹ Associations of IBS-D with other taxa including *Dorea* sp. CAG:317, *Blautia* sp. CAG:257, and *Proteobacteria* bacterium CAG:139 remained and a new association between *B. pseudocatenulatum* and IBS-D emerged. In our analyses of microbe-SCFA relationships, strongest contributions to SCFA in the overall cohort were observed with *Bacteroides plebeius*, *Prevotella* sp. CAG:1031, and *B. pseudocatenulatum* suggesting these taxa as major drivers of microbial metabolism which is consistent with their known functions. *B. plebeius* harbors a specialized polysaccharide utilization locus acquired through horizontal gene transfer from marine bacteria.^{52, 53} *Prevotella* spp. are highly prevalent within the human gut, and is largely regarded as a fiber degrader that is associated with agrarian or plant-based diets, although strain-level variations may impact their contributions to health and disease.⁵⁴ *B. pseudocatenulatum* is recognized as an adult-type *Bifidobacterium* that is metabolically adapted toward dietary carbohydrates including long-chain xylans.⁵⁵ Among the three clinical groups, we found microbe-SCFA associations were most pronounced in IBS-D and that fewer microbes were linked to acetate and butyrate ratios. These observations may indicate that microbial contributions to SCFA profiles are particularly important in IBS-D and may also imply a greater degree of microbial specialization and reduced functional redundancy. Previously, Jacobs et al.⁵⁶ reported significant increases in transcript abundances for fructooligosaccharide and polyol utilization and upregulation of transcripts for fructose and glucan metabolism as well as the succinate pathway for carbohydrate fermentation in IBS.

Comparisons between IBS-D and IBS-C demonstrated differences in microbial metabolism with upregulation of multiple metabolic pathways including transcripts for fructose, mannose, and polyol metabolism in IBS-D. Together, our findings suggest that differences microbial metabolism between IBS subtypes may underlie variations in clinical symptoms or that changes in microbial metabolism could play a particularly prominent role in IBS-D. These findings could explain the higher prevalence of mixed- and diarrhea-predominant symptoms in post-infection IBS^{57, 58} as well as the limited evidence for efficacy of antimicrobial treatments such as rifaximin in IBS-C⁵⁹ relative to IBS-D.^{60, 61}

Dorea sp. CAG:317 and *B. pseudocatenulatum* were identified as top ranked bacteria associated with acetate to butyrate ratio and acetate concentrations. Mucin degradation has been described among several *Dorea* species and the genus *Dorea* belongs to the Lachnospiraceae family, a major producer of SCFA,⁶² Notably, Wang et al.⁶³ recently demonstrated that while xylan supplementation following fiber deprivation alleviated gut dysbiosis through selective promotion of *B. pseudocatenulatum*, these changes were found to be associated with lower community diversity. *Dorea* sp. CAG:317 and *B. pseudocatenulatum* may represent keystone taxa linked to altered microbial metabolism or persistent impairment in community recovery among patients with IBS-D. Only two bacterial species, *A. muciniphila* and *P. copri* were significantly increased in IBS-C. Although relationships between microbes and SCFA were less clear in IBS-C. *P. copri* was further identified as top ranked bacteria that was positively associated with propionate and negatively associated with acetate to butyrate ratio in IBS-C. Previous studies have reported increased *A. muciniphila* in IBS-C to be protective against experimentally-induced colitis in mice⁶⁴ and a positive association between mucosal *P. copri* abundance and abdominal pain in patients with IBS without prior history of infection.⁶⁵ Others

have demonstrated that IBS patients with high levels of propionate present with worse gastrointestinal symptoms, quality of life, and negative emotions.¹⁵ *P. copri* is a prominent gut commensal that contributes to propionate production via the succinate pathway⁶⁶ and has been linked to both beneficial and harmful effects on human health. Rolhion et al. found *P. copri* to be associated with mucus erosion and increased propionate. In another study by Jiang et. al, *P. copri* colonization was associated with proinflammatory effects following a high fiber diet in patients with rheumatoid arthritis through succinate production.⁶⁷ Exacerbation of rheumatoid with the high fiber diet in the presence of *P. copri* colonization led to microbial dysbiosis characterized by increased abundance of *Akkermansia*, which was also increased in our own IBS-C cohort.

In addition to identify major microbial features associated with individual SCFA, we made several important observations related to the pattern or direction of observed microbe-SCFA relationships. Interestingly, negative correlations were observed in the IBS-D subgroup between several SCFA-producing species including *R. hominis*, *B. adolescentis*, *R. inulinivorans* and individual SCFA. *B. adolescentis* utilizes fructo-oligosaccharides and starch and leads to the production of lactate and acetate which is cross-fed to butyrate-producing anaerobes,⁶⁸ but have also been identified as a potential GABA producer and correlated with anxiety and/or depression,⁶⁹ which are common comorbid conditions in patients with IBS. *R. inulinivorans* is a known fiber fermenter⁴⁵ that also possesses the ability to forage mucin through a metabolic interplay with other members of the intestinal microbiota⁷⁰ and is also known for net acetate uptake,⁶⁶ which may explain its correlation with reduced SCFA. In general, analysis of bacterial associations with individual SCFAs demonstrated the largest number of negative associations in IBS-D. Many of these negative correlations involved bacterial species that have been implicated in SCFA production and polysaccharide utilization. Findings may imply disrupted cross-feeding

mechanisms, modifications in substrate utilization, competitive inhibition, or other context-specific changes leading to altered microbial metabolism in IBS. Upon analyzing relationships of bacterial species with acetate to butyrate ratios, which we studied as a marker for overall SCFA production and intestinal SCFA availability,⁷¹ we identified fewer top ranked bacteria in IBS compared to healthy controls to suggest a reduction in overall functional redundancy within the IBS microbiome. Future work should investigate the effects of bacterial cross-feeding, microbial conversions, and interactions of the colonic microbiome with diet- and host-derived substrates that may destabilize the microbiota-mucosal interface to expose vulnerabilities and promote pathophysiological mechanisms that drive IBS.

A few bacterial species were associated with SCFA in both IBS subtypes including *R. torques*, *C. comes*, *Clostridium* sp. CAG:299, *B. eggerthii*, and *Adlercreutzia equolifaciens* which may suggest these bacterial species represent shared microbial features across IBS subtypes. *R. torques* is a mucin-degrading bacteria that has previously been associated with IBS⁷² as well as depression.⁷³ Similarly, recent studies have demonstrated associations between *C. comes*⁷⁴ and *B. eggerthii*⁷⁵ depression and PTSD, which have both been described as comorbid conditions in IBS. *R. A. equolifaciens* has been described as an anti-inflammatory commensal in non-alcoholic fatty liver disease.⁷⁶ These data could point towards shared microbial features as novel biomarker of common pathogenic mechanisms in IBS involving the microbiota-gut-brain axis, possibly through the effects of microbially-derived SCFA.

In our comparisons across IBS-D patients by with and without BAM, we observed BAM to be associated with reduced abundances of several SCFA-producing bacterial species. Similar observations have been made in studies linking biofilm formation to bile acid accumulation and reduced SCFA-producing bacteria.²¹ Results suggest that BAM exerts indirect effects on the

intestinal metabolome or the overall metabolic potential of the colonic microbiota characterized by a decline in SCFA-producing capacity. We also assessed of functional variations in the GI microbiome to reveal differential abundances of KO identifiers belonging to metabolic processes related to degradation of carbohydrates and serine as well as SCFA production. In a subset of participants selected by stool form characteristics, we identified changes in substrate degradation potential and fermentative capacity between IBS and controls and between IBS subtypes that were characterized by increased capacity for starch degradation in controls, enhancement of distinct mucin-degrading functions in IBS-D vs. IBS-C, and reduced pectin degrading potential in IBS-C. Results suggest that abnormal stool form in IBS is influenced by microbiota-encoded substrate preferences.

Collectively, our findings indicate there are distinct changes in microbiome composition may underlie clinical heterogeneity in IBS. Using a dual-omics approach, we demonstrate that the gut microbiota in IBS-D is characterized by increased metabolic activity, reduced functional redundancy, and persistently reduced diversity. Several taxa including *R. gnavus*, *Dorea* sp. CAG:317 and *B. pseudocatenulatum* are identified as the leading candidates for further study as potential microbial targets in IBS-D. A few unique microbial features including *P. copri*-associated propionate production and *Akkermansia* expansion may play a crucial role in IBS-C. Meanwhile, microbial features that are common between subtypes may offer insights into shared pathophysiological mechanisms involving the microbiota-gut-brain axis in IBS. Identifying the major microbial features (taxa or functions) that drive metabolic output may be crucial in defining rational approaches to modulate the microbiome in IBS.

Despite study strengths, we recognize the limitations of this work including cross-sectional sampling, which overlooks the fluctuating nature of the microbiome. To address this,

we assessed relationships between microbial abundances and quantifiable IBS endophenotypes that were defined at the time of specimen collection while accounting for baseline covariates. All participants were instructed to consume a 4-day high fat diet while otherwise maintaining their usual diet. Previously, we examined the effect of both habitual diet and real-time intake on excreted stool SCFA to find that that modest variations in macronutrient intake including polysaccharides did not exert substantial effects on excreted SCFA.^{25, 77} We enrolled patients with opposing IBS phenotypes from both the university clinics and the surrounding communities to increase our ability to detect differences between clinical cohorts and limit referral bias. We also acknowledge that causal microbial mechanisms cannot be determined through the current work. However, we applied a dual-omics approach to investigate biologically relevant changes in intestinal microbiome composition and function. Finally, we cannot discount the possibility we may have missed significant taxa due to the moderate sample size and a risk for Type I errors as we did not adjust for false discovery rate in our analysis of taxonomic differences. However, our analytic approach was based on endpoints that were determined *a priori* and we applied a conservative strategy by focusing on taxa that were associated with SCFA while accounting for transit. We further limited analyses to high abundant taxa and to species that exhibited >3-fold change in abundance while focusing on the biological plausibility of our results. Findings from this study should be validated in larger, longitudinal cohorts, but provide a framework that may be used to define key microbial interactions and targets in IBS.

In conclusion, main findings from this study highlight microbiota-SCFA patterns vary across clinical IBS phenotypes and endophenotypes. Our observations suggest that a paired microbe-metabolite approach may represent a viable strategy for identifying actionable microbial features in heterogenous patient populations. Prominent shifts in microbial composition are

observed in IBS-D. These changes appear to affect metabolic capacity, substrate preferences, and metabolite profiles. In addition, altered microbial substrate uptake may impact stool form or bowel function in IBS. These findings generate new hypotheses regarding candidate microbes that could be pursued as actionable microbial biomarkers or metabolically influential features in the effort to develop rational microbiota-based therapies in IBS.

References:

1. Sperber AD, Bangdiwala SI, Drossman DA, et al. Worldwide Prevalence and Burden of Functional Gastrointestinal Disorders, Results of Rome Foundation Global Study. *Gastroenterology* 2021;160:99-114 e3.
2. Ford AC, Sperber AD, Corsetti M, Camilleri M. Irritable bowel syndrome. *Lancet* 2020;396:1675-1688.
3. Tap J, Derrien M, Tornblom H, et al. Identification of an Intestinal Microbiota Signature Associated With Severity of Irritable Bowel Syndrome. *Gastroenterology* 2017;152:111-123 e8.
4. Anitha M, Vijay-Kumar M, Sitaraman SV, et al. Gut microbial products regulate murine gastrointestinal motility via Toll-like receptor 4 signaling. *Gastroenterology* 2012;143:1006-16 e4.
5. Heitkemper MM, Cain KC, Shulman RJ, et al. Stool and urine trefoil factor 3 levels: associations with symptoms, intestinal permeability, and microbial diversity in irritable bowel syndrome. *Benef Microbes* 2018;9:345-355.
6. Sundin J, Rangel I, Repsilber D, Brummer RJ. Cytokine Response after Stimulation with Key Commensal Bacteria Differ in Post-Infectious Irritable Bowel Syndrome (PI-IBS) Patients Compared to Healthy Controls. *PLoS One* 2015;10:e0134836.
7. McCracken VJ, Lorenz RG. The gastrointestinal ecosystem: a precarious alliance among epithelium, immunity and microbiota. *Cell Microbiol* 2001;3:1-11.
8. Liu Y, Zhang L, Wang X, et al. Similar Fecal Microbiota Signatures in Patients With Diarrhea-Predominant Irritable Bowel Syndrome and Patients With Depression. *Clin Gastroenterol Hepatol* 2016;14:1602-1611 e5.

- 513 9. Crouzet L, Gaultier E, Del'Homme C, et al. The hypersensitivity to colonic distension of
514 IBS patients can be transferred to rats through their fecal microbiota. *Neurogastroenterol*
515 *Motil* 2013;25:e272-82.
- 516 10. Mars RAT, Frith M, Kashyap PC. Functional Gastrointestinal Disorders and the
517 Microbiome-What Is the Best Strategy for Moving Microbiome-based Therapies for
518 Functional Gastrointestinal Disorders into the Clinic? *Gastroenterology* 2021;160:538-
519 555.
- 520 11. Pittayanon R, Lau JT, Yuan Y, et al. Gut Microbiota in Patients With Irritable Bowel
521 Syndrome-A Systematic Review. *Gastroenterology* 2019;157:97-108.
- 522 12. Shin A, Camilleri M, Vijayvargiya P, et al. Bowel functions, fecal unconjugated primary
523 and secondary bile acids, and colonic transit in patients with irritable bowel syndrome.
524 *Clin Gastroenterol Hepatol* 2013;11:1270-1275 e1.
- 525 13. Vijayvargiya P, Camilleri M, Burton D, et al. Bile and fat excretion are biomarkers of
526 clinically significant diarrhoea and constipation in irritable bowel syndrome. *Aliment*
527 *Pharmacol Ther* 2019;49:744-758.
- 528 14. Mars RAT, Yang Y, Ward T, et al. Longitudinal Multi-omics Reveals Subset-Specific
529 Mechanisms Underlying Irritable Bowel Syndrome. *Cell* 2020;182:1460-1473 e17.
- 530 15. Tana C, Umesaki Y, Imaoka A, et al. Altered profiles of intestinal microbiota and organic
531 acids may be the origin of symptoms in irritable bowel syndrome. *Neurogastroenterol*
532 *Motil* 2010;22:512-9, e114-5.
- 533 16. Vijayvargiya P, Camilleri M, Chedid V, et al. Analysis of Fecal Primary Bile Acids
534 Detects Increased Stool Weight and Colonic Transit in Patients With Chronic Functional
535 Diarrhea. *Clin Gastroenterol Hepatol* 2019;17:922-929 e2.

- 536 17. BouSaba J, Sannaa W, McKinzie S, et al. Impact of Bile Acid Diarrhea in Patients With
537 Diarrhea-Predominant Irritable Bowel Syndrome on Symptoms and Quality of Life. Clin
538 Gastroenterol Hepatol 2022;20:2083-2090 e1.
- 539 18. BouSaba J, Zheng T, Dilmaghani S, et al. Effect of rapid colonic transit on stool
540 microbiome and short-chain fatty acids in diarrhoea-predominant irritable bowel
541 syndrome. Gut 2023.
- 542 19. Camilleri M, Carlson P, BouSaba J, et al. Comparison of biochemical, microbial and
543 mucosal mRNA expression in bile acid diarrhoea and irritable bowel syndrome with
544 diarrhoea. Gut 2023;72:54-65.
- 545 20. Zhao L, Yang W, Chen Y, et al. A Clostridia-rich microbiota enhances bile acid excretion
546 in diarrhea-predominant irritable bowel syndrome. J Clin Invest 2020;130:438-450.
- 547 21. Baumgartner M, Lang M, Holley H, et al. Mucosal Biofilms Are an Endoscopic Feature
548 of Irritable Bowel Syndrome and Ulcerative Colitis. Gastroenterology 2021;161:1245-
549 1256 e20.
- 550 22. Silva YP, Bernardi A, Frozza RL. The Role of Short-Chain Fatty Acids From Gut
551 Microbiota in Gut-Brain Communication. Front Endocrinol (Lausanne) 2020;11:25.
- 552 23. Ringel-Kulka T, Choi CH, Tamas D, et al. Altered Colonic Bacterial Fermentation as a
553 Potential Pathophysiological Factor in Irritable Bowel Syndrome. Am J Gastroenterol
554 2015;110:1339-46.
- 555 24. Gargari G, Taverniti V, Gardana C, et al. Fecal Clostridiales distribution and short-chain
556 fatty acids reflect bowel habits in irritable bowel syndrome. Environ Microbiol
557 2018;20:3201-3213.

- 558 25. Waseem MR, Shin A, Siwiec R, et al. Associations of Fecal Short Chain Fatty Acids
559 With Colonic Transit, Fecal Bile Acid, and Food Intake in Irritable Bowel Syndrome.
560 Clin Transl Gastroenterol 2023;14:e00541.
- 561 26. Deehan EC, Yang C, Perez-Munoz ME, et al. Precision Microbiome Modulation with
562 Discrete Dietary Fiber Structures Directs Short-Chain Fatty Acid Production. Cell Host
563 Microbe 2020;27:389-404 e6.
- 564 27. Mearin F, Lacy BE, Chang L, et al. Bowel Disorders. Gastroenterology 2016.
- 565 28. Mulligan AA, Luben RN, Bhaniani A, et al. A new tool for converting food frequency
566 questionnaire data into nutrient and food group values: FETA research methods and
567 availability. BMJ Open 2014;4:e004503.
- 568 29. Heaton KW, Radvan J, Cripps H, et al. Defecation frequency and timing, and stool form
569 in the general population: a prospective study. Gut 1992;33:818-24.
- 570 30. Sadik R, Abrahamsson H, Ung KA, Stotzer PO. Accelerated regional bowel transit and
571 overweight shown in idiopathic bile acid malabsorption. Am J Gastroenterol
572 2004;99:711-8.
- 573 31. Park J, Kim M, Kang SG, et al. Short-chain fatty acids induce both effector and
574 regulatory T cells by suppression of histone deacetylases and regulation of the mTOR-
575 S6K pathway. Mucosal Immunol 2015;8:80-93.
- 576 32. Tagliacozzi D, Mozzi AF, Casetta B, et al. Quantitative analysis of bile acids in human
577 plasma by liquid chromatography-electrospray tandem mass spectrometry: a simple and
578 rapid one-step method. Clin Chem Lab Med 2003;41:1633-41.
- 579 33. Vijayvargiya P, Camilleri M, Shin A, Saenger A. Methods for diagnosis of bile acid
580 malabsorption in clinical practice. Clin Gastroenterol Hepatol 2013;11:1232-9.

- 581 34. Louis P, Hold GL, Flint HJ. The gut microbiota, bacterial metabolites and colorectal
582 cancer. *Nat Rev Microbiol* 2014;12:661-72.
- 583 35. McMurdie PJ, Holmes S. Phyloseq: a bioconductor package for handling and analysis of
584 high-throughput phylogenetic sequence data. *Pac Symp Biocomput* 2012:235-46.
- 585 36. Oksanen J, Simpson GL, Blanchet F, et al. *vegan: Community Ecology Package*.
586 Ordination methods, diversity analysis and other functions for community and vegetation
587 ecologists. 2.6-4 ed, 2022.
- 588 37. Vieira-Silva S, Falony G, Darzi Y, et al. Species-function relationships shape ecological
589 properties of the human gut microbiome. *Nat Microbiol* 2016;1:16088.
- 590 38. Vujkovic-Cvijin I, Sklar J, Jiang L, et al. Host variables confound gut microbiota studies
591 of human disease. *Nature* 2020;587:448-454.
- 592 39. Saad RJ, Rao SS, Koch KL, et al. Do stool form and frequency correlate with whole-gut
593 and colonic transit? Results from a multicenter study in constipated individuals and
594 healthy controls. *Am J Gastroenterol* 2010;105:403-11.
- 595 40. Jaruvongvanich V, Patcharatrakul T, Gonlachanvit S. Prediction of Delayed Colonic
596 Transit Using Bristol Stool Form and Stool Frequency in Eastern Constipated Patients: A
597 Difference From the West. *J Neurogastroenterol Motil* 2017;23:561-568.
- 598 41. Zhai L, Huang C, Ning Z, et al. *Ruminococcus gnavus* plays a pathogenic role in
599 diarrhea-predominant irritable bowel syndrome by increasing serotonin biosynthesis. *Cell*
600 *Host Microbe* 2023;31:33-44 e5.
- 601 42. Yun EJ, Yu S, Park NJ, et al. Metabolic and enzymatic elucidation of cooperative
602 degradation of red seaweed agarose by two human gut bacteria. *Sci Rep* 2021;11:13955.

43. Machiels K, Joossens M, Sabino J, et al. A decrease of the butyrate-producing species *Roseburia hominis* and *Faecalibacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut* 2014;63:1275-83.
44. Quan J, Wu Z, Ye Y, et al. Metagenomic Characterization of Intestinal Regions in Pigs With Contrasting Feed Efficiency. *Front Microbiol* 2020;11:32.
45. Armstrong HK, Bording-Jorgensen M, Santer DM, et al. Unfermented beta-fructan Fibers Fuel Inflammation in Select Inflammatory Bowel Disease Patients. *Gastroenterology* 2023;164:228-240.
46. Martens EC, Chiang HC, Gordon JI. Mucosal glycan foraging enhances fitness and transmission of a saccharolytic human gut bacterial symbiont. *Cell Host Microbe* 2008;4:447-57.
47. Porter NT, Luis AS, Martens EC. *Bacteroides thetaiotaomicron*. *Trends Microbiol* 2018;26:966-967.
48. Shetty SA, Boeren S, Bui TPN, et al. Unravelling lactate-acetate and sugar conversion into butyrate by intestinal *Anaerobutyricum* and *Anaerostipes* species by comparative proteogenomics. *Environ Microbiol* 2020;22:4863-4875.
49. Henke MT, Kenny DJ, Cassilly CD, et al. *Ruminococcus gnavus*, a member of the human gut microbiome associated with Crohn's disease, produces an inflammatory polysaccharide. *Proc Natl Acad Sci U S A* 2019;116:12672-12677.
50. Bell A, Brunt J, Crost E, et al. Elucidation of a sialic acid metabolism pathway in mucus-foraging *Ruminococcus gnavus* unravels mechanisms of bacterial adaptation to the gut. *Nat Microbiol* 2019;4:2393-2404.

- 625 51. Shin A, Xu H, Imperiale TF. Associations of chronic diarrhoea with non-alcoholic fatty
626 liver disease and obesity-related disorders among US adults. *BMJ Open Gastroenterol*
627 2019;6:e000322.
- 628 52. Hehemann JH, Correc G, Barbeyron T, et al. Transfer of carbohydrate-active enzymes
629 from marine bacteria to Japanese gut microbiota. *Nature* 2010;464:908-12.
- 630 53. Hehemann JH, Kelly AG, Pudlo NA, et al. Bacteria of the human gut microbiome
631 catabolize red seaweed glycans with carbohydrate-active enzyme updates from extrinsic
632 microbes. *Proc Natl Acad Sci U S A* 2012;109:19786-91.
- 633 54. De Filippis F, Pasolli E, Tett A, et al. Distinct Genetic and Functional Traits of Human
634 Intestinal *Prevotella copri* Strains Are Associated with Different Habitual Diets. *Cell*
635 *Host Microbe* 2019;25:444-453 e3.
- 636 55. Derrien M, Turroni F, Ventura M, van Sinderen D. Insights into endogenous
637 *Bifidobacterium* species in the human gut microbiota during adulthood. *Trends Microbiol*
638 2022;30:940-947.
- 639 56. Jacobs JP, Lagishetty V, Hauer MC, et al. Multi-omics profiles of the intestinal
640 microbiome in irritable bowel syndrome and its bowel habit subtypes. *Microbiome*
641 2023;11:5.
- 642 57. Klem F, Wadhwa A, Prokop LJ, et al. Prevalence, Risk Factors, and Outcomes of
643 Irritable Bowel Syndrome After Infectious Enteritis: A Systematic Review and Meta-
644 analysis. *Gastroenterology* 2017;152:1042-1054 e1.
- 645 58. Breen-Lyles M, Decuir M, Byale A, et al. Impact of Rome IV criteria on the prevalence
646 of post-infection irritable bowel syndrome. *Neurogastroenterol Motil* 2023;35:e14532.

- 647 59. Pimentel M, Chang C, Chua KS, et al. Antibiotic treatment of constipation-predominant
648 irritable bowel syndrome. *Dig Dis Sci* 2014;59:1278-85.
- 649 60. Pimentel M, Lembo A, Chey WD, et al. Rifaximin therapy for patients with irritable
650 bowel syndrome without constipation. *N Engl J Med* 2011;364:22-32.
- 651 61. Lembo A, Sultan S, Chang L, et al. AGA Clinical Practice Guideline on the
652 Pharmacological Management of Irritable Bowel Syndrome With Diarrhea.
653 *Gastroenterology* 2022;163:137-151.
- 654 62. Vacca M, Celano G, Calabrese FM, et al. The Controversial Role of Human Gut
655 Lachnospiraceae. *Microorganisms* 2020;8.
- 656 63. Wang Z, Bai Y, Pi Y, et al. Xylan alleviates dietary fiber deprivation-induced dysbiosis
657 by selectively promoting *Bifidobacterium pseudocatenulatum* in pigs. *Microbiome*
658 2021;9:227.
- 659 64. Gobert AP, Sagrestani G, Delmas E, et al. The human intestinal microbiota of
660 constipated-predominant irritable bowel syndrome patients exhibits anti-inflammatory
661 properties. *Sci Rep* 2016;6:39399.
- 662 65. Choo C, Mahurkar-Joshi S, Dong TS, et al. Colonic mucosal microbiota is associated
663 with bowel habit subtype and abdominal pain in patients with irritable bowel syndrome.
664 *Am J Physiol Gastrointest Liver Physiol* 2022;323:G134-G143.
- 665 66. Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic
666 microbiota. *Environ Microbiol* 2017;19:29-41.
- 667 67. Jiang L, Shang M, Yu S, et al. A high-fiber diet synergizes with *Prevotella copri* and
668 exacerbates rheumatoid arthritis. *Cell Mol Immunol* 2022;19:1414-1424.

68. Belenguer A, Duncan SH, Calder AG, et al. Two routes of metabolic cross-feeding between *Bifidobacterium adolescentis* and butyrate-producing anaerobes from the human gut. *Appl Environ Microbiol* 2006;72:3593-9.
69. Duranti S, Ruiz L, Lugli GA, et al. *Bifidobacterium adolescentis* as a key member of the human gut microbiota in the production of GABA. *Sci Rep* 2020;10:14112.
70. Pichler MJ, Yamada C, Shuoker B, et al. Butyrate producing colonic Clostridiales metabolise human milk oligosaccharides and cross feed on mucin via conserved pathways. *Nat Commun* 2020;11:3285.
71. Wijdeveld M, Schranter A, Hagemeijer A, et al. Intestinal acetate and butyrate availability is associated with glucose metabolism in healthy individuals. *iScience* 2023;26:108478.
72. Carco C, Young W, Gearry RB, et al. Increasing Evidence That Irritable Bowel Syndrome and Functional Gastrointestinal Disorders Have a Microbial Pathogenesis. *Front Cell Infect Microbiol* 2020;10:468.
73. Mulder D, Jakobi B, Shi Y, et al. Gut microbiota composition links to variation in functional domains across psychiatric disorders. *Brain Behav Immun* 2024;120:275-287.
74. Yu L, Chen X, Bai X, et al. Microbiota Alters and Its Correlation with Molecular Regulation Underlying Depression in PCOS Patients. *Mol Neurobiol* 2023.
75. Xiao L, Liu S, Wu Y, et al. The interactions between host genome and gut microbiome increase the risk of psychiatric disorders: Mendelian randomization and biological annotation. *Brain Behav Immun* 2023;113:389-400.

- 690 76. Onate FP, Chamignon C, Burz SD, et al. Adlercreutzia equolifaciens Is an Anti-
691 Inflammatory Commensal Bacterium with Decreased Abundance in Gut Microbiota of
692 Patients with Metabolic Liver Disease. Int J Mol Sci 2023;24.
- 693 77. Calderon G, Patel C, Camilleri M, et al. Associations of Habitual Dietary Intake With
694 Fecal Short-Chain Fatty Acids and Bowel Functions in Irritable Bowel Syndrome. J Clin
695 Gastroenterol 2022;56:234-242.

696

697

698

699

Table 1: Clinical Characteristics of Patients with Irritable Bowel Syndrome (IBS) and Controls

<i>Data show mean (standard deviation [SD]) except where specified</i>	Controls (n=17)	IBS-C (n=15)	IBS- D (n=26)
Age (years)	31.7 (13.6)	34.3 (10.5)	38.7 (15.3)
Women, n (%)	12 (71%)	13 (87%)	18 (70%)
BMI, kg/m²	26.3 (5.9)	25.3 (5.2)	26.7 (9.4)
White, n (%)	9 (53%)	10 (67%)	23 (88%)
Baseline dietary intake*			
Energy (kcal)	1946.8 (1025.7)	2028.4 (1717.3)	1514.1 (557.4)
Total carbohydrates (g)	219.4 (155.6)	191.1 (153.7)	160.1 (62.9)
Starch (g)	104.7 (54.9)	107.9 (110.9)	87.7 (46.9)
Protein (g)	100.9 (47.1)	96.8 (75.3)	75.1 (23.4)
Fat (g)	77.6 (34.7)	99.6 (101.9)	63.8 (30.3)
Fiber (g)	17.9 (20.9)	14.3 (10.0)	11.6 (4.1)

Group comparisons were conducted using the ANOVA F- and Fisher exact tests. IBS-C, IBS with constipation; IBS-D, IBS with diarrhea; *Diet data missing for two participants with IBS-C and four participants with IBS-D

Table 2: Quantitative Traits in Patients with Irritable Bowel Syndrome (IBS) and Controls

<i>Data show median (interquartile range)</i>	Controls (n=17)	IBS-C (n=15)	IBS- D (n=26)
Total stool bile acids (μmol/48h)*#	342 (130-640)	190 (110-449)	607 (447-1235)
Primary stool bile acids (%)	2.1 (0.8-9.1)	1.8 (0.8-2.4)	3.1 (0.8-12.4)
Total stool SCFA (μg/mg) @**	9.8 (4.5-13.8)	11.6 (7.7-20.0)	14.3 (10.0-20.8)
Stool acetate@*	6.3 (3.3-8.5)	7.1 (5.3-12.8)	9.4 (5.9-14.1)
Stool propionate	1.7 (0.8-2.4)	2.2 (1.5-2.9)	2.4 (1.6-4.0)
Stool butyrate	1.2 (0.5-2.4)	2.2 (1.0-3.7)	1.9 (1.0-3.6)
Acetate to butyrate ratio	5.0 (3.3-6.7)	3.7 (2.7-4.9)	4.5 (3.3-6.7)
Total colonic transit time (CTT), days	1.4 (0.2-2.5)	1.5 (1.0-2.1)	0.9 (0.5-1.5)
Right CTT, days	0.6 (0.3-0.6)	0.4 (0.2-0.8)	0.4 (0.2-0.6)
Transverse CTT, days	0.1 (0-0.4)	0.3 (0-0.8)	0.1 (0-0.3)
Left CTT, days**#	0.5 (0.3-1.9)	0.5 (0.2-1.1)	0.3 (0.1-0.5)

IBS-C=IBS with constipation; IBS-D=IBS with diarrhea; SCFA=short chain fatty acids. Comparisons across groups were conducted using the Kruskal-Wallis test. For traits showing significant differences (* $p < 0.05$; ** $p = 0.05$), pairwise comparisons were conducted with Dunn's test, applying Bonferroni correction for multiple tests. # $p < 0.05$ for Controls vs. IBS-D; \$ $p = 0.001$ for IBS-C vs. IBS-D; all other pairwise comparisons were non-significant (e.g., IBS-C vs. controls); @* $p < 0.05$ for Controls vs. IBS. @** $p = 0.05$ for Controls vs. IBS

FIGURE LEGENDS

Figure 1: Consort Flow Diagram

Figure 2: Stool Short Chain Fatty Acid Concentrations in Healthy Volunteers and Participants

with Irritable Bowel Syndrome (IBS). *Histograms are shown for total short chain fatty acids,*

acetate, butyrate, propionate, and acetate to butyrate ratios within clinical groups including

healthy volunteers (HV), IBS with constipation (IBS-C), and IBS with diarrhea (IBS-D). Groups

are denoted by color (HV = red, IBS-C = green, IBS-D = blue).

Figure 2: Stool Bile Acids in Healthy Volunteers and Participants with Irritable Bowel

Syndrome (IBS). *Histograms are shown for % primary bile acids (bile.acids) and total bile acids*

within clinical groups including healthy volunteers (HV), IBS with constipation (IBS-C), and IBS

with diarrhea (IBS-D). Groups are denoted by color (HV = red, IBS-C = green, IBS-D = blue).

Figure 4: Bacterial Taxa with Differential Abundance of ≥ 3 -fold Between Groups. *Data show*

mean percent abundances (standard deviation) of significantly different bacterial taxa with ≥ 3 -

fold differences in pairwise comparisons of clinical cohorts.

Figures 5: Correlations of bacterial taxa with short chain fatty acids (SCFA) based on partial

canonical correspondence analysis. *Panels show associations between bacterial taxa and specific*

SCFA([a] acetate, [b] butyrate, [c] propionate, and [d] and acetate to butyrate ratio) within

cohorts. The strength of the correlation is represented projections >0.5 with purple indicating

negative correlations and red indicating positive correlations.

Figure 6: Variations in Microbial Substrate Utilization in Healthy Volunteers and Participants

with IBS. *Figure shows significant differences in functional potential for carbohydrate*

degradation, short chain fatty acid production or metabolism, and mucin degradation across

735 *clinical groups in representative subset selected by stool form characteristics. Groups are*
736 *denoted by color (HV = blue, IBS-C = red, IBS-D = gray).*

737

738

Assessed for eligibility (n=96)


Excluded due to screen fail (n=16)

Selected and enrolled (n=80)


Lost to follow-up or missing
sequencing data (n=24)

Study follow-up


Final data analysis (n=58)




Histograms with Density Curves by Cohort




Histograms with Density Curves by Cohort




Histograms with Density Curves by Cohort




Histograms with Density Curves by Cohort




Histograms with Density Curves by Cohort



Histograms with Density Curves by Cohort



Histograms with Density Curves by Cohort



Mean percent abundances (standard deviation) of significantly different high abundant bacterial taxa in pairwise comparisons

Taxa	Controls	IBS-C	IBS-D	IBS-C vs. Controls (p-value)	IBS-D vs. Controls (p-value)	IBS-C vs. IBS-D (p-value)
<i>Akkermansia muciniphila</i>	0.294 (±1.022)	1.890 (±3.138)	1.241 (±2.676)	0.029	ns	ns
<i>Bifidobacterium pseudocatenulatum</i>	0.064 (±0.167)	0.041 (±0.154)	1.510 (±6.706)	ns	0.034	0.006
<i>Blautia</i> sp. CAG:257*	0.016 (±0.034)	0.043 (±0.122)	0.121 (±0.220)	ns	0.03	ns
<i>Clostridium</i> sp. CAG:58*	0.013 (±0.022)	0.174 (±0.201)	0.253 (±0.546)	0.004	0.025	ns
<i>Dorea</i> sp. CAG:317*	0.008 (±0.025)	0.006 (±0.025)	0.132 (±0.306)	ns	0.03	0.011
<i>Firmicutes</i> bacterium CAG:110	0.273 (±0.519)	0.030 (±0.096)	0.037 (±0.110)	0.047	ns	ns
<i>Firmicutes</i> bacterium CAG:83*	1.046 (±3.048)	0.135 (±0.185)	0.551 (±1.845)	ns	0.002	0.038
<i>Monoglobus pectinilyticus</i> **	0.164 (±0.540)	0.198 (±0.303)	0.081 (±0.374)	0.028	ns	0.008
<i>Prevotella copri</i>	2.037 (±5.027)	8.401 (±19.932)	3.986 (±12.817)	0.049	ns	ns
<i>Proteobacteria</i> bacterium CAG:139*	0.020 (±0.036)	0.364 (±0.810)	0.635 (±1.512)	ns	0.008	ns
<i>Ruminococcus torques</i>	3.299 (±5.879)	0.755 (±1.570)	0.971 (±1.683)	ns	0.044	ns
<i>Streptococcus salivarius</i>	0.092 (±0.230)	0.121 (±0.177)	0.200 (±0.343)	0.021	ns	ns

*bacterial species that were significantly different in overall comparisons; ** differential increase in *M. pectinilyticus* abundance in IBS-C vs. controls or IBS-D <3-fold; color scheme represents relative abundance with higher values noted in darker blue

