

Title: Non-Consensual Sex among Japanese Women in the COVID-19 Pandemic: A Large-Scale Nationwide Survey-Based Study

Authors:

Tomoya Suzuki,¹ Yudai Kaneda,² Yasuhiro Kotera,³ Akihiko Ozaki,^{1,4} Tetsuya Tanimoto,^{1,5} Divya Bhandari,¹ Sayaka Horiuchi,⁶ Takahiro Tabuchi⁷

Affiliations:

1. Medical Governance Research Institute, Minato, Tokyo, Japan
2. School of Medicine, Hokkaido University, Sapporo, Hokkaido, Japan
3. Faculty of Medicine and Health Sciences, University of Nottingham, Nottingham, UK
4. Department of Breast and Thyroid Surgery, Jyoban Hospital of Tokiwa Foundation, Iwaki, Fukushima, Japan
5. Department of Internal Medicine, Navitas Clinic, Tachikawa, Tokyo, Japan
6. Graduate Faculty of Interdisciplinary Research Faculty of Medicine Basic Medicine (Epidemiology/Environmental Medicine), University of Yamanashi
7. Cancer Control Center, Osaka International Cancer Institute, Osaka, Japan

Corresponding author:

Tomoya Suzuki

Medical Governance Research Institute, Minato, Tokyo, Japan

Tel: +81-80-9873-8361

E-mail: tomo1502@gmail.com

Acknowledgments:

None

Authors' Contributions

Conceptualization: TS, AO, and TakahiroT

Data analysis: TS and AO

Writing original draft: TS and AO

Funding acquisition: AO and TakahiroT

Project Administration: AO and TakahiroT

Review and editing: YudaiK, TetsuyaT, YasuhiroK, BD, SH, and TakahiroT

All authors read and approved the final manuscript for publication.

Funding Information

This study was funded by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grants (grant number 17H03589; 19K10671; 19K10446; 18H03107; 18H03062; 21H04856], the JSPS Grant-in-Aid for Young Scientists (grant number 19K19439], Research Support Program to Apply the Wisdom of the University to tackle COVID-19-Related Emergency Problems, University of Tsukuba and Health Labor Sciences Research Grant (grant number 19FA1005; 19FG2001].

Conflict of interest statement:

TT reports personal fees from Medical Network Systems, MNES Inc., and Bionics co. ltd., outside the submitted work. AO reports personal fees from Medical Network Systems, MNES Inc., outside the submitted work. All remaining authors have nothing to disclose.

Keywords: COVID-19, Non-consensual sex, sex consent, sexual abuse, sexual violence, rape, sexual education

ABSTRACT

Background: Non-consensual sex including rape and sexual assault has been a global concern and may have been influenced by the COVID-19 pandemic, however the information on this topic is limited. Therefore, our objective was to survey the incidence rate of non-consensual sex among Japanese women aged 15-79 years between April to September 2020, following the COVID-19 pandemic in Japan.

Materials and Methods: We utilized the data obtained from a nationwide, cross-sectional internet survey conducted in Japan between August and September 2020. Sampling weights were applied to calculate national estimates, and multivariable logistic regression was performed to identify factors associated with non-consensual sex. Data was extracted from a cross-sectional, web-based, self-administered survey of approximately 2.2 million individuals from the general public, including in men and women.

Results: Excluding men and responses with inconsistencies, the final analysis included 12,809 women participants, with 138 (1.1%) reporting experiencing non-consensual sex within a five-month period. Being aged 15–29 years and having a worsened mental or economic status were associated with experiencing non-consensual sex.

71

72 **Conclusions:** Early intervention to prevent individuals from becoming victims of sexual harm
73 should be extended to economically vulnerable and young women, especially during times of
74 societal upheaval such as the COVID-19 pandemic. Additionally, Japan should prioritize the
75 implementation of comprehensive education on the concept of sexual consent.

76

INTRODUCTION

Non-consensual sex refers to engaging in sexual behavior without obtaining consent to touch, observe, or perform sexual acts involving private body parts that exceed the boundaries of another person's body or mind (1,2). Non-consensual sex can take many forms, including rape, transactional sex, cross-generational sex, unwanted touch, and molestation. In the European Union, between 45% to 55% of women have experienced sexual harassment since the age of 15 (3). Previous research has identified common factors associated with victims, including ages between 10 and 30 years, mental vulnerability, and financial hardship (4). Women with these characteristics are at a higher risk of experiencing non-consensual sex compared to those without these traits (4). Furthermore, women with these characteristics often find it challenging to disclose incidents of non-consensual sex to others, making it difficult to address these issues (5).

Of note, non-consensual sex is reported to increase due to disruptions in social conditions, such as natural disasters and worsened economic conditions. For instance, incidents of non-consensual sex against women dramatically increased during times of economic and psychological turmoil, such as the 2005 Hurricane Katrina in the United States and the 2011 Great East Japan Earthquake (6–8). Therefore, it is crucial to investigate the prevalence of non-

consensual sex after the COVID-19 pandemic.

At the onset of the COVID-19 pandemic, it was reported that 243 million girls and women aged 15-49 years had been subjected to non-consensual sex worldwide (9). This was attributed to the increase in the mandatory stay-at-home measure due to lockdowns, and resulting in reduced access to social support from teachers, friends and caregivers (10). Awareness of the status of non-consensual sex during the COVID-19 pandemic has grown in many countries (11,12). However, the frequency and characteristics of non-consensual sex remain to be elucidated, not only after but also before the COVID-19 pandemic in Japan to date, partly due to traditional social norms that discourage open discussions about sexual issues (13).

The purpose of our study was to report the incidence of non-consensual during the early stages of the COVID-19 pandemic in Japan. Additionally, we explored potential risk factors such as mental health conditions, economic changes, and concerns about the lockdown, and anxiety about the disease.

METHODS

Context

This cross-sectional study was conducted among a sample of approximately 2.2 million

individuals from the general public who participated in a web-based self-reported questionnaire survey for the Japan COVID-19 and Society Internet Survey (JACSIS) project (14). The survey was administered by the well-established internet research agency, Rakuten Insight, Inc., which has been utilized in prior research studies (14,15). Data were collected from April 2020 to September 30, 2020. The prefectures where the selected individuals lived represented all prefectures in Japan, as indicated in previous studies (16).

Patient and public involvement

No patients were involved in this study.

Participants

The survey invitation was extended to 28,000 individuals out of 2.2 million registered with the survey agency, Rakuten Insight, Inc. Using a computer algorithm, study participants were selected through random sampling.

Analysis Subjects

Respondents who self-identified as women in the questionnaire were included in our analysis. Among all the survey respondents totaling 28,000, the number of valid women participants for

analysis was 12,809.

Outcome Variable: Experience of Non-Consensual Sex

The focus of this study was to examine instances of non-consensual sex occurring between from April 2020 to 30 September 2020. Participants were not questioned about specific details but were instead asked to respond with a simple “yes” or “no” to indicate whether they had experienced such incidents.

Exposure Variables

Sociodemographic characteristics

As independent variables, demographics were included, encompassing gender, age groups (15–19, 20–29, 30–39, 40–49, 50–59, 60–69, 70–79 years), marital status (unmarried, married and widowed/separated), having children (none or more than one), educational attainment (high school educated, college educated or higher), employment status (any type of employment, unemployed), household income level, which was calculated as dividing the household income by the square root of household size (categorized by the tertials of household equivalent income (low, <JPY2.5 million/US\$25 000/£16 667; intermediate, JPY2.5–JPY4.3 million/US\$25 000–US\$43 000/£16 667–£28 667; high, >JPY4.3

million/<US\$43 000/<£28 667; unknown/declined to answer)), and smoking status (not smoker or current smoker).(14)

Living Area Classification

Regarding living regions, all 47 prefectures were divided into the following three areas based on the date of the Declaration of the State of Emergency (DSE area) (17,18). “Designated as DSE on April 7, 2020”, “Designated as DSE on April 16, 2020”, and “Others”, “Specific alert designated on April 7, 2020” included Tokyo, Kanagawa, Chiba, Saitama, Osaka, Hyogo, and Fukuoka. “Specific alert designated on April 16, 2020” included Hokkaido, Ibaraki, Ishikawa, Gifu, Aichi, and Kyoto. “Others” comprised the remaining 34 prefectures. These categorization were determined by the Japanese Government using three indicators: the cumulative number of infected people, epidemiological trend, medical capacity and surveillance system (17). During the target period, people living in the relevant prefectures were advised to avoid unnecessarily outings, limit the use of entertainment facilities, and adhere to restrictions on economic activities (19).

Fear of COVID-19 Scale

The Japanese version Fear of COVID-19 Scale (FCV-19S) was used to assess participants’

fear of the COVID-19 infection (20). This instrument comprises seven items rated on a 5-point scale (1 = “strongly disagree”; 5 = “strongly agree”). Consequently, the total score ranged from 7 to 35, with a higher score indicating a greater fear of COVID-19. Satisfactory internal consistency and validation of the scale were confirmed in the original seven-item scale ($\alpha = .82$) (21).

Personal Health and Economic Statuses

Based on previous research, an examination was conducted to determine whether health and economic statuses were associated with non-consensual sex (4). The health-related questions included the following: Self-rated health (good or other than good), “Any change in mental state in the last month compared to before January 2020 ?” (Worse, No change, Getting better, I do not know)”, Suicidal thoughts (Happened since before COVID-19 pandemic, First experienced during the COVID-19 pandemic or Never), Desire not to talk to anyone due to worries (Yes or No), Feeling isolated (Yes or No), Any cancellation of a family gathering due to the COVID-19 pandemic (Yes or No), Non-payment of salary (Happened since before COVID-19 pandemic, First happened during the COVID-19 pandemic, Never), and Lack of money to buy necessities of life (Happened since before COVID-19 pandemic, First Happened during the COVID-19 pandemic or Never).

187

188 **Statistical Procedure**

189 To achieve the purpose of identifying the incidence of non-consensual sexual activity and its
190 associated factors during the COVID-19 outbreak, two types of analysis methods were
191 employed.

192

193 First, to determine the incidence rate, we divided the sample into two groups: those who
194 responded “Yes” to the question of “Any non-consensual sex” and who did not. The number
195 and percentage of individuals in these two categories, along with the χ -square test, were
196 calculated for each independent variable. The number, percentage, and p-value of each item
197 are described in Table 1.

198

199 Second, a multivariate logistic regression analysis was conducted to identify potential
200 predictive factors in the occurrence of non-consensual sex. This analysis encompassed all the
201 aforementioned exposure variables, and odds ratios with 95% confidence intervals (CIs) were
202 estimated (Table 2). As a sub-analysis, the incidence of suicidal ideation by age was also
203 calculated.

204

It should be noted that because the characteristics of the participants in this Internet survey may differ from the general population (i.e., individuals with low digital skills might be excluded), a weighted analysis was performed to minimize differences in demographics, economic status, and health-related characteristics. This approach aimed to approximate national estimates from a nationally representative survey.

Statistical significance was defined as $p < 0.05$. The data were analyzed using STATA V.16.1 (Stata Corp, College Station, Texas, USA).

Ethics

This study adhered to ethical standards in accordance with the 1975 Declaration of Helsinki, as revised in 2013. Prior to participating in the online survey, all participants provided informed consent through a web-based consent form. The Research Ethics Committee of the Osaka International Cancer Institute reviewed and approved the study (No. 20084).

Results

The total number of respondents, including both men and women, was 28,000. Out of these, 2,518 participants who provided invalid responses (e.g., the same number is used for all survey items; incomplete responses) and 12,673 men were excluded. Consequently, the final analysis sample consisted of 12,809 women, whose ages ranged from 15 and 79 years old.

Table 1 shows the characteristics of the weighted proportion of non-consensual sex among all participants. The overall weighted incidence rate for non-consensual sex during the five-month period (April to September 2020) was 1.1 %. The incidence rate of non-consensual sex was higher among individuals aged 15-19 years (1.1%) and 20-29 years (2.4%), which exceeded the rates in other age groups. Women with jobs (1.5%) had a higher incidence rate than those without jobs (0.7%).

Table 1: Demographics and Descriptive Statistics of Weighted Women Participants during the Five-Month from April to September 2020

Variable	Non-Consensual Sex April-Sept 2020	
	Total (n=12,809)	
	Yes N (%)	No N (%)
Total	138 (1.1)	12,671 (98.9)
Age		
15-19	8.0 (1.1)	707 (98.9)
20-29	37 (2.4)	1543 (97.7)
30-39	38 (2.0)	1845 (98.0)
40-49	25 (1.0)	2395 (98.9)
50-59	15 (0.7)	2097 (99.3)
60-69	15 (0.7)	2127 (99.3)
70-79	1.0 (0.1)	1956 (99.9)
Marital status		
Married	88 (1.1)	7929 (98.9)
Never married	29 (0.9)	3276 (99.1)
Widowed or separated	20 (1.4)	1467 (98.7)

Having children		
No children	56 (0.9)	6006 (99.1)
At least one or more	99 (1.5)	6648 (98.5)
Educational attainment		
High school or lower	54 (1.0)	4261 (99.0)
College/university or graduate school	83 (1.2)	7136 (98.9)
Employment status		
Unemployed	43 (0.7)	6259 (99.3)
In employment	95 (1.5)	6412 (98.5)
Smoking status		
Not smoker	117 (1.0)	11404 (99.0)
Current smoker	20 (1.6)	1268 (98.4)
Household Income (million JPY)		
Low	53 (1.7)	3144 (98.3)
Moderate	43 (1.3)	3226 (98.7)
High	26 (1.0)	2669 (99.0)
Unknown	14 (0.4)	3632 (99.6)
Region		
Other	86 (1.3)	6628 (98.7)
DSE (April 16-)	16 (0.8)	2117 (99.2)
DSE (April 7-)	31 (0.9)	3568 (99.1)
Fear of COVID-19 [†]		
7–15	41 (1.2)	3402 (98.8)
16–20	26 (0.6)	4173 (99.4)
21–25	34 (1.0)	3405 (99.0)
26–35	37 (2.1)	1691 (97.9)
Self-rated health		
Other than good	94 (1.3)	7349 (98.7)
Good	44 (0.8)	5322 (99.2)
Mental state change in the last month compared to before January 2020		
Worse	59 (2.2)	2662 (97.8)
None	56 (0.6)	8998 (99.4)
Better	19 (3.3)	547 (96.7)
Unknown	4 (0.7)	500 (99.3)
Suicidal thoughts		
Yes, since before COVID-19	43 (4.0)	1039 (95.1)
First happened	20 (3.5)	539 (96.5)
Never	75 (0.7)	11093 (99.3)
Desire not to talk to anyone due to worries		
Yes	76 (4.0)	2487 (97.0)
No	62 (0.6)	10184 (99.4)
Feeling isolated		
Yes	74 (3.7)	1925 (96.3)
No	64 (0.6)	10747 (99.4)
Cancellation of a family gathering due to the pandemic		
Yes	10 (0.6)	1730 (99.4)
No	127 (1.2)	10940 (98.9)
Non-payment of salary		
Yes, since before COVID-19	15 (10.3)	133 (89.7)
First experience	8 (10.4)	67 (89.6)
Never	115 (0.9)	12471 (99.1)
Lack of money to buy necessities		
Yes, since before COVID-19	35 (4.8)	692 (95.2)
First experience	22 (6.1)	340 (94.0)
Never	81 (0.7)	11639 (99.3)

235

236 [†] Classification of the Fear of COVID-19 scale scores refers to previous studies (22).

237 All estimates account for survey weights.

238

239 There was no significant difference in the incidence rate of non-consensual sex among those

240 living in areas designated as DSE (0.9 % on April 7, 2020, 0.8 % on April 16, 2020, 1.3%

241 others). A higher FCV-19S (2.1%, [26-35]) was associated with a higher incidence rate

compared to other scores (1.2% [score from 7 to 15], 0.6% [score from 16 to 20], 1.0% [score from 21 to 25]).

The incidence rate of participants who reported feeling “worse” (2.2%) or “getting better” (4.4%) before the COVID-19 pandemic was higher than those who “No change” (0.6%) and “I do not know” (0.7%). Participants who had suicidal thoughts (4.0% since before COVID-19, 3.5% first happened) had a higher incidence rate than those who had never experienced them (0.7%). Participants who reported feeling isolated (3.7%) had a higher incidence rate than those who had never felt isolated (0.6%). Additionally, participants who had experienced non-payment of salary (10.3% since before COVID-19, 10.4% first experience) had a higher incidence rate than those who had never experienced it (0.9%). Similarly, participants who had experienced a lack of money to buy necessities (4.8% since before COVID-19, 6.1% first experience) had a higher incidence rate than those who had never experienced it (0.7%).

Table 2 shows the factors associated with non-consensual sex among participants. Participants aged 15 to 19 years and those aged 20 to 29 years had higher odds of experiencing non-consensual sex compared to participants in other aging groups. (Aged 15- 19 years old: OR 4.74, 95%CI 1.37 – 16.4, Aged 20-29 years old: OR 3.20, 95%CI 1.09 – 9.38). Employed women participants were associated with higher odds of experiencing non-consensual sex than

261 unemployed participants (OR 1.68, 95% CI 1.36-2.07).

262

263 Table 2: Factors Associated with Non-Consensual Sex Among Weighted Participants During
264 the-Five-Month from April to September 2020

	Weighted sample	Weighted incidence	Crude Odds (95% CI)	Adjusted rate % (95% CI)	Adjusted odds ratio (95% CI)	P-value
Total	12,809	138 (1.1)				
Age						
15- 19	715	8.0 (1.1)	2.26(1.29, 3.94)	3.4 (1.3, 5.5)	4.74(1.37, 16.4)	0.014
20-29	1,580	37.1(2.4)	1.74(1.07, 2.81)	2.4 (1.3, 3.4)	3.2 (1.09, 9.38)	0.034
30-39	1,883	37.7 (2.0)	1.37(0.84, 2.23)	1.5 (0.9, 2.1)	1.88 (0.63, 5.55)	0.256
40-49	2,420	24.5 (1.0)	Reference	0.8 (0.3, 1.3)	Reference	-
50-59	2,112	14.5 (0.7)	0.43(0.22, 0.83)	0.8 (0.5, 1.0)	0.90 (0.37, 2.19)	0.822
60-69	2,142	14.8 (0.7)	0.49(0.26, 0.92)	1.1 (0.6, 1.7)	1.35 (0.45, 4.06)	0.585
70-79	1,957	1.0 (0.1)	0.23(0.10, 0.55)	0.05 (0.0, 0.1)	0.05 (0.02, 0.19)	<0.001
Marital status						
Married	8,017	88.3 (1.2)	Reference	1.6 (1.2, 2.0)	Reference	
Never married	3,304	29.0 (0.8)	1.35(0.96, 1.90)	0.5 (0.1, 0.8)	0.26 (0.10, 0.70)	0.007
Widowed or separated	1,487	20.3 (1.2)	0.83(0.48, 1.43)	1.5 (0.8, 2.1)	0.92 (0.69, 1.24)	0.599
Having children						
No children	6,302	42.8 (0.7)	Reference	0.8 (0.7, 0.9)	Reference	
One or more	6,507	94.9 (1.4)	1.60(1.15, 2.22)	1.2 (1.1, 1.4)	1.58 (1.23, 2.02)	<0.001
Educational Attainment						
College/university or graduate school	7,219	83.3 (1.0)	Reference	0.9 (0.8, 0.9)	Reference	
High school or less	5,590	54.4 (1.3)	0.76(0.53, 1.08)	1.3 (1.2, 1.3)	0.66 (0.57, 0.75)	<0.001
Employment Status						
Unemployed	6,237	43.2 (0.7)	Reference	0.8 (0.7, 0.9)	Reference	
Employed	6,572	94.5 (1.5)	1.701(1.23, 2.37)	1.3 (1.2, 1.4)	1.68 (1.36, 2.07)	<0.001
Smoking status						
Not smoker	11,521	117.4 (1.0)	Reference	1.1 (1.0,1.2)	Reference	
Current smoker	1,288	20.3 (1.6)	1.63 (1.05, 2.53)	1.0 (0.3,1.6)	0.89 (0.39, 2.04)	0.779
Household Income (million JPY)						
Low	3,579	57.2 (1.5)	Reference	1.4 (0.9, 1.8)	Reference	
Moderate	3,271	49.3 (1.6)	1.28(0.68, 1.56)	1.4 (1.4, 1.5)	1.06 (0.77, 1.46)	0.715
High	2,313	16.6 (0.6)	1.01(0.66, 1.53)	0.8 (0.3, 1.3)	0.57 (0.21, 1.56)	0.275
Unknown	3,647	14.5 (0.5)	0.467(0.28, 0.79)	0.4 (0.2, 0.7)	0.29 (0.11, 0.75)	0.011
Region						
Other	6,715	86.4 (2.0)	Reference	1.1 (1.0, 1.1)	Reference	
DES (April 13-)	2,133	16.3 (0.8)	1.12(0.70, 1.78)	1.1 (1.1, 1.2)	1.07 (0.99, 1.15)	0.100
DES (April 7-)	3,599	31.2 (0.5)	1.03(0.72, 1.48)	1.0 (1.0, 1.1)	0.94 (0.68, 1.24)	0.261
Fear of COVID-19						
7-15	3,443	40.9 (1.2)	Reference	1.4 (1.1, 1.7)	Reference	
16-20	4,199	25.9 (0.6)	0.781(0.50, 1.21)	0.7 (0.4, 1.0)	0.45 (0.23, 0.88)	0.019
21-25	3,439	34.3 (0.9)	0.832(0.53, 1.30)	0.9 (0.9, 1.0)	0.62 (0.47, 0.81)	0.001
26-35	1,728	36.5 (2.3)	2.00(1.28, 3.11)	1.4 (1.3, 1.5)	1.01 (0.84, 1.21)	0.927
Self-rated health						
Good	5,366	43.8 (0.7)	Reference	0.9 (0.8, 1.1)	Reference	
Other than good	7,443	93.9 (1.5)	1.39(1.01, 1.91)	1.2 (1.1, 1.2)	1.25 (0.98, 1.48)	0.069
Mental state change in the last month compared to before January 2020						
Worse	9,054	56.0 (0.6)	Reference	0.9 (0.7, 1.2)	Reference	
None	2,686	59.3 (2.2)	3.14(2.19, 4.49)	1.1 (0.8, 1.5)	1.25 (0.67, 2.33)	0.480
Better	566	18.8 (3.0)	6.53(4.15, 10.30)	1.8 (1.4, 2.3)	2.11 (1.39, 3.18)	< 0.001
Unknown	504	3.5 (0.8)	1.55(0.622, 3.88)	0.8 (0.7, 0.8)	0.80 (0.53, 1.20)	0.286
Suicidal thoughts						
Yes, since before COVID-19	11,169	75.2 (0.7)	Reference	0.9 (0.8, 1.0)	Reference	
First happened	1,082	42.8 (3.9)	5.85(4.06, 8.43)	1.6 (1.4, 1.8)	1.85 (1.50, 2.27)	<0.001
Never	558	19.7 (4.3)	7.66(4.85, 12.01)	1.1 (0.7, 1.5)	1.25 (0.79, 1.97)	0.337
Desire not to talk to anyone due to worries						
Yes	10,246	61.5 (0.6)	Reference	0.8 (0.7, 0.9)	Reference	
No	2,563	76.2 (3.2)	7.27 (5.24, 10.08)	1.5 (1.3, 1.7)	1.98 (1.56, 2.51)	<0.001
Feeling isolated						
Yes	10,811	64.1 (0.6)	Reference	0.8 (0.3, 1.2)	Reference	

No Cancellation of a family gathering due to the pandemic	1,988	73.6 (3.4)	8.35 (6.02, 11.60)	1.8 (0.6, 2.9)	2.52 (0.70, 9.04)	0.156
Yes	1,741	83.3 (1.0)	Reference	0.5 (0.1, 0.9)	Reference	
No	11,068	54.4 (1.3)	1.035(0.64, 1.68)	1.2 (1.1, 1.3)	2.58 (1.07, 6.25)	0.035
Non-payment of salary						
Yes, since before COVID-19	12,586	7.8 (7.3)	Reference	1.0 (0.9, 1.0)	Reference	
First experience	148	114.6 (0.9)	12.15(7.02, 21.01)	3.9 (2.9, 4.9)	4.88 (3.42, 6.97)	<0.001
Never	75	15.3 (10.4)	15.13(8.40, 27.28)	1.9 (1.1, 2.8)	2.17 (1.36, 3.48)	0.001
Lack of money to buy necessities						
Yes, since before COVID-19	11,720	80.5 (0.7)	Reference	0.8 (0.7, 0.9)	Reference	
First experience	727	34.9 (5.7)	6.97(4.68, 10.38)	2.2 (1.9, 2.6)	2.94 (2.27, 3.81)	<0.001
Never	362	22.2 (6.1)	7.62(4.74, 12.25)	1.9 (1.1, 2.8)	2.50 (1.76, 3.55)	<0.001

265

266 Participants who did not receive salaries or compensations were more likely to experience non-
267 consensual sex than those who did: (experienced since before COVID-19) OR 4.88, 95% CI
268 3.42-6.97; (first experience during COVID-19) OR 2.17, 95% CI 1.36-3.48. Similarly,
269 participants who could not afford to buy life necessities were more likely to experiencing non-
270 consensual sex than the others (OR 2.5, 95% CI 1.76-3.55).

271

272 Lastly, participants who had suicidal thoughts were more likely to experience non-consensual
273 sex than those who did not: (since before COVID-19: OR 1.85, 95% CI 1.50-2.27). Participants
274 with reported feeling isolated were more likely to experience non-consensual sex than those
275 who did not (OR 1.98, 95% CI 1.56-2.51). The result of the sub-analysis showed suicidal
276 ideation was present in 20% of all women aged between 15 and 19 years.

277

278 Figure 1: Forest plot illustrating incidence rate of non-consensual sex for each factor
279 (reference) among the participants.

[Insert Figure 1]

Note: Significant positive factors associated with non-consensual sex included the age group “Aged 15–29 years old”, as well as worsened mental or economic status were associated with an increased likelihood of experiencing non-consensual sex.

Discussion

This study assessed the incidence of non-consensual sex in Japan during COVID-19 and explored associated factors while applying weighing to mitigate biases related to online participation. The overall rate of participants reporting non-consensual sex was 1.2% over a five-month period from April to September 2020. Factors associated with experiencing non-consensual sex among participants included ages between 15 and 29, employment status, deteriorating mental health condition, suicidal thoughts, a desire not to talk to anyone, feeling isolated, and challenging economic circumstances.

In Japan, research related to sexual matters is limited. We believe this limitation stems from the fact that sexual topics are often considered taboo with Japanese education institutions, and sexual education in Japan lags behind that in Western countries (22–24). Additionally, the understanding of obtaining sexual consent, which is essential in all sexual activities, lags behind compared to countries that are known to be advanced in sex education in some of Europe countries (13,22,25). For instance, the Netherlands had sex education compulsory from

elementary school, and Denmark was one of the European countries that criminalized non-consensual sexual acts in 2020 at the time of our survey. Despite the efforts in Europe, conducting such a comprehensive study on a large scale was of significant importance as it ventured into uncharted territory in Japan.

Globally, concern arose about an increase in sexual violence against women around April 2020, during the early stages of the COVID-19 outbreak. However, specific incidence data were limited (12,26). While some countries reported an uptick in sexual violence cases (e.g., South Africa, Bangladesh (12,27)), others reported a significant decrease after implementing lockdown measures, as seen in Australia and Canada (28,29). In Japan, the available data included two key pieces of information from the police office: the reported prevalence of sexual violence among women, which stood at 6.9% (30), and a notable increase in the number of sexual assault victims seeking consultations, approximately 1.2 times higher in 2020 compared to 2019 (31,32). However, the incidence rate had not been thoroughly investigated during the COVID-19 pandemic, resulting in a scarcity of information regarding the incidence of sexual violence against women (33). It is important to note that our study did not have access to a comparative dataset from a different time point to evaluate percentage change in the incidence of sexual violence. Nevertheless, we were able to fundamental data concerning non-consensual

sex for the first time in Japan.

Approximately 80% of the women who reported experiencing non-consensual sex fell within the age range of 15 and 49 years old. Notably, the incidence of reported non-consensual sex dramatically decreased among women aged 50 to 79 years. There was a significant increase in the risk of non-consensual sex within the age group 15 to 29 years. This pattern is consistent with reports from the United States and Japan, where individuals in their late teens to early thirties are identified as the age group at the highest risk of experiencing rape (30,34).

It is crucial to recognize that young women who have been victims of sexual assault are also more likely to experience lifetime suicide attempts and post-traumatic stress symptom. Specifically, research indicates that teenage sexual trauma is strongly correlate with suicide attempts (35–38). Additionally, our sub-analysis results revealed that suicidal ideation was present in 20% of all women aged between 15 and 19 years. Based on the aforementioned findings and considerations, we firmly believe that measures to protect the younger generation are imperative (39). As mentioned previously, Japan falls behind comprehensive sexual education for the younger generation. We contend that it is necessary to incorporate detailed explanations and the factors and circumstances that lead to unwanted sexual experiences

involving the non-consensual sex into Japanese school curriculum.

Concretely, employed participants were more likely to be in contact with others than those who are unemployed. This time, we have not specifically investigated the locations where non-consensual sex has occurred. Therefore, it can be said that employed women may become victims of non-consensual sex, potentially by someone within their workplace or close social circle. Indeed, reports from both Japan and the United States have indicated that non-consensual sex is most frequently perpetrated by individuals known to the victims (30,40). Employed people have an extra circle of known individuals from their workplace compared to unemployed people.

Intriguingly, women who did not receive a salary and lacked the financial means to purchase necessities were at an elevated risk of experiencing non-consensual sex. While this appears to be the opposite of the above finding suggesting that being employed could be associated with a higher odds of experiencing non-consensual sex, this aligns with previous studies showing a connection between deteriorating economic conditions and an increased incidence of sexual assault that have established a connection between deteriorating economic conditions and an increased incidence of sexual assault (41,42), which is consistent with our findings. There are

two possible explanations for this association: First, worsening personal economic circumstances may lead women to engage in occupations with a higher risk of non-consensual sex, such as working in the sex industry. Second, women in precarious financial situation often experience psychological pressure to secure and maintain employment (42). Under such pressure, they may become psychologically vulnerable to sexual coercion from their employers in exchange for job security (42). Future research should explore people in these circumstances in depth to better under-explored areas in Japan.

While 66% of the participants lived in areas under DSE, our study did not find a significant association between non-consensual sex and living in the DSE area. However, we did identify a significant correlation between experiencing a moderate level of fear related to the COVID-19 pandemic (scoring 16-25 points) and instances of non-consensual sex. This suggests that participants who stayed at home more frequently might have become more vulnerable to non-consensual sexual experiences by their partner. In fact, non-consensual sex was more frequently reported as being perpetrated by partner during the COVID-19 pandemic, and there was an observed increase in the number of domestic problems during this period (31).

Based on the considerations outlined above, there is a heightened risk of non-consensual sexual

activity occurred during the COVID-19 pandemic, particularly among young, impoverished women. Consequently, it is of utmost importance to identify and provide economic and psychosocial support to these vulnerable individuals during times of social disruption. Furthermore, a notable deficiency in the current landscape of Japan is the absence of comprehensive education regarding the concept of sexual consent. Specifically, there is an urgent need for thorough instruction emphasizing the importance of men obtaining consent from women before engaging in any sexual activities (43). This education should be disseminated at both the local and national levels, as it may necessitate a societal shift in awareness at all levels.

In a noteworthy development, on June 16, 2023, Japan amended the legal term “Forced Sexual Intercourse Crime” to “Non-consensual Sexual Intercourse Crime”, explicitly stipulating that any sexual activity without consent is now considered a criminal offense (44). We anticipate that this alteration will provide a means of support to victims who have endured their suffering in silence and were unable to report their experiences.

Limitations

This study has several limitations. First, as it is a web-based survey, participants in an internet-

based study may not be representative of the general population. To address this potential bias, we performed statistical adjustments. However, it is important to note that this method may not fully account for the differences between participants in an internet survey and those in a nationwide representative survey, which poses challenges to the generalizability of our findings. To mitigate this limitation, harmonizing our data with a major national and representative cross-sectional study would enable us to pool data and potentially adjust for the factor of “being a respondent in an internet survey”, as has been done in other JACSIS studies (45).

The second limitation is related to the cross-sectional nature of survey. This design does not allow us to establish causal relationships between the presence or absence of non-consensual sex and factors such as economic status or suicidal ideation (46).

The third limitation concerns the definition of “Non-consensual sex”. We did not explicitly seek a specific definition of non-consensual sexual activity, which may vary by generations or community cultures (e.g., workplace community, religious community). Sex education in Japan is known to be under-developed, therefore respondents may not necessarily share a relatively standardized understanding of this concept. We did inquire about employment status, but we did not specifically ask about non-consensual sexual activity in the workplace. However,

we observed a significant correlation between being employed and experiencing non-consensual sexual activity. Also, we did not inquire the place of experienced non-consensual sex such as workplace. We observed a significant correlation between being employed and experiencing non-consensual sexual activity. However, it was not possible to ascertain whether non-consensual sex had occurred in the workplace.

The Forth, as mentioned in the section on “Analysis Subjects”, our questionnaire only asked whether respondents self-identified as male or female. Therefore, among those who self-identified as female, there is a possibility that transgender women or individuals who identify as bigender may also be included. This follows the convention in Japan, where many national surveys still only inquire about male or female gender identities.

Conclusion

Early intervention to prevent individuals from becoming victims of sexual harm should be extended to economically vulnerable and young women, especially during times of societal upheaval such as the COVID-19 pandemic. Additionally, Japan should prioritize the implementation of comprehensive education on the concept of sexual consent.

References

1. Jozkowski KN, Sanders S, Peterson ZD, Dennis B, Reece M. Consenting to sexual activity: The development and psychometric assessment of dual measures of consent. Arch Sex Behav [Internet]. 2014 Jan 23 [cited 2022 Nov 27];43(3):437–50. Available from: <https://link.springer.com/article/10.1007/s10508-013-0225-7>
2. Jozkowski KN, Peterson ZD. Assessing the Validity and Reliability of the Perceptions of the Consent to Sex Scale. <http://dx.doi.org/10.1080/002244992012757282> [Internet]. 2014 Aug 1 [cited 2022 Nov 27];51(6):632–45. Available from: <https://www.tandfonline.com/doi/abs/10.1080/00224499.2012.757282>
3. European Union Agency for Fundamental Rights. Violence against women : an EU-wide survey. Main results report. Hum Rights Doc online [Internet]. 2014 [cited 2022 Dec 15]; Available from: https://www.academia.edu/31782914/Violence_against_women_an_EU_wide_survey_Main_results_report
4. Tharp AT, DeGue S, Valle LA, Brookmeyer KA, Massetti GM, Matjasko JL. A Systematic Qualitative Review of Risk and Protective Factors for Sexual Violence Perpetration. Trauma, Violence, Abus. 2022 Apr;14(2):133–67.
5. Delker BC, Salton R, McLean KC, Syed M. Who has to tell their trauma story and

how hard will it be? Influence of cultural stigma and narrative redemption on the
storying of sexual violence. PLoS One [Internet]. 2020 Jun 5 [cited 2022 Nov
2];15(6):e0234201. Available from:
<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0234201>

6. Anastario M, Lawry L, Shehab N. Increased Gender-based Violence Among Women
Internally Displaced in Mississippi 2 Years Post–Hurricane Katrina. Disaster Med
Public Health Prep [Internet]. 2009 Mar [cited 2022 Jun 23];3(1):18–26. Available
from: [https://www.cambridge.org/core/journals/disaster-medicine-and-public-health-
preparedness/article/abs/increased-genderbased-violence-among-women-internally-
displaced-in-mississippi-2-years-posthurricane-
katrina/D576CBEB73639E69823A1391B5BF160B](https://www.cambridge.org/core/journals/disaster-medicine-and-public-health-preparedness/article/abs/increased-genderbased-violence-among-women-internally-displaced-in-mississippi-2-years-posthurricane-katrina/D576CBEB73639E69823A1391B5BF160B)

7. Thornton W, Voigt L. Disaster rape: Vulnerability of women to sexual assaults during
Hurricane Katrina. J Public Manag Soc ... [Internet]. 2007 [cited 2022 Jun 23];13:23–
49. Available from: [http://sites.google.com/a/jpmsp.com/www/Vol13Iss2-
DisasterRape-ThorntonandVo.pdf](http://sites.google.com/a/jpmsp.com/www/Vol13Iss2-DisasterRape-ThorntonandVo.pdf)

8. Picardo CW, Burton S, Naponick J, Katrina Reproductive Assessment Team.
Physically and sexually violent experiences of reproductive-aged women displaced by
Hurricane Katrina. J La State Med Soc [Internet]. 2010 Sep 1 [cited 2022 Jun

23];162(5):282, 284–8, 290. Available from:

<https://europepmc.org/article/med/21141260>

9. Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on College

Students' Mental Health in the United States: Interview Survey Study. J Med Internet

Res 2020;22(9)e21279 <https://www.jmir.org/2020/9/e21279> [Internet]. 2020 Sep 3

[cited 2022 Feb 14];22(9):e21279. Available from:

<https://www.jmir.org/2020/9/e21279>

10. Wakata S, Nishioka D, Takaki Y. Changes in health-related quality of life scores

among low-income patients on social welfare programs in Japan during the COVID-19

pandemic: a single-center repeated cross-sectional study. BMC Public Health

[Internet]. 2022 Dec 1 [cited 2022 Dec 7];22(1):1–8. Available from:

<https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-14597-5>

11. European Union Agency for Fundamental Rights. Violence against women: An EU-

wide survey main results, European Union Agency for Fundamental Rights [Internet].

2014 [cited 2022 Jun 23]. Available from:

[https://fra.europa.eu/en/publication/2014/violence-against-women-eu-wide-survey-](https://fra.europa.eu/en/publication/2014/violence-against-women-eu-wide-survey-main-results-report)

[main-results-report](https://fra.europa.eu/en/publication/2014/violence-against-women-eu-wide-survey-main-results-report)

12. al Mamun F, Hosen I, Mamun MA. Sexual violence and rapes' increment during the

- 482 COVID-19 pandemic in Bangladesh [Internet]. Vol. 34, EClinicalMedicine. Lancet
483 Publishing Group; 2021 [cited 2022 Nov 2]. Available from:
484 <http://www.thelancet.com/article/S2589537021000973/fulltext>
- 485 13. Nishioka E. Historical Transition of Sexuality Education in Japan and Outline of
486 Reproductive Health/Rights. Nihon Eiseigaku Zasshi [Internet]. 2018 [cited 2022 Nov
487 27];73(2):178–84. Available from:
488 https://www.researchgate.net/publication/325459300_Historical_Transition_of_Sexuality_Education_in_Japan_and_Outline_of_Reproductive_HealthRights
- 489 ity_Education_in_Japan_and_Outline_of_Reproductive_HealthRights
- 490 14. Miyawaki A, Tabuchi T, Tomata Y, Tsugawa Y. Association between participation in
491 the government subsidy programme for domestic travel and symptoms indicative of
492 COVID-19 infection in Japan: Cross-sectional study. BMJ Open [Internet]. 2021 Apr
493 13 [cited 2021 Jun 24];11(4):49069. Available from:
494 <http://dx.doi.org/10.1136/bmjopen-2021-049069>
- 495 15. Rakuten Insight. Rakuten Insight - Asia's Leader in Online Market Research
496 [Internet]. 2021 [cited 2022 Nov 2]. p. 1. Available from: <https://insight.rakuten.com/>
- 497 16. Hosokawa Y, Okawa S, Hori A, Morisaki N, Takahashi Y, Fujiwara T, et al. The
498 Prevalence of COVID-19 Vaccination and Vaccine Hesitancy in Pregnant Women: An
499 Internet-based Cross-sectional Study in Japan. J Epidemiol [Internet]. 2022 [cited 2023

- 500 Nov 12];32(4):188–94. Available from: <https://doi.org/10.2188/jea.JE20210458>
- 501 17. Prime Minister of Japan and His Cabinet. Guidelines for Lifting the State of
502 Emergency [Internet]. 2020 [cited 2021 Jul 7]. Available from:
503 https://japan.kantei.go.jp/ongoingtopics/_00025.html
- 504 18. Kyodo News+. Abe declares coronavirus emergency over in Japan [Internet]. [cited
505 2021 Jul 7]. Available from:
506 [https://english.kyodonews.net/news/2020/05/a1f00cf165ae-japan-poised-to-end-state-](https://english.kyodonews.net/news/2020/05/a1f00cf165ae-japan-poised-to-end-state-of-emergency-over-coronavirus-crisis.html)
507 [of-emergency-over-coronavirus-crisis.html](https://english.kyodonews.net/news/2020/05/a1f00cf165ae-japan-poised-to-end-state-of-emergency-over-coronavirus-crisis.html)
- 508 19. Sieg L. Explainer: As COVID-19 cases surge, Japan sticks to “lockdown-lite” |
509 Reuters [Internet]. 2021 [cited 2022 Dec 15]. Available from:
510 [https://www.reuters.com/world/asia-pacific/covid-19-cases-surge-japan-sticks-](https://www.reuters.com/world/asia-pacific/covid-19-cases-surge-japan-sticks-lockdown-lite-2021-08-06/)
511 [lockdown-lite-2021-08-06/](https://www.reuters.com/world/asia-pacific/covid-19-cases-surge-japan-sticks-lockdown-lite-2021-08-06/)
- 512 20. Wakashima K, Asai K, Kobayashi D, Koiwa K, Kamoshida S, Sakuraba M. The
513 Japanese version of the Fear of COVID-19 scale: Reliability, validity, and relation to
514 coping behavior. PLoS One [Internet]. 2020 Nov 5 [cited 2022 Dec
515 15];15(11):e0241958. Available from:
516 <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0241958>
- 517 21. Ahorsu DK, Lin C-Y, Imani V, Saffari M, Griffiths MD, Pakpour AH. The Fear of

- 518 COVID-19 Scale: Development and Initial Validation. Int J Ment Heal Addict 2020
- 519 [Internet]. 2020 Mar 27 [cited 2021 Jul 9];1–9. Available from:
- 520 <https://link.springer.com/article/10.1007/s11469-020-00270-8>
- 521 22. Hashimoto N, Ushitora K, Morioka M, Motegi T, Tanaka K, Tashiro M, et al. Sex
- 522 Education Sexuality, Society and Learning School education and development of
- 523 gender perspectives and sexuality in Japan. 2017 [cited 2022 Nov 27]; Available from:
- 524 <http://www.tandfonline.com/action/journalInformation?journalCode=csed20>
- 525 23. MARCH 2021 Global Gender Gap Report The analysis presented in the Global
- 526 Gender Gap Report 2021. 2021 [cited 2022 Nov 27]; Available from:
- 527 <http://reports.weforum.org/global->
- 528 24. Fu H. The bumpy road to socialise nature: sex education in Japan. Cult Health Sex
- 529 [Internet]. 2011 Sep [cited 2022 Dec 15];13(8):903–15. Available from:
- 530 <https://pubmed.ncbi.nlm.nih.gov/21787254/>
- 531 25. Ponzetti JJ. Evidence-based approaches to sexuality education: A global perspective.
- 532 Evidence-based Approaches to Sex Educ A Glob Perspect. 2015 Jan 1;359–84.
- 533 26. UN Women. COVID-19 and Ending Violence Against Women and Girls [Internet].
- 534 United Nations Entity for Gender Equality and the Empowerment of Women (UN
- 535 Women); 2020 [cited 2021 Jul 7]. Available from: <https://www.unwomen.org/>

- 536 /media/headquarters/attachments/sections/library/publications/2020/issue-brief-covid-
537 19-and-ending-violence-against-women-and-girls-en.pdf?la=en&vs=5006
- 538 27. Ditekemena JD, Luhata C, Mavoko HM, Siewe Fodjo JN, Nkamba DM, Van Damme
539 W, et al. Intimate Partners Violence against Women during a COVID-19 Lockdown
540 Period: Results of an Online Survey in 7 Provinces of the Democratic Republic of
541 Congo. Int J Environ Res Public Heal 2021, Vol 18, Page 5108 [Internet]. 2021 May
542 12 [cited 2022 Nov 2];18(10):5108. Available from: [https://www.mdpi.com/1660-](https://www.mdpi.com/1660-4601/18/10/5108/htm)
543 4601/18/10/5108/htm
- 544 28. Payne JL, Morgan A, Piquero AR. COVID-19 and social distancing measures in
545 Queensland, Australia, are associated with short-term decreases in recorded violent
546 crime. J Exp Criminol [Internet]. 2022 Mar 1 [cited 2022 Nov 2];18(1):89–113.
547 Available from: <https://link.springer.com/article/10.1007/s11292-020-09441-y>
- 548 29. Muldoon KA, Denize KM, Talarico R, Fell DB, Sobiesiak A, Heimerl M, et al.
549 COVID-19 pandemic and violence: rising risks and decreasing urgent care-seeking for
550 sexual assault and domestic violence survivors. BMC Med [Internet]. 2021 Dec 1
551 [cited 2021 Jun 24];19(1):1–9. Available from: [https://doi.org/10.1186/s12916-020-](https://doi.org/10.1186/s12916-020-01897-z)
552 01897-z
- 553 30. Gender Equality Bureau, Cabinet Office J. Survey on Violence between Men and

- 554 Women [Internet]. 2018 [cited 2022 Dec 15]. Available from:
555 http://www.gender.go.jp/policy/no_violence/e-vaw/chousa/h11_top.html
- 556 31. Hashimoto S. Sexual violence consultations in Japan up 15 % during pandemic. 2020
557 [cited 2022 Dec 4];1–2. Available from:
558 [https://english.kyodonews.net/news/2020/11/946cdc3cfe27-sexual-violence-](https://english.kyodonews.net/news/2020/11/946cdc3cfe27-sexual-violence-consultations-in-japan-up-15-during-pandemic.html?phrase=sexcrimes&words=sex,crime,crimes,Sex)
559 [consultations-in-japan-up-15-during-pandemic.html?phrase=sex](https://english.kyodonews.net/news/2020/11/946cdc3cfe27-sexual-violence-consultations-in-japan-up-15-during-pandemic.html?phrase=sexcrimes&words=sex,crime,crimes,Sex)
560 [crimes&words=sex,crime,crimes,Sex](https://english.kyodonews.net/news/2020/11/946cdc3cfe27-sexual-violence-consultations-in-japan-up-15-during-pandemic.html?phrase=sexcrimes&words=sex,crime,crimes,Sex)
- 561 32. The White Paper on Gender Equality 2021 [Internet]. 2021 Jun [cited 2022 Dec 15].
562 Available from:
563 https://www.gender.go.jp/english_contents/about_danjo/whitepaper/pdf/ewp2021.pdf
- 564 33. Mittal S, Singh T. Gender-Based Violence During COVID-19 Pandemic: A Mini-
565 Review. Front Glob Women’s Heal. 2020 Sep 8;0:4.
- 566 34. Morgan RE, Statistician B, Mason BJ, Intern B. Crimes Against the Elderly, 2003–
567 2013. 2014 Nov.
- 568 35. Olshen E, McVeigh KH, Wunsch-Hitzig RA, Rickert VI. Dating violence, sexual
569 assault, and suicide attempts among urban teenagers. Arch Pediatr Adolesc Med
570 [Internet]. 2007 Jun 1 [cited 2021 Jun 24];161(6):539–45. Available from:
571 <https://jamanetwork.com/journals/jamapediatrics/fullarticle/570505>

- 590 wants-talk-sex-education/
- 591 40. Breiding MJ, Smith SG, Basile KC, Walters ML, Chen J, Merrick MT. Prevalence and
592 Characteristics of Sexual Violence, Stalking, and Intimate Partner Violence
593 Victimization—National Intimate Partner and Sexual Violence Survey, United States,
594 2011. MMWR Surveill Summ [Internet]. 2014 Sep 9 [cited 2022 Aug 23];63(8):1.
595 Available from: /pmc/articles/PMC4692457/
- 596 41. Markowitz FE. Socioeconomic disadvantage and violence: Recent research on culture
597 and neighborhood control as explanatory mechanisms. Aggress Violent Behav. 2003
598 Mar 1;8(2):145–54.
- 599 42. Department of Reproductive Health and Research World Health Organization.
600 Violence against women [Internet]. 2014 [cited 2021 Jun 24]. Available from:
601 https://apps.who.int/iris/bitstream/handle/10665/112325/WHO_RHR_14.11_eng.pdf
- 602 43. Willis M, Hunt M, Wodika A, Rhodes DL, Goodman J, Jozkowski KN. Explicit
603 Verbal Sexual Consent Communication: Effects of Gender, Relationship Status, and
604 Type of Sexual Behavior. <https://doi.org/10.1080/1931761120191565793> [Internet].
605 2019 Jan 2 [cited 2022 Nov 27];31(1):60–70. Available from:
606 <https://www.tandfonline.com/doi/abs/10.1080/19317611.2019.1565793>
- 607 44. Ministry of Justice. Ministry of Justice: FAQs on Amendments to Laws Related to

- 608 Sexual Crimes [Internet]. [cited 2023 Nov 12]. Available from:
- 609 https://www.moj.go.jp/keiji1/keiji12_00200.html
- 610 45. Tabuchi T, Shinozaki T, Kunugita N, Nakamura M, Tsuji I. Study Profile: The Japan
- 611 “Society and New Tobacco” Internet Survey (JASTIS): A Longitudinal Internet
- 612 Cohort Study of Heat-Not-Burn Tobacco Products, Electronic Cigarettes, and
- 613 Conventional Tobacco Products in Japan. J Epidemiol [Internet]. 2019 Nov 5 [cited
- 614 2022 Dec 21];29(11):444–50. Available from: <https://doi.org/10.2188/jea.JE20180116>
- 615 46. Kotera Y, Van Laethem M, Ohshima R. Cross-cultural comparison of mental health
- 616 between Japanese and Dutch workers: relationships with mental health shame, self-
- 617 compassion, work engagement and motivation. Cross Cult Strateg Manag. 2020 Sep
- 618 3;27(3):511–30.
- 619

Figure 1. Forest plot of the incidence rate of non-consensual sex of each factor (reference) among the participants.

