

1 A Breast Cancer Polygenic Risk Score Validation in 15,490 Brazilians

2 using Exome Sequencing

3

4 Flávia Eichemberger Rius^{1,2}, Rodrigo Guindalini³, Danilo Viana¹, Júlia Salomão¹,
 5 Laila Gallo¹, Renata Freitas¹, Cláudia Bertolacini¹, Lucas Taniguti¹, Danilo
 6 Imparato¹, Flávia Antunes¹, Gabriel Sousa¹, Renan Achjian¹, Eric Fukuyama¹,
 7 Cleandra Gregório¹, Iuri Ventura¹, Juliana Gomes¹, Nathália Taniguti¹, Simone
 8 Maistro², José Eduardo Krieger⁴, Yonglan Zheng⁵, Dezheng Huo⁶, Olufunmilayo I.
 9 Olopade⁵, Maria Aparecida Koike², David Schlesinger¹

10

11 1. Mendelics, São Paulo, Brazil.

12 2. Comprehensive Center for Precision Oncology - C2PO, Centro de Investigação
 13 Translacional em Oncologia (CTO), Departamento de Radiologia e Oncologia, Instituto
 14 do Cancer do Estado de Sao Paulo (ICESP), Hospital das Clinicas HCFMUSP,
 15 Faculdade de Medicina, Universidade de Sao Paulo, Sao Paulo, Brazil.

16 3. Instituto D'Or de Pesquisa e Ensino (IDOR), São Paulo, Brazil.

17 4. Instituto do Coração, Hospital das Clínicas da Faculdade de Medicina da
 18 Universidade de São Paulo - FMUSP, São Paulo, Brazil.

19 5. Medicine and Human Genetics, Center for Clinical Cancer Genetics and Global
 20 Health, University of Chicago Medical Center, Chicago, USA.

21 6. Department of Public Health Sciences, University of Chicago, Chicago, USA.

22

23 Abstract

24 Purpose

25 Brazil has a highly admixed population. Polygenic Risk Scores (PRS) have been mostly
26 developed from European population studies and applying them to other populations is
27 challenging. To assess the use of PRS for breast cancer (BC) risk in Brazil, we validated
28 four PRSs developed in the Brazilian population.

29 Patients and Methods

30 We analyzed 6,362 women with a history of breast cancer and 9,128 unphenotyped
31 adults as controls in a sample obtained from a clinical laboratory. Genomic variants
32 were imputed from exomes and scores were calculated for all samples.

33 Results

34 After excluding individuals with known pathogenic or likely pathogenic variants in
35 *BRCA1*, *BRCA2*, *PALB2*, *PTEN*, or *TP53*, and first-degree relatives of the probands,
36 5,730 cases and 8,847 controls remained. Four PRS models were compared, and PRS
37 3820 from Mavaddat *et al.* 2019 performed best, with an Odds Ratio (OR) of 1.41 per
38 standard deviation (SD) increase (p-value: < 0.0001) and an OR of 1.94 (p-value: <
39 0.0001) for the individuals in the top risk decile. PRS 3820 also performed well for
40 different ancestry groups: East Asian majority (Group 1), Non-European majority (Group
41 2), and European majority (Group 3), showing significant effect sizes for all groups:
42 (Group 1: OR 1.54, p-value 0.006; Group 2: OR 1.44, p-value: <0.001; Group 3 OR:

1.43, p-value: <0.001). PRS 90% compares with monogenic moderate BC risk genes (PRS90 OR: 1.94; CHEK2 OR: 1.89; ATM OR: 1.99).

Conclusion

PRS 3820 can be accurately used in the Brazilian population. This will allow a more precise BC risk assessment of mutation-negative women in Brazil.

Introduction

Breast cancer (BC) is a critical global health concern, representing the most common cancer diagnosed among women¹. In Brazil, over 70,000 women are diagnosed with BC every year, accounting for 30% of all cancers in the female population².

Approximately 10% of all BC cases are attributable to germline pathogenic variants in susceptibility genes³. Rare variants in high penetrance genes (*BRCA1*, *BRCA2*, *TP53*, *PTEN*, and *PALB2*) and in moderate penetrance genes (*CHEK2* and *ATM*) are associated with a more than 4-fold and 1.5–4 fold increased risk of BC, respectively^{4,5}. Rare variants in these genes account for approximately 25% of the genetic risk. The remaining genetic risk (~75%) is derived from common, low penetrance variants that individually confer small risk, but which combined effect can be substantial^{4–6}.

Genome-wide association studies (GWASs) have been predominantly carried out in European populations^{7–10}. Evaluation of PRS across different genetic and environmental backgrounds is essential to enable the implementation of genetic risk stratification strategies for individuals from non-European populations¹¹.

The Brazilian population exhibits a unique, highly admixed, genetic composition. It is mostly derived from a combination of Native Americans, Southern Europeans (Portuguese, Spanish, and Italian) that immigrated in the period 1500-1900, and Sub-Saharan Africans brought through extensive slave trading until the 1800s. More recently, from 1822 to the first half of the 1900s, other smaller waves of immigration also contributed to Brazil's remarkable diversity, including Japanese, Lebanese, German, and Eastern Europeans¹². Three in every four Brazilians have multiple genetic ancestries^{13,14}. Given Brazil's genetic diversity, any PRS developed in predominantly European populations requires validation before it can be used in clinical settings. Several laboratory methods are available for genotyping variants directly or indirectly (imputation), including microarrays, whole exome sequencing (WES), and whole genome sequencing (WGS). WES offers an affordable and scalable alternative to arrays and WGS, while allowing for simultaneous rare and common variant genotyping. In this study, we evaluate four BC PRSs^{7,8,15} developed using WES in 15,490 Brazilians.

Methods

Study population

A total of 15,490 individuals were selected for this study, including 6,362 women with breast cancer history, and 9,128 adult unphenotyped controls. Both clinical and genetic data were collected from a database of a College of American Pathology (CAP)–accredited laboratory (Mendelics, São Paulo, SP, Brazil). All BC and control subjects provided Informed Consentment for use of retrospective anonymized data for

research purposes. Samples were anonymized before analysis. Clinical records such as BC histological type and age of diagnosis were obtained from genetic test requisitions. The study was IRB-approved (CAAE: 70112423.3.0000.0068).

Relatedness calculation and data filtering

Relatedness of individuals was obtained from the exomes using somalier software¹⁶, following the standard protocol required for a VCF file (<https://github.com/brentp/somalier#readme>). Concerning related individuals removal, if two individuals had a first-degree relationship, one of them was randomly selected to be included in the dataset. However, if individuals had two or more first-degree relationships, all related individuals were excluded from the dataset. This process resulted in a total of 211 removals. Furthermore, 73 individuals were removed from the sample due to unavailability of files necessary for genome imputation.

PRS analyses were performed after filtering out cases and controls with pathogenic or likely-pathogenic (P/LP) variants in BC genes *BRCA1*, *BRCA2*, *TP53*, *PALB2*, and *PTEN*.

Exome sequencing and imputation

Exome sequencing data were generated from buccal swab or venous blood samples with standard protocol for Illumina Flex Exome Prep, using a custom probe set from Twist Biosciences. Sequencing was conducted in Illumina sequencers and the bioinformatics pipeline for data analysis followed Broad Institute's GATK best practices

107 (<https://gatk.broadinstitute.org/hc/en-us/sections/360007226651-Best-Practices-Workflo>
108 ws), with alignment to GRCh38.

109 Imputation of exomes was based on a panel of 2,504 individuals of all ancestries from
110 the 1000 Genomes Project (1KGP)¹⁷ on GRCh38 (2017 release)
111 (<https://www.internationalgenome.org/data-portal/data-collection/grch38>). All regions
112 captured from the exome sequencing comprehending at least 1x coverage, as well as
113 off-target regions, were considered for the imputation, performed using Glimpse (v1.1.0)
114 software¹⁸.

115

116 Polygenic Risk Score calculation

117 Four BC PRSs with publicly available summary statistics, from three different studies,
118 were evaluated in this work: Khera *et al.* 2018⁷, with 5,218 variants; Mavaddat *et al.*
119 2019⁸ PRSs (with 313 and 3,820 variants); and UK Biobank¹⁵ (UKBB) PRS obtained
120 from a variant thresholding (p-value < 10e-5) on summary statistics for phenotype code
121 20001_1002, with 7,538 variants.

122 The PRS variants were selected based on exome bed kit distance and minor allele
123 frequency (MAF). Additionally, the PRS from Mavaddat study, originally with 3,820
124 variants, had a pathogenic variant of moderate-penetrance in *CHEK2* gene (*CHEK2*
125 p.Ile157Thr - Clinvar: RCV000144596) that was removed to avoid conflation with
126 monogenic risk.

127 PRS calculation was performed using a software developed by Mendelics, evaluating
128 the weighted sum of beta values, in which weights are based on the number of the

individual's alleles containing the variant of the PRS file. The sum is normalized by all beta positive and negative values so the final value can be between zero and one.

Genetic Principal Component Analyses (PCA) Assessment

PCA was calculated for exomes from a projection in 1KGP¹⁷ and Human Genome Diversity Project (HGDP)¹⁹ samples. Only variants with MAF > 1% and that could have been directly genotyped using WES were included for the PCA analysis in 1KGP and HGDP samples using plink2²⁰. Exomes were converted to plink bfile format (bed, bim, and fam files) and had duplicated variants removed. PCA projection for 10 PCs was calculated using plink2 –score method, with allele frequencies from the breast cancer case-control sample.

Ancestry evaluation

Admixture²¹ was used to extract continental ancestries from all non-related and data completed exomes. The analysis was supervised by the 1KGP samples, after removal of South Asian, Oceania, and admixed Americans from the GRCh38 1KGP release of 2017. South Asian and Oceania ancestries were removed because they are not a significant part of Brazilian ancestral composition. Latin American admixed populations (Colombian, Peruvian, Puerto Rican, and Mexican) were removed to avoid confounding with the native americans belonging to the same population label. Continents evaluated were: Africa - AFR, America - AMR, East Asia - EAS, and Europe - EUR. Ancestry

150 results were further used for splitting individuals into groups according to their ancestry
151 composition, to further analyze the effect size of PRS on each group.

152

153 Paired imputed and sequenced genomes analysis

154 Exome-imputed variants and directly sequenced variants from WGS were compared
155 using 1001 samples from an independent Brazilian population dataset
156 (<http://elsabrazil.org/>) that had both WES and WGS available. The WES were
157 sequenced and imputed also using the same method previously described. BC
158 PRS-3820 from Mavaddat *et al.* study was calculated for both imputed and sequenced
159 genomes, and their Spearman correlation was calculated using R software base
160 function *cor.test*.

161

162 Statistical analyses

163 PRS values were standardized according to the control values prior to all statistical
164 analyses. PCs were Z-scored prior to analyses. To assess the effect size of PRS on
165 breast cancer status (0 = control, 1 = case) corrected for PCs, Odds Ratio (OR) per
166 standard deviation of PRS was calculated by performing a logistic regression of BC
167 status with PRS and PCs 1 to 10 as predictors. AUC for the full dataset evaluation was
168 obtained using the yardstick R package (yardstick.tidymodels.org/) *roc_auc* function, in
169 the testing data split (25%). In order to find segmentation effect-sizes, individuals were
170 classified into deciles or percentiles following the left-open and right-closed intervals.
171 OR for deciles was calculated by first selecting only the decile analyzed and the interval

from 40-60% individuals as the control section, and binarizing it (0 = belongs to the control interval 40-60%, 1 = belongs to the decile analyzed, for example, 10%); and performing a logistic regression analysis on the binarized decile information with correction for PCs 1 to 10. A similar approach was conducted for calculating the OR on percentiles for comparison with Mavaddat's⁸ PRS validation. For each ancestry proportion group, AUC was estimated using 10-fold cross-validation with the R package *caret*²². All PRS 95% confidence intervals (CI) were obtained from the logistic regression output from the R function *glm* (stats package²³). OR and CI for genes *BRCA1*, *BRCA2*, *PALB2*, *TP53*, *ATM* and *CHEK2* were obtained using *epitools* R package²⁴. All statistical tests performed were two-tailed.

Results

Case-control sample selection and characteristics

After removal of 211 subjects with a first-degree relationship and 73 with missing files necessary for imputation, a total of 15,206 subjects remained (**Supplementary Table 1**). Four percent of all cases and controls were removed from the analysis due to their presence of pathogenic or likely-pathogenic (P/LP) variants in high penetrance genes with OR > 5 for breast cancer: *BRCA1*, *BRCA2*, *TP53*, *PALB2*, and *PTEN* (n = 629). Therefore, the sample used for PRS evaluation consisted of 5,730 women with a BC history, and 8,847 unphenotyped controls, both with known sex and age (**Table 1**).

Table 1. Demographics of cases and controls in BC dataset used for PRS evaluation

		Case	Control	Total	p value
	Total	5,730	8,847	14,577	-
Sex	F	4,225	5,730	9,955	-
	M	-	4,662	4,662	-
Age	Total	49.5 (11.7)	41.6 (13.3)	44.8 (13.3)	0.000
	F	49.5 (11.7)	42 (13.7)	46.4 (13.1)	0.000
	M	-	41.3 (12.9)	41.3 (12.9)	-

194 p-values obtained from two-tailed t-tests.

195

196 Ancestry composition of our sample was obtained using ADMIXTURE analysis²¹,
197 supervised by EUR, EAS, AFR and non-admixed AMR populations of 1KGP and HGDP.
198 The results show that the majority of individuals have EUR as their greatest ancestry
199 proportion (median 84%, SD 18%). Besides that, a significant portion of AFR (median
200 6%, std. dev. 12%) and AMR (median 8%, SD 7%) ancestries are present,
201 complemented with a variety of EUR proportions. A small quota of EAS is also observed
202 (median < 1%, SD 12%), composed by 214 individuals with over 70% of this ancestry.



203

204 **Figure 1. Ancestry composition of our Brazilian cohort.** Estimated ancestries are
 205 shown as proportions per individual. Each thin bar represents one individual and their
 206 ancestry proportion. Europe (EUR) in purple, Africa (AFR) in blue, East Asia (EAS) in
 207 green and America (AMR) in yellow.

208

209 Effect sizes of four different PRSs in the Brazilian population

210 Four PRS files from three studies were selected for initial effect size investigation in our
 211 cohort (**Supplementary Table 2**). All four PRS files had their variants further filtered to
 212 address only variants covered by the imputation of our exomes.

213


PCA was performed on the exomes to capture the population genetic structure. PRSs were calculated for the imputed genomes (details described in the **Methods**) and standardized by z-score to improve interpretability. To avoid confounding from P/LP variants on PRS effect, we have evaluated only individuals without those rare variants ($n = 14,577$). Effects were corrected for the ten first PCs, and results are all reported in **Supplementary Table 3**.

Both PRS_{Broad} and PRS_{3820} performed well, with very significant effect sizes (both p-values < 0.0001) following the direction of risk rise as the PRS increases (OR_{Broad} : 1.52; OR_{3820} : 1.41). PRS_{313} and PRS_{UKBB} have not reached significance level for their OR results (p-value₃₁₃: 0.315 and p-value_{UKBB}: 0.985). Goodness of fit of the model is also greater for PRS_{3820} (Nagelkerke pseudo- R^2 : 0.061) and PRS_{Broad} (Nagelkerke pseudo- R^2 : 0.051). Note that pseudo- R^2 values should not be interpreted as a linear regression R^2 value, but as a metric of improvement from null model to fitted model, which has its value mainly by being compared between different PRS models in which a greater pseudo- R^2 indicates a better goodness of fit to the data.

Since PRS_{Broad} and PRS_{3820} showed significant results per standard deviation, they were used to split the data into deciles to evaluate BC risk conferred by PRS in each strata. These analyses were also corrected for the first ten PCs. Interestingly, shorter confidence intervals and a better “staircase” shape can be seen for PRS_{3820} plot in comparison to PRS_{Broad} (**Figure 2**). Moreover, especially the top 10% (90-100% interval) present a much greater effect for PRS_{3820} (OR_{90-100} : 1.94; CI: 1.71 - 2.20) compared to

PR_{Broad} (OR₉₀₋₁₀₀: 1.77; CI: 1.51 - 2.10) (Supplementary Table 4), indicating a better performance of the former in identifying women with increased risk of BC. Therefore we decided to focus our next analyses on PR₃₈₂₀, which was the best PRS to identify BC risk in our Brazilian population.

241



242

Figure 2. Effect sizes by decile of PR₃₈₂₀ and PR_{Broad}. Odds Ratios (OR) and Confidence Intervals (CI) for PR₃₈₂₀ (red) and PR_{Broad} (blue). ORs for both PRS deciles were corrected for the first ten PCs.


246

PR₃₈₂₀ performance compared with the original study

As seen in the previous results, the PR₃₈₂₀ showed a positive association with increased risk of BC (OR per standard deviation: 1.41; CI: 1.36 - 1.47) after correction

for the first 10 principal components (PCs). This association was slightly lower when compared to the original study test set, composed of only Europeans (OR: 1.66; CI: 1.61 - 1.70). Besides that, performance of our model with PRS₃₈₂₀ in identifying BC cases was very similar to the original study (AUC_{Brazilians}: 0.610 vs. AUC_{Europeans}: 0.636). After calculating OR per percentiles, we observed that the PRS₃₈₂₀ exhibited an expressive risk increase for our admixed population, although the increase was smaller than the original study, which applied the PRS₃₈₂₀ to a population with the same ancestry it was originated from (OR_{Brazilians} >99: 2.72; OR_{Europeans} >99: 3.95).

258



259

Figure 3. Comparison of PRS₃₈₂₀ performance for Europeans and Brazilians. The plot shows the PRS₃₈₂₀ adapted in this study (orange), with 2,892 variants, compared with the original from Mavaddat *et al.* study (blue), with 3,820 variants.


The lower interval, comprehending the lowest 1% of PRS values, showed a smaller decrease in BC risk compared to the original study. This result is probably related to the small sample size of this section, with only 31 cases and 88 controls available to calculate OR. In addition, the 95th to 99th percentile interval exhibited marginal growth in odds ratio (OR) when contrasted with the interval immediately below (OR 90th-95th: 1.75, OR 95th-99th: 1.83). Besides that, both effect sizes show an expressive increase in BC risk due to PRS results. This might be partly due to the cohort sample size. Our study evaluated a total of 14,577 individuals, while Mavaddat's evaluated twice this number in their test dataset composed of joined cohorts ($n = 29,751$).

PRS evaluation by ancestry composition

Since our sample contains a great majority of EUR ancestry proportion, we decided to evaluate the PRS effect size in different ancestry compositions. We have created three groups: EAS majority ($> 50\%$ EAS, $n = 217$), 0 - 50% EUR ($n = 763$) and 51 - 100% EUR ($n = 13,597$). All three bins had statistically significant ($p < 0.001$) ORs above 1.40 (1.54, 1.44 and 1.43, respectively) per PRS standard deviation, showing a positive association of the PRS value with increased BC risk. The EAS majority group shows a wider confidence interval due to the small sample size (cases = 64, controls = 153). Besides that, the lower tail of the 95% confidence interval has an OR of 1.14

(Supplementary Table 5), which means at least 14% risk rise for each unit of standardized PRS increase.

A.



B.




Figure 4. Breast cancer Odds Ratio by ancestry proportion. The cohort was split into three groups based on main ancestry: EAS majority (>50% EAS), 0 - 50% EUR and 51 - 100% EUR (A) Ancestry composition of each group, with colors representing continental ancestries for each subject. (B) Breast cancer ORs by PRS₃₈₂₀ standard deviation for the three groups. p-values displayed were corrected for multiple-hypothesis testing using Bonferroni method.

294

295 **Comparison of PRS derived from genomes imputed from exomes with WGS**

296 A correlation of 0.76 (p value < 2.2e-16) was obtained between BC PRS₃₈₂₀ values
297 calculated from imputed genomes and WGS, showing a consistent concordance
298 between both methods (**Supplementary Figure 1A**).


299

300 When we compared imputed (exome) and sequenced genomes (WGS), most of the
301 extreme PRS₃₈₂₀ values were concordant (decile 1: 56%; decile 10: 60%)
302 (**Supplementary Figure 1B**). Furthermore, most of the proportion which is not in the
303 same decile is in the surrounding deciles, which indicates a low deviation from the
304 purpose of predicting risk.

305

306 **Comparison of PRS and breast cancer genes effect size**

307 For the purpose of understanding how the PRS₃₈₂₀ effect size compare to known high
308 and moderate risk genes for BC, we have compared OR of the top PRS₃₈₂₀ decile
309 (PRS90) with all pathogenic variants located in *TP53*, *BRCA1*, *BRCA2*, *PALB2*, *ATM*
310 and *CHEK2* genes (**Figure 5**) in this cohort of individuals.



311

312 **Figure 5. Effect sizes of 90th percentile of PRS and BC genes in BC risk.** Effect
313 sizes (OR and CI) were obtained according to the presence of pathogenic variants in
314 the genes *TP53*, *BRCA1*, *BRCA2*, *PALB2*, *ATM* and *CHEK2*, or belonging to the 90th to
315 100th percentiles of PRS₃₈₂₀.

316

317 As expected, *TP53*, *BRCA1* and *BRCA2* present the most extreme BC risks (OR: 14.05,
318 CI: 4.1-95.05; OR: 13.43, CI: 9.25-20.32; and OR: 8.77, CI: 6.15-12.93, respectively).
319 PRS90 risk (OR: 1.94, CI: 1.71-2.2) can be compared with moderate risk BC genes *ATM*
320 (OR: 1.99, CI: 1.42-2.78) and *CHEK2* (OR: 1.89, CI: 1.35-2.66). This result indicates
321 how an increased risk for BC due to PRS90 could be interpreted in the clinical context,
322 potentially following the same care protocols as for a moderate risk monogenic variant
323 for BC.

324

325 Discussion

326 In the present study we have validated a breast cancer PRS developed from Europeans
327 in the highly admixed Brazilian population. The PRS adapted from Mavaddat *et al.* study
328 with 2,892 variants⁸ showed a statistically significant risk prediction value (OR: 1.41 per
329 SD). Furthermore, individuals classified in the top decile had an expressive effect size
330 (OR: 1.94; CI: 1.71 - 2.20) of almost one-fold increased risk of BC compared to the
331 middle percentiles (40-60%). This PRS highest decile risk is comparable with the
332 previously reported risks for moderate-penetrance monogenic variants in *ATM*, *NF1*,
333 and *CHEK2* genes (1.82, 1.93, and 2.47 OR, respectively)²⁵, and also with risks in *ATM*
334 and *CHEK2* calculated in our sample (1.99 and 1.89 OR, respectively).

335

336 This study is based on a previous study from Mavaddat *et al.* 2019, which developed
337 and validated a PRS with 3,820 variants evaluating aggressive BC risk (metastatic BC).
338 For all BC subtypes (ER+ and ER-) they found an OR of 1.71 per SD (CI: 1.64 - 1.79) in
339 the validation set (n = 29,751; cases = 11,428), and OR 1.66 per SD (CI: 1.61 - 1.70) in
340 the prospective set (n = 190,040; cases = 3,215). These values are even greater
341 compared to the widely used 313 PRS (OR: 1.65 per SD; CI: 1.59 - 1.72 in validation
342 set). However, they included a *CHEK2* gene pathogenic variant in the PRS and worked
343 with only aggressive BC, which may have led to overestimating their OR values. A study
344 from Liu and colleagues has evaluated another modification of the same PRS with
345 3,820 variants developed from Mavaddat *et al.* for African, Latin, and European

populations²⁶. According to the study, the effect size of this PRS to a BC risk in an European sample (n = 33,594) was 1.40 per standard deviation, a result very similar to ours for a Brazilian sample (OR 1.41 per SD; n = 14,477). They deliberately have included women with *in situ* ductal BC as well as women with metastatic BC, what they claim to be a reason for OR decline compared to the original study, which included only metastatic BC women both in their discovery and validation sets. Our study, however, does not distinguish BC types, therefore we hypothesize that both metastatic and *in situ* BC are included, which may be a factor, together with genetic population structure, that decreased the OR compared to the original study.

Furthermore, significant effect sizes per PRS standard deviation were obtained for distinct ancestry compositions within our sample. Due to the high proportion of EUR (median 84%, std. dev. 18%), we separated the sample into groups with different ancestry compositions. Despite the small sample size (n = 217) of the EAS majority group (**Supplementary Table 5**), there was a statistical significance (adjusted p-value: 0.006) for the effect size in this group, which had similar magnitude (OR: 1.54, CI: 1.40-2.12) of the full sample (OR: 1.41, CI: 1.36-1.47, p-value: < 0.0001). Also, PRS₃₈₂₀ had significant and expressive effect sizes on BC risk for both EUR proportion groups (0-50% EUR OR: 1.44, CI: 1.23-1.69; 51-100% EUR OR: 1.43, CI: 1.38-1.49). These results evidence that, for individuals with a more prominent East Asian ancestry, for admixed individuals, and for predominantly Europeans, PRS₃₈₂₀ is still effective in stratifying BC risk.

369 All of our PRS values were calculated according to a new methodology: the imputation
 370 of genomes from exomes. This approach has demonstrated to be very successful for
 371 PRS calculation and assessment of BC risk in our study, and could be very interesting
 372 for laboratories that already perform exome sequencing as a cost-effective methodology
 373 to identify P/LP variants for BC. A variety of studies have compared low-pass genome
 374 sequencing with arrays for different applications, such as pharmacogenetics, GWAS,
 375 CNV detection, and PRS calculation^{27,28,29}. The study of Li *et al.*²⁸ reported improved
 376 accuracy for polygenic risk prediction of imputed low-pass genome compared to array
 377 imputation for both coronary artery disease and BC. Despite the slight difference we
 378 found between PRS values calculated from sequenced genomes and imputed genomes
 379 from exomes (Spearman correlation: 0.76), decile classification showed satisfactory
 380 concordance between both methods for the majority of results in the extreme deciles (1
 381 and 10th), which are the most important to define decreased or increased risk.
 382 Unfortunately, it was not possible to assess the predictive power of PRS values
 383 calculated from genomes of BC patients due to unavailability of paired exome and
 384 genome data.

385

386 Among familial BC cases, approximately 25% have a P/LP germline variant reported³⁰.
 387 In the Brazilian population, a robust study with 1,663 breast cancer patients detected
 388 20.1% of P/LP germline variants using multigene panel testing^{4,6}. A 2017 study reported
 389 that 18% of the hereditary BC can be explained by a polygenic effect of variants
 390 discovered in a GWAS³¹. Therefore, employing this PRS in the clinical practice might
 391 bring an elucidation to BC Brazilian families without high or moderate-effect germline

392 variants detected. Moreover, women without prior knowledge of their familial BC
393 condition, or even those with a high PRS risk by chance, will have the possibility to be
394 informed of their results and share them with their physicians to adopt preventive
395 actions accordingly to their risk strata, such as intensifying surveillance adding breast
396 magnetic resonance imaging to mammography screening³².

397

398 In conclusion, our work was able to validate a PRS developed in Europeans in the
399 Brazilian population, using imputed genomes from exomes. The top decile of this PRS
400 presents a risk comparable to moderate-risk monogenic variants for BC. Future studies
401 will be required to evaluate the combination of PRS with P/LP variants and clinical
402 factors in order to deliver more informative results to patients, thus physicians can
403 recommend prevention strategies based on their combined polygenic and monogenic
404 BC risk.

405

406 Ethics Statement

407 This work was approved by the Ethics Committee Comissão para análise de projeto de
408 pesquisa of Hospital das Clínicas da FMUSP - CAPPesq under the CAAE number
409 70112423.3.0000.0068.

410

411 **Acknowledgements**

412 We thank all individuals once sequenced in Mendelics laboratory who have consented
413 to participate in this research. We also thank all UKBB participants for their contribution
414 to the PRS hereby analyzed, and all authors from previous studies on BC PRSs in
415 which we based our validation (Khera *et al.* 2018 and Mavaddat *et al.* 2019). Maria
416 Aparecida Azevedo Koike Folgueira received research support from Conselho Nacional
417 de Desenvolvimento Científico e Tecnológico, Brazil (CNPq-308052/2022-6).

418

419 **Data Availability**

420 All variants and betas which compose the four evaluated PRSs are available as
421 Supplementary Information. Individual cases and controls data are not publicly available
422 due to the confidentiality consentment agreement signed by all included in the study.

423

424 **Competing Interests**

425 Flávia Eichemberger Rius, Danilo Viana, Júlia Salomão, Laila Gallo, Renata Freitas,
426 Cláudia Bertolacini, Lucas Taniguti, Danilo Imparato, Flávia Antunes, Gabriel Sousa,
427 Renan Achjian, Eric Fukuyama, Cleandra Gregório, Iuri Ventura, Juliana Gomes,
428 Nathália Taniguti, and David Schlesinger are currently employed by Mendelics, or were
429 employed at the time of the study.

Rodrigo Guindalini acted as a consultant for AstraZeneca, Janssen Oncology, Roche/Genentech and Igenomix; received speaker honoraria from AstraZeneca, Bristol Myers Squibb, GlaxoSmithKline, Merck Sharpe & Dohme Brasil, Novartis, and Roche outside the submitted work; and has equity in Mendelics Análise Genômica. Olufunmilayo I. Olopade is co-founder at CancerIQ; serves as scientific advisor at Tempus; and has received research funding from Color Genomics and Roche/Genentech. José Eduardo Krieger, Yonglan Zheng, Dezheng Huo, Simone Maistro and Maria Aparecida Koike declare no competing interests.

Author Contributions

Generated Main Data: Flávia Eichemberger Rius, Danilo Viana, Júlia Salomão, Laila Gallo, Renata Freitas, Cláudia Bertolacini, Lucas Taniguti, Danilo Imparato, Flávia Antunes, Gabriel Sousa, Renan Achjian, Eric Fukuyama, David Schlesinger. Analyzed Data: Flávia Eichemberger Rius, Rodrigo Guindalini, Danilo Viana, Lucas Taniguti, Danilo Imparato, Flávia Antunes, Gabriel Sousa, Renan Achjian, Eric Fukuyama, Yonglan Zheng, Dezheng Huo, Olufunmilayo I. Olopade, Maria Aparecida Koike, David Schlesinger. Other Contributions: Cleandra Gregório, Iuri Ventura, Juliana Gomes, Nathália Taniguti, Simone Maistro, José Eduardo Krieger.

References

1. Sung, H. *et al.* Global cancer statistics 2020: GLOBOCAN estimates of incidence

- 452 and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **71**,
453 209–249 (2021).
- 454 2. Instituto Nacional de Câncer. *Estimativa 2023 : incidência de câncer no Brasil*.
455 (Ministério da Saúde, 2023).
- 456 3. Nielsen, F. C., van Overeem Hansen, T. & Sørensen, C. S. Hereditary breast and
457 ovarian cancer: new genes in confined pathways. *Nat. Rev. Cancer* **16**, 599–612
458 (2016).
- 459 4. Guindalini, R. S. C. *et al.* Detection of germline variants in Brazilian breast cancer
460 patients using multigene panel testing. *Sci. Rep.* **12**, 4190 (2022).
- 461 5. Shiovitz, S. & Korde, L. A. Genetics of breast cancer: a topic in evolution. *Ann.*
462 *Oncol.* **26**, 1291–1299 (2015).
- 463 6. Melchor, L. & Benítez, J. The complex genetic landscape of familial breast cancer.
464 *Hum. Genet.* **132**, 845–863 (2013).
- 465 7. Khera, A. V. *et al.* Genome-wide polygenic scores for common diseases identify
466 individuals with risk equivalent to monogenic mutations. *Nat. Genet.* **50**, 1219–1224
467 (2018).
- 468 8. Mavaddat, N. *et al.* Polygenic risk scores for prediction of breast cancer and breast
469 cancer subtypes. *Am. J. Hum. Genet.* **104**, 21–34 (2019).
- 470 9. Zhang, H. *et al.* Genome-wide association study identifies 32 novel breast cancer
471 susceptibility loci from overall and subtype-specific analyses. *Nat. Genet.* **52**,
472 572–581 (2020).
- 473 10. Morra, A. *et al.* Association of germline genetic variants with breast cancer-specific
474 survival in patient subgroups defined by clinic-pathological variables related to

- tumor biology and type of systemic treatment. *Breast Cancer Res.* **23**, 86 (2021).
11. Mars, N. *et al.* Genome-wide risk prediction of common diseases across ancestries in one million people. *Cell Genomics* **2**, None (2022).
12. Salzano, Freire-Maia, F. M. N. As origens. in *Populações Brasileiras: Aspectos Demográficos, Genéticos e Antropológicos* (1967).
13. Souza, A. M. de, Resende, S. S., Sousa, T. N. de & Brito, C. F. A. de. A systematic scoping review of the genetic ancestry of the Brazilian population. *Genet. Mol. Biol.* **42**, 495–508 (2019).
14. Naslavsky, M. S. *et al.* Whole-genome sequencing of 1,171 elderly admixed individuals from São Paulo, Brazil. *Nat. Commun.* **13**, 1004 (2022).
15. Sudlow, C. *et al.* UK Biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* **12**, e1001779 (2015).
16. Pedersen, B. S. *et al.* Somalier: rapid relatedness estimation for cancer and germline studies using efficient genome sketches. *Genome Med.* **12**, 62 (2020).
17. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).
18. Rubinacci, S., Ribeiro, D. M., Hofmeister, R. J. & Delaneau, O. Efficient phasing and imputation of low-coverage sequencing data using large reference panels. *Nat. Genet.* **53**, 120–126 (2021).
19. Bergström, A. *et al.* Insights into human genetic variation and population history from 929 diverse genomes. *Science* **367**, (2020).
20. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and

- richer datasets. *Gigascience* **4**, 7 (2015).
21. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).
22. Kuhn, M. Building Predictive Models in R Using the caret Package. *J. Stat. Softw.* **28**, (2008).
23. R Foundation for Statistical Computing. *R: A Language and Environment for Statistical Computing*. (<https://www.R-project.org/>, 2023).
24. Aragon, T. J., Fay, M. P., Wollschlaeger, D. & Omidpanah, A. *epitools: Epidemiology Tools. Tools for training and practicing epidemiologists including methods for two-way and multi-way contingency tables*. (CRAN, 2020).
25. Hu, C. *et al.* A Population-Based Study of Genes Previously Implicated in Breast Cancer. *N. Engl. J. Med.* **384**, 440–451 (2021).
26. Liu, C. *et al.* Generalizability of polygenic risk scores for breast cancer among women with european, african, and latinx ancestry. *JAMA Netw. Open* **4**, e2119084 (2021).
27. Wasik, K. *et al.* Comparing low-pass sequencing and genotyping for trait mapping in pharmacogenetics. *BMC Genomics* **22**, 197 (2021).
28. Li, J. H., Mazur, C. A., Berisa, T. & Pickrell, J. K. Low-pass sequencing increases the power of GWAS and decreases measurement error of polygenic risk scores compared to genotyping arrays. *Genome Res.* **31**, 529–537 (2021).
29. Chaubey, A. *et al.* Low-Pass Genome Sequencing: Validation and Diagnostic Utility from 409 Clinical Cases of Low-Pass Genome Sequencing for the Detection of Copy Number Variants to Replace Constitutional Microarray. *J. Mol. Diagn.* **22**,

- 521 823–840 (2020).
- 522 30. Bahcall, O. Common variation and heritability estimates for breast, ovarian and
523 prostate cancers. *Nat. Genet.* (2019) doi:10.1038/ngicogs.1.
- 524 31. Michailidou, K. *et al.* Association analysis identifies 65 new breast cancer risk loci.
525 *Nature* **551**, 92–94 (2017).
- 526 32. Monticciolo, D. L., Newell, M. S., Moy, L., Lee, C. S. & Destounis, S. V. Breast
527 Cancer Screening for Women at Higher-Than-Average Risk: Updated
528 Recommendations From the ACR. *J. Am. Coll. Radiol.* (2023)
529 doi:10.1016/j.jacr.2023.04.002.