

Type of the Paper: Article

Title: HIV and Pregnancy among Adolescents in Sub-Saharan Africa: A Scoping Review

Folahanmi Tomiwa Akinsolu^{1,2*}, Ifeoluwa Eunice Adewole¹, Abisola Lawale¹, Mobolaji Timothy Olagunju³, Olunike Rebecca Abodunrin^{1,3}, Oluwabukola Mary Ola¹, Abel Nnamdi Chukwuemeka¹, Aisha Oluwaseun Gambari^{1,2}, George Uchenna Eleje⁷, and Oliver Chukwujekwu Ezechi^{1,2}

¹Lead City University, Ibadan

²Nigerian Institute of Medical Research, Lagos

³Nanjing Medical University, Jiangning, Jiangsu Province, China.

⁴University of Cape Town, South Africa

⁵Health Economics Unit, School of Public Health and Family Medicine, Faculty of Health Sciences, University of Cape Town

⁶Department of Medical Microbiology, Babcock University Teaching Hospital, Ogun State, Nigeria

⁷Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria

Folahanmi Tomiwa. Akinsolu^{1,2,*},

¹Lead City University, Ibadan

²Nigerian Institute of Medical Research, Lagos

Email: akinsolu.folahanmi@lcu.edu.ng, Folahanmi.tomiwa@gmail.com

ORCID: 0000-0002-6782-9820

Abisola Lawale¹,

¹Lead City University, Ibadan

Email: abisola.lawale@gmail.com

ORCID: 0009-0006-2449-5433

Ifeoluwa Eunice Adewole¹,

¹Lead City University, Ibadan

Email: theifeoluwa@gmail.com

ORCID: 0000-0003-3992-1205

Olunike Rebecca Abodunrin^{1,3},

¹Lead City University, Ibadan

³Nanjing Medical University, Jiangsu Province, China.

Email: abodunrinolunike@gmail.com

ORCID: 0009-0000-5258-1220

Mobolaji Timothy Olagunju³,
³Nanjing Medical University, Jiangsu Province, China.

Email: olagunjumobolaji@gmail.com

ORCID: 0009-0001-5064-9444

Oluwabukola Mary Ola¹

¹Lead City University, Ibadan

Email: olaoluwabukolamary@gmail.com

ORCID: 0009-0005-8766-2852

Abel Nnamdi Chukwuemeka¹

¹Lead City University, Ibadan

Email: adoncollion@gmail.com

ORCID: 0000-0003-1461-5711

Aisha Oluwaseun Gambari^{1,2}

¹Lead City University, Ibadan

²Nigerian Institute of Medical Research, Lagos

Email: ayeeshagambari@gmail.com

George Uchenna Eleje⁷,

⁷Nnamdi Azikiwe University Teaching Hospital (NAUTH), Nnewi, Nigeria,

Email: elejegeorge1@gmail.com

Oliver Chukwujekwu Ezechi^{1,2},

¹Lead City University, Ibadan

²Nigerian Institute of Medical Research, Lagos

Email: oezechi@nimr.gov.ng, oezechi@yahoo.co.uk

ORCID: 0000-0003-0688-3898

*Corresponding Author

Ifeoluwa Eunice Adewole

theifeoluwa@gmail.com (I.A)

ORCID: 0000-0003-3992-1205

ABSTRACT

The interaction of adolescent pregnancy with HIV is a complex and crucial public health issue worldwide. This is evident in sub-Saharan Africa, where the co-occurrence of HIV and adolescent pregnancies represents a high risk for additional health and social consequences for both adolescent mothers and their children. This study aims to explore literature to provide a clear synthesis of data related to pregnancies among adolescents living with HIV in Sub-Saharan Africa.

A literature search was conducted in July 2023 using PubMed, Scopus, CINAHL, Cochrane Library, African Journals Online, and World Health Organization Global Index Medicus. Two authors screened potential studies. Articles selected were those published in or translated into the English language between 2002 and 2022.

Among 1,560 identified results, 25 peer-reviewed publications on pregnancy among adolescents living with HIV in Sub-Saharan Africa were included after screening. The studies, included both qualitative and quantitative data, with a total of 178,227 participants in the age range of 12-18 years. The majority of the studies reported quantitative data, while a few presented qualitative or mixed-methods findings.

This scoping review highlights disparities in pregnancy prevalence, revealing varied rates among different age groups. The underrepresentation of adolescents in demographic studies suggests diverse vulnerabilities and access to healthcare. The study also emphasizes the need for tailored healthcare services and comprehensive interventions addressing HIV risk, mental health challenges, and socio-economic barriers.

Keywords: Adolescent, Pregnancy, Sub-Saharan Africa, Human Immunodeficiency Virus

INTRODUCTION

Global estimates show that every year, approximately 21 million pregnancies and 12 million births occur among adolescents in low- and middle-income countries [1]. Sub-Saharan Africa is responsible for some of the highest rates of adolescent pregnancies globally [2], with almost one in five females experiencing pregnancy between the ages of 10 and 19 [3].

Adolescent parents and pregnant adolescents face the challenge of managing pregnancy, parenting, and child-rearing responsibilities amid a comprehensive developmental phase often characterized by substantial psychological, social, and biological changes [4].

Furthermore, adolescents make up a large proportion of the people living with HIV globally (1.18 million-2.19 million). In 2022, 140,000 adolescents aged 10 to 19 were newly infected with HIV, with girls accounting for about 80% of them [5]. Particularly in Sub-Saharan Africa, nearly six times the number of newly acquired HIV infections occur among adolescent girls.

Adolescents are considerably more vulnerable to HIV infection compared to adults, given their ongoing developmental stage and the necessity to adapt to rapid biological, physical, and structural changes in their lives [5, 6]. The interaction of adolescent pregnancy with HIV is a complex and crucial public health issue worldwide. This is evident in sub-Saharan Africa, where the co-occurrence of HIV and adolescent pregnancies represents a high risk for additional health and social consequences for both adolescent mothers and their children [7].

With the high rates of early motherhood and HIV incidence in sub-Saharan Africa, designing targeted interventions, enhancing reproductive health services, and encouraging improved overall health outcomes are largely hinged on having this clear understanding of pregnancies among adolescents living with HIV in Sub-Saharan Africa. Thus, this scoping review aims to

- 123 explore existing literature to provide a clear synthesis of data related to pregnancies among
- 124 adolescents living with HIV in Sub-Saharan Africa.

MATERIALS AND METHODS

Overview

This review was guided by the methodological framework of Arksey and O'Malley [8]. Thus, the five steps followed were: (i) identifying the research question, (ii) identifying relevant studies, (iii) selecting eligible studies, (iv) charting the data, and (v) collating and summarizing the results. Appraisal of data quality was not done as the intention was to map all research activities in the field. The preferred reporting items for systematic reviews and meta-analyses extension for scoping reviews (PRISMA-ScR) checklist [9] was utilized with mapping done using the PRISMA-P chart [10].

We conducted a scoping review to synthesize all available evidence on pregnancies among adolescents aged 10 to 19 living with HIV in various Sub-Saharan African countries. A comprehensive search was conducted across major electronic databases, including PubMed, Scopus, CINAHL, Web of Science, and the World Health Organization Global Index Medicus, for studies published from 2002 up to December 2022. The scoping review adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews (PRISMA-ScR) guidelines [10] to ensure methodological rigour and transparency.

Articles published in English and reporting original data on pregnancies among adolescents living with HIV in Sub-Saharan Africa were included. Study selection and data extraction were conducted independently by four reviewers (O.O, O.A, A.L and I.A) to ensure the reliability of the findings. Conflicts were resolved by a fifth reviewer (F.T).

Articles Identification

The initial search was conducted in July 2023 across major electronic databases, including PubMed, Scopus, Google Scholar, and the World Health Organization Global Index Medicus, for studies published from 2002 up to December 2022. The search was performed using the following search strategies enumerated in Appendix 1, combining keywords and Medical Subject Heading (MeSH) terms: “pregnancy”, “adolescents”, “HIV”, and “Sub-Saharan Africa”. The search strategy was refined to further check the appropriateness of the keywords and was expanded to include “females”, “prevalence”, “SSA”, “teenage pregnancy”, and “teenager”. A hand search was also conducted on the references of the included studies and websites of various postgraduate and undergraduate medical colleges to identify potentially relevant literature. The search and screening details are summarized with a graphical presentation using the PRISMA flow diagram in Figure 1. No protocol was published for this review.

The identification and analysis of data in this review was based on the Population/Exposure/Outcome (PEO) framework [11]. This framework was used to structure the review process by identifying key components of the main research question, which are population, exposure and outcome. In this review, the population of interest were adolescents living with HIV in Sub-Saharan Africa, the exposure is HIV status among adolescents in Africa, and the outcome is the prevalence of pregnancy among adolescents living with HIV in Africa.

Search strategy

To achieve our study objectives, a comprehensive search was conducted across major electronic databases, including PubMed, Scopus, Google Scholar, and the World Health Organization Global Index Medicus, for studies published from 2002 up to December 2022.

In this scoping review, we searched for studies that reported pregnancy among female adolescents living with HIV in Sub-Saharan Africa. The search was conducted on studies published between 2002 and 2022 to obtain recent data on pregnancies among female adolescents living with HIV in Sub-Saharan Africa and its associated risk factors (Appendix 1).

Eligibility Criteria and Article Selection

The literature obtained through database searches was imported into the reference management website, Rayyan.ai. In Rayyan, duplicates were removed using the "duplicates items" function. Two independent reviewers (M. O and I.A) conducted title and abstract screening, following the eligibility criteria set for this review, according to the PEO framework [11]. Full-text review of the remaining publications was then completed independently by four researchers (O.O, O.A, A.L and I.A) with reasons for inclusion or exclusion noted. Conflicts were resolved by F.T. Inclusion criteria required that the publications be in English and have full texts available to extract all relevant information. Studies were excluded if they were grey literature, theses, abstracts only or published before January 2002. Studies published between January 2002 and December 2022 were considered for inclusion. The titles and abstracts were used to screen the articles. Studies that do not report the following were excluded: pregnancy in adolescents living with HIV in a Sub-Saharan African Country; HIV status; sample size; lack primary data or well-

defined methodology; duplicated samples and case-control studies and clinical trials; participants not adolescents; non-Sub-Saharan African women of known HIV status.

Data Charting

From the publications included in this review, information on the paper identifiers (title, author, link), the country, year of publication, study aim, study design, study location, HIV status, adolescent pregnancy, and study findings were extracted. The extracted information from each publication was compiled and summarized in Table 1.

RESULTS

Socio-demographic Characteristics of Participants

Figure 1 provides a Prisma Flow diagram for the study. Of the 1,560 results identified through databases (n=1,533) and hand-searching (n=27), 68 duplicates were removed, title and abstract screening for keywords relating to the topic of interest was undertaken for all articles and 52 documents were identified for full-text review. Ultimately, 27 full-text documents were excluded due to lack of relevance, a lack of data from relevant geographical locations, or not reporting on the age range of interest. In total, 25 peer-reviewed publications reported data on pregnancy among HIV-positive adolescent mothers (10–19 years) that spoke to the domains of focus (Figure 1).

Characteristics of Included Studies

The studies that were included reported qualitative and quantitative data on the experiences of adolescent mothers affected by HIV from sub-Saharan African countries: 12 studies from South Africa [4,11,12,13,15,22,23,24,25,26,29,32], 2 from Malawi [17,21], 1 from Zimbabwe [16], 6 Kenyan studies [18,20,27,31,33,34] (n=6), 1 from Lesotho [28] 1 from Uganda [19], and 2 multi-country studies [14, 30].

The majority of the studies reported on quantitative data [4,11,12,14,15,17,18,19,20,22,24,26,28,29,31,32,33,34], five reported qualitative data [13,21,25,27,30], and one study used mixed methods [16]. The total number of participants included is 178,227, and the age range is between 12-18 years.

214 **Table 1: Summary of included studies**

S/ N	Title	Reference	Country	Aim	Study type	Adolescent Pregnancy Prevalence	HIV Prevalence	Key Findings
1	Prevalence and Risk Factors of Repeat Pregnancy among South African Adolescent Females	Govender et al, 2019 [12]	South Africa	This study set out to establish the prevalence and risk factors of repeat pregnancies among South African adolescent females.	Quantitative study	100%	17.8%	<ul style="list-style-type: none"> • 19.9% had experienced a repeat pregnancy. • The risk factors include a history of spontaneous abortion, previous contraceptive use. • A higher level of education and emotional support from family were found to be significant protective factors

								against adolescent repeat pregnancy.
2	Prevalence of repeat pregnancies and associated factors among teenagers in KwaZulu-Natal, South Africa	Mphatswe et al, 2016 [13]	South Africa	To determine the prevalence of repeat teenage pregnancy and the interval between first/most recent and repeat pregnancies, as well as to evaluate the sexual/reproductive health characteristics of teenagers with repeat pregnancies.	Quantitative study	100%	13.3%	<ul style="list-style-type: none"> 82.4% were seen for a first pregnancy and 17.6% for a repeat pregnancy. Only 17.0% participants had previously used contraception 93.1% of whom stopped within 12months and 8.2% had used emergency contraception.

3	The Needs of HIV-positive Teenage Mothers within a South African socio-cultural Context: Experiences of social service providers	Slabbert et al, 2022 [14]	South Africa	To explore and describe the experiences of social service providers in supporting the basic needs of HIV-positive teenage mothers in a rural area in South Africa	Qualitative Study			<ul style="list-style-type: none"> • HIV positive teenage mothers have several needs and lack basic human needs and services • Socio-cultural factors such as early marriage, lack of parental guidance, and gender power imbalances also contribute to teenage pregnancies and positive HIV status.
4	Adolescent Mothers in	Groves et al, 2021	Eswatini, Kenya,	To examine the intersection between	Quantitative Study	17.4%	3.3%	<ul style="list-style-type: none"> • Nearly one fifth of adolescent girls were

	Eastern and Southern Africa: An Overlooked and Uniquely Vulnerable Subpopulation in the Fight Against HIV	[15]	Lesotho, Malawi, Mozambique, South Africa, Tanzania, Uganda, Zambia, and Zimbabwe.	adolescent motherhood and HIV infection across 10 Eastern and Southern African countries, where over half of all HIV infections occur among adolescent girls.				<p>mothers, and the HIV prevalence among all adolescent girls was 3.3%.</p> <ul style="list-style-type: none"> Relative to nonmothers, adolescent mothers were, on average, older, poorer, and more likely to be married, rural dwellers, and household heads. Adolescent motherhood was positively associated with HIV infection.
5	The next generation:	Anderson et al, 2021	South	To describe characteristics of	Quantitative Study	38.8% (≤ 16 years)	100%	<ul style="list-style-type: none"> 258 pregnancies were identified, with 38.8%,

Pregnancy in adolescents and women living with perinatally acquired HIV in South Africa	[16]	Africa	pregnant South African (SA) YLPHIV and their pregnancy outcomes.		39.1% (17 - 19 years)		<p>39.1%, and 22.1% occurring in young females aged ≤ 16 years, 17–19 years, and >20 years.</p> <ul style="list-style-type: none"> Antiretroviral therapy initiation was observed in 84.9% before pregnancy, 6.6% during pregnancy, and 8.5% by pregnancy end. Outcomes revealed that 80.3% resulted in live births, 14.4% in terminations, and 2.2% in stillbirths, with a 2.2% mother-to-child transmission rate.
---	------	--------	--	--	-----------------------	--	--

6	Pregnancy and childbearing among Adolescents and Young Persons Living with HIV (YPLWH): The Case of Chitungwiza, Harare	Makunika, 2021 [17]	Zimbabwe	To highlight the sexual and reproductive health issues affecting young people living with HIV pertaining to childbearing and pregnancy.	Mixed Methods Study		100%	<ul style="list-style-type: none"> sexual and reproductive health (SRH) knowledge heavily impacts YPLWH's decision to have children. Status disclosure has direct impact on decision for condom use.
7	HIV, sexual violence, and termination	Bossard et al, 2022 [18]	Malawi	To assess sexually transmitted infection (STI) prevalence,	Quantitative	52.3%	14.7%	<ul style="list-style-type: none"> The prevalence of HIV, other STIs, unwanted pregnancy, unsafe abortion,

	of pregnancy among adolescent and adult female sex workers in Malawi: A respondent-driven sampling study			progress towards UNAIDS 90-90-90 HIV targets, and SRH care behaviours in a region with established HIV and SRH services tailored to the distinct needs of Female Sex Worker (FSWs).				and sexual violence are extremely high. <ul style="list-style-type: none"> • Peer-led approaches contributed to levels of ART coverage and HIV status awareness similar to those found in the general district population, despite the challenges and risks faced by FSWs.
8	Prevalence of Pregnancy among Adolescent	Awuor, 2021 [19]	Kenya	To find out the prevalence of pregnancy in adolescent living	Quantitative Study	25.5%	100%	<ul style="list-style-type: none"> • 25.5% had pregnancy history in which, 10.2% had knowledge of PMTCT, while only 4.1% were

	Living with HIV in Muhoroni Sub County Kisumu County, Kenya			with HIV, review evidence about Antenatal Care (ANC) service delivery and outcomes for HIV-infected pregnant adolescents in Muhoroni Sub County				<p>having knowledge of PNS</p> <ul style="list-style-type: none"> • All the pregnant girls had knowledge on both ANC and Drug adherence • 10.2% of pregnancies were planned while 15.3% were unplanned which 10.2% of pregnancies were aborted, and 11 (11.2%) were delivered safely, while 4 (4.1%) of the girls were currently pregnant.
9	Adolescent Motherhood and HIV in South Africa:	Roberts, 2021 [4]	South Africa	To identify the prevalence of probable common mental disorder	Quantitative Study	15.2%	70.5%	<ul style="list-style-type: none"> • Results identify higher mental health burden among adolescent mothers

	Examining Prevalence of Common Mental Disorder			among adolescent mothers and, among adolescents experiencing the syndemic of motherhood and HIV.				compared to never-pregnant adolescents • An increased prevalence of mental health burden among adolescent mothers living with HIV compared to other groups. • An elevated prevalence of mental health comorbidities among adolescent mothers irrespective of HIV status.
10	Prevalence and predictors for unintended	Napyo et al, 2020 [20]	Uganda	To determine the prevalence and predictors for unintended pregnancy among	Quantitative Study	6.8%	100%	• The prevalence of unintended pregnancy was 41.1%. • The predictors for

	pregnancy among HIV-infected pregnant women in Lira, Northern Uganda: a cross-sectional study			HIV-infected pregnant women in Northern Uganda				unintended pregnancy were: being single (not living with a partner or being in a marital union), having five or more children and taking antiretroviral drugs for long periods of time
11	Maternal Health Care Utilization Among HIV-Positive Female	Harriet Birungi et al 2011 [21]	Kenya	To examine maternal health care utilization among HIV-positive female adolescents enrolled in HIV/AIDS	Quantitative Study	51.9%	100%	<ul style="list-style-type: none"> The use of PMTCT services was less common than prenatal care services among HIV-positive female adolescents. HIV-positive adolescents

	Adolescents in Kenya			programs in Kenya.				were less likely to use maternal health care services for higher-order pregnancies than for lower-order pregnancies. They were, however, more likely to receive prenatal care and PMTCT services when their husband, rather than someone else, was responsible for the pregnancy.
12	“I would love if there was a young woman to	Carbone et al 2109 [22]	Malawi	examining barriers to, and facilitators of, PMTCT care for HIV-infected	Qualitative Study	100%	100%	<ul style="list-style-type: none"> HIV-infected adolescent mothers expressed a preference for peer-led, non-judgmental PMTCT

encourage us, to ease our anxiety which we would have if we were alone”:	Adapting the Mothers2Mo thers Mentor Mother Model for adolescent mothers living with HIV in			AGYW in Malawi, and explore strategies for adapting the mothers2mothers (m2m) Mentor Mother Model to better meet AGYW service delivery- related needs and preferences				support services that bridge communities and facilities to pragmatically address barriers of stigma, poverty, health system complexity, and food insecurity.
--	---	--	--	---	--	--	--	---

	Malawi							
13	Early adolescent pregnancy increases risk of incident HIV infection in the Eastern Cape, South Africa: a longitudinal study	Christofide et al 2013 [23]	South Africa	To investigate the hypothesis that early adolescent pregnancy predicts incident HIV infections over two years of follow-up.	Quantitative Study			<ul style="list-style-type: none"> • Early adolescent pregnancies increase the incidence of HIV among South African women. • The higher risk is associated with sexual risk behaviours such as higher partner numbers and a greater age difference with partners rather than a biological explanation of hormonal changes during pregnancy.
14	Adolescent and young	Fatti et al 2014 [24]	South Africa	To explore risks associated with	Quantitative Study	20.8%	100%	<ul style="list-style-type: none"> • An increasing proportion of pregnant HIV-positive

pregnant women at increased risk of mother-to-child transmission of HIV and poorer maternal and infant health outcomes: A cohort study at public facilities in				pregnancy in young HIV-infected women, we compared mother-to-child transmission (MTCT) of HIV and maternal and infant health outcomes according to maternal age categories.				women in NMBM were young, and they had increased MTCT and poorer maternal and infant outcomes than older women.
--	--	--	--	---	--	--	--	---

	the Nelson Mandela Bay Metropolitan district, Eastern Cape, South Africa							
15.	Autonomy and infant feeding decision-making among teenage mothers in a rural and	Jama et al 2018 [25]	South Africa	To explore autonomy and infant feeding decision-making among teenage mothers in a rural and urban setting in KwaZulu-Natal, South Africa	Qualitative Study	100%		<ul style="list-style-type: none"> • Teenage mothers had knowledge about recommended feeding practices. • Most feeding advice shared by family members was inappropriate, leading to poor infant feeding practices among teenage mothers.

	urban setting in KwaZulu-Natal, South Africa							Returning to school and fear of breastfeeding in public were also barriers to exclusive breastfeeding.
16	“Motherhood Is Hard”: Exploring the Complexities of Unplanned Motherhood Among HIV-Positive Adolescents in South Africa	Josephine 2019 [26]	South Africa	To examine the complexities of unplanned motherhood among HIV-positive adolescent mothers and their coping strategies through drawing out aspects of their personal experiences	Qualitative Study	100%	100%	<ul style="list-style-type: none"> HIV-positive adolescent mothers perceive unplanned motherhood as difficult and this negatively affects their future childbearing decisions.

17	A longitudinal cohort study of rural adolescent vs adult South African mothers and their children from birth to 24 months	Roux et al 2019 [27]	South Africa	To explore risk factors associated with unplanned teenage pregnancy in Zomba district, Malawi	Quantitative Study	17%	6%	<ul style="list-style-type: none"> • In rural South Africa, almost one in five pregnant women is an adolescent. • Caretaking tasks influencing child growth, especially breastfeeding and securing the child grant appear as the greatest problems for adolescent compared to adult mothers.
18	“If the mother does not know, there is no way she can	Luseno et al 2019 [28]	Kenya	To explore the challenges in promoting the health of these adolescents and preventing	Qualitative study	100%	100%	<ul style="list-style-type: none"> • Adolescents frequently reported being alone during testing, experiencing fear and denial on receiving their results, and delaying

<p>tell the adolescent to go for drugs”:</p> <p>Challenges in promoting health and preventing transmission among pregnant and parenting Kenyan adolescents living with HIV</p>			<p>onward transmission</p>				<p>disclosure to family and linkage to treatment.</p> <ul style="list-style-type: none"> • They also mentioned a lack of contraceptive counseling, with some reporting multiple pregnancies. • Providers voiced misgivings and uncertainty about disclosing HIV diagnoses to minor adolescents without a family member present and reported severe shortages of personnel and resources to adequately serve ALHIV in rural clinics
--	--	--	----------------------------	--	--	--	--

19	HIV incidence among pregnant and postpartum women in a high prevalence setting	Machekan o et al 2018 [29]	Lesotho	To estimate HIV incidence during pregnancy and postpartum and factors associated with HIV acquisition in a cohort of HIV-negative women presenting for antenatal care services at selected health facilities in Lesotho	Quantitative Study	21.3%	4.4%	<ul style="list-style-type: none"> The estimated HIV incidence rate during pregnancy was almost double the estimated HIV incidence during postpartum.
20	Adolescent pregnancy and	Mchunu et al 2016 [30]	South Africa	To assess the prevalence of adolescent	Quantitative Study	19.2%	8.5%	<ul style="list-style-type: none"> 19.2% said that they had an adolescent pregnancy, while 5.8% of male youth

	associated factors in South African youth			<p>pregnancy and associated factors in the South African context, as part of a population-based household survey that formed part of an evaluation of the impact of loveLife, South Africa's national HIV prevention campaign for young people.</p>				<p>indicated that they had impregnated a girl when they were an adolescent.</p> <ul style="list-style-type: none"> • 16.2% of the women indicated that they ever had an unwanted pregnancy and 6.7% had ever terminated a pregnancy. • It was found that being employed or unemployed, greater poverty, having higher sexually permissive attitudes and scoring higher on the contraceptive or the condom use index were associated with adolescent
--	---	--	--	---	--	--	--	---

								pregnancy, and among men wanting the pregnancy and having a sense of the future were associated with adolescent pregnancy.
21	Adolescent pregnancy at antiretroviral therapy (ART) initiation: a critical barrier to retention on ART	Nuwagaba - Biribonwoha et al 2018 [31]	Kenya, Tanzania, and Uganda	To explore and quantify the impact and interactions between age, pregnancy, and site-level factors on LTFU among Adolescents Initiating ART in East Africa		23%	100%	<ul style="list-style-type: none"> Older adolescents experienced higher risk of LTFU compared to YA and adults.

22	Factors associated with unintended pregnancy, poor birth outcomes and post-partum contraceptive use among HIV-positive female adolescents in Kenya	Obare et al 2012 [32]	Kenya	To examine the factors associated with experiencing unintended pregnancies, poor birth outcomes, and post-partum contraceptive use among HIV-positive female adolescents in Kenya.	Quantitative Study	49.4%	100%	<ul style="list-style-type: none"> The experience of repeat unintended pregnancies among HIV-positive female adolescents in the sample is partly due to inconsistent use of contraception to prevent recurrence while poor birth outcomes among higher order pregnancies are partly due to abortion.
23	Adolescent	Ramraj et	South	To quantify the risk	Quantitative	100%	100%	<ul style="list-style-type: none"> Adolescent mothers,

	Access to Care and Risk of Early Mother-to-Child HIV Transmission	al 2018 [33]	Africa	of early mother-to-child transmission (MTCT) of HIV among adolescents enrolled in nationally representative MTCT surveillance studies in South Africa.	Study			<p>compared with adult mothers, have almost three times less planned pregnancies 14.4%</p> <ul style="list-style-type: none"> Gestational age at first antenatal care booking was the only significant predictor of early MTCT among adolescents.
24	Teen pregnancy in rural western Kenya: a public health issue	Omoro et al 2018 [34]	Kenya	To determine among western Kenyan teenage girls: (1) the proportion who reported a history of pregnancy and the	Quantitative Study	23.3%	4.6%	<ul style="list-style-type: none"> Over a third (37.2%) were sexually active and 23.3% had ever been pregnant. Adjusted odds of reporting a history of pregnancy were

				<p>proportion who reported an unwanted pregnancy in the last 12 months; (2) demographic and risk factors associated with those who reported a history of pregnancy and demographic and risk factors associated with those who reported an unwanted pregnancy; and (3)</p>				<p>greater for older compared to younger teens, teens who were ever married or cohabiting compared to those who were single, teens with a primary education or less compared to those with a higher level of education, and teens who experienced partner violence in the last 12 months.</p>
--	--	--	--	---	--	--	--	---

				contraceptive methods used by those with an unwanted pregnancy in the last 12 months.				
25	Gaps in Adolescent Engagement in Antenatal Care and Prevention of Mother-to-child HIV Transmission Services in	Ronen et al 2017 [35]	Kenya	To compare engagement in the PMTCT cascade between adult and adolescent mothers in Kenya.		12.8%	4.2%	<ul style="list-style-type: none"> • Adolescents were less likely than adults to have intended pregnancy disclose their HIV status. Adolescents were less likely than adults to attend ≥ 4 ANC visits. • Among 2359 women without previous HIV testing, 96.1% received testing during pregnancy;

	Kenya							testing levels did not differ between adolescents and adults. Adolescents were less likely than adults to be on ARVs.
--	-------	--	--	--	--	--	--	---

Prevalence of Adolescent Pregnancies among those with HIV in Sub-Saharan Africa

Generally, the included studies consistently highlight a positive association between adolescent motherhood and HIV infection [12, 14, 22, 27]. Specifically, young women who underwent early adolescent pregnancy showed a higher likelihood of later acquiring HIV.

In a South African study carried out among 326 pregnant/parenting adolescents attending prenatal clinics and postnatal and gynecology wards, HIV prevalence was 17.8% [12]. Another study in South Africa, where 341 teenagers were interviewed, reported 82.4% having their first pregnancy and 17.6% having a repeat pregnancy. Notably, a majority of the teenagers with repeat pregnancies conceived again within 24 months of the initial or most recent pregnancy [13]. A study that tested the association between adolescent motherhood and HIV across 10 countries in Eastern and Southern Africa showed that nearly one in five (17.4%) of the adolescent girls included in the analysis were mothers and 5.8% were adolescent mothers with HIV [15].

In South Africa, the prevalence of pregnant adolescent girls who are living with HIV increased every year. In 2016, 38.8% of pregnancies occurred in adolescents ≤ 16 years, while 39.1% were within the 17-19 age bracket [16]. A comparable trend was observed in Zimbabwe, where 33% of young people living with HIV, aged 15 to 24, had experienced pregnancy [17]. Conversely, a study in Malawi focused on female sex workers (FSWs) revealed that 14.7% of them were between the ages of 13–19 and were living with HIV. Additionally, among the 363 participants, 52.3% had encountered their first pregnancy before turning 19 years old [18].

In studies carried out in Kenya, the highest prevalence of pregnancy in adolescents living with HIV was observed between the ages of 17 to 19 years [19,33]. Other studies conducted within

the country reported that the majority of the pregnancies were reported by adolescents aged 18 or older, although most of the pregnancies had occurred when the respondents were 17 years and below [21,32]. Similarly, a longitudinal study carried out in South Africa over two years of follow-up reported that two hundred and fifty-one young women (17.74%) experienced a pregnancy when they were under the age of 19 [23].

Within Uganda, a study involving 518 HIV-positive pregnant women revealed that 6.8% were under the age of 20 [20]. In Kenya, a sampling of 498 HIV-positive women from both primary national and secondary Nyanza oversample surveys found that 21 individuals (4.2%) were adolescents [34].

Socioeconomic factors were also found to be associated with adolescent pregnancy. Low-income earners, those with no income, and those with lower levels of education were more likely to have early pregnancy in comparison to their counterparts [12,16,20,29,33,34]. They also experienced more physical and sexual violence [22, 26,33].

Moreover, the impact of early childbirth extended to subsequent pregnancies among adolescents living with HIV. Studies indicated that having children at a young age significantly influenced the likelihood of repeat pregnancies among this demographic [11, 12]. As the age of adolescents increased, so did the odds of experiencing repeated pregnancies, suggesting a correlation between age and the occurrence of repeat pregnancies.

Impact of Pregnancies on Clinical Outcomes and Well-being

Poor mental health has been found to be higher among adolescent mothers. Probable common mental disorder was higher among adolescent mothers living with HIV (23.0%) in comparison to non-mothers and mothers who were not living with HIV. Adolescent mothers were shown to

have higher prevalence of probable mental health comorbidities regardless of their HIV status, indicating that the experience of pregnancy and motherhood may contribute to a heavier burden on their mental health [4].

Qualitative data revealed that adolescent mothers living with HIV often cannot remain at home with their parents, either for reasons of overcrowding in the home, abuse, neglect, or financial difficulties. They are often averse to pregnancies because it brings emotional and physical burden to the already burdened person [14, 17]. Adolescent mothers' safety needs are not met. Some of them have had to stay on their own with their babies or move to unsafe communities because of the threat of violent acts against them [14].

Reports from included studies show that HIV-infected pregnant adolescents have challenges with long-term contraceptive continuation and consequently poorer prevention of mother-to-child HIV transmission (PMTCT) service outcomes and higher rates of mother-to-child HIV transmission among infants compared to HIV-infected pregnant adults, likely due to the poor knowledge on care among them [13, 15, 16, 19]. Adolescent mothers have lower adherence to ANC visits and ART treatments, [16,21,31, 34, 35]. This is corroborated by another study which reported a significant number of pregnant adolescents with HIV had severe immunosuppression (CD4 count 200 cells/L) or increased VLs (400 copies/mL) [15]. Adolescents also had increased risks of maternal mortality, first presentation in labour, and stillbirth. [23,31]

Seven studies reported the low use of maternal health care services, including use of PMTCT, skilled attendance and postnatal/postabortion care for pregnancies which might end in adverse outcomes such as miscarriage, stillbirth or abortion among HIV-positive female adolescents [12, 14, 15, 18, 20, 30, 32, 34].

DISCUSSION

The scoping review has presented an extensive synopsis of all available literature on adolescent mothers living with HIV in sub-Saharan Africa through the prevalence of adolescent pregnancies, including the risk factors, impacts on clinical outcomes and well-being, and the availability and effectiveness of reproductive health services.

The prevalence of pregnancies among adolescents living with HIV in Africa has long been a recognized concern. While prior studies have identified the co-occurrence of adolescent pregnancies and HIV, this review delves deeper into the dynamics of prevalence. For example, included studies reported the ages of 17 to 19 years to have the highest prevalence rates of adolescent pregnancy [18,34], while others reported that the majority of the pregnancies had occurred when the respondents were aged 17 years and below [20,22,31]. These disparities could be attributed to differences in healthcare infrastructure, HIV testing, and antiretroviral therapy availability [36,37].

This study sheds light on a significant disparity in the representation of adolescents within certain demographics, particularly in studies involving pregnant individuals living with HIV in South Africa. In both cases cited, adolescents comprised a notably lower percentage compared to their older counterparts. In the first instance, among pregnant adolescents and young pregnant women living with HIV, only 20.8% fell within the adolescent age range (13-19 years), while the older cohort (20-24 years) represented a higher proportion at 32.6%. This disparity indicates a potential discrepancy in the impact of HIV within different age groups, suggesting varying levels of vulnerability or access to healthcare services between adolescents and slightly older women.

Similarly, in the longitudinal study that assessed the challenges faced by adolescent versus adult mothers in rural South Africa during the initial 24 months post-birth, adolescents accounted for only 17% of the mothers. This percentage was notably lower when considering HIV seropositive cases, where only 6% of adolescent mothers were identified as HIV positive. This finding, in contrast to their adult counterparts, highlights a significant difference in HIV prevalence between adolescent and adult mothers. The lower representation of adolescents in these studies might reflect several underlying factors. There could be differences in healthcare-seeking behaviors, access to reproductive health education, or social support systems between adolescents and older individuals. It could also signify varying levels of vulnerability or risk perception among these age groups regarding HIV and pregnancy.

Understanding these disparities is crucial for developing targeted interventions and support systems that address the specific needs and challenges faced by adolescents, especially pregnant adolescents living with HIV. It emphasizes the necessity of tailored healthcare services and comprehensive support frameworks to bridge the gap and ensure equitable care and resources for all demographics, irrespective of age.

The study also showed that HIV prevalence was higher among adolescent mothers in comparison to adolescent non-mothers. This is in contrast to a study in South Africa where adolescent mothers were less likely to be living with HIV compared to never-pregnant adolescents within the sample (55.5% vs. 73.3%). The prevalence of adolescent motherhood in the sample was 15.2%, and almost all (98.8%) pregnancies were unplanned or unwanted [4]. Studies have shown that adolescents were more likely to report having had at least one unwanted pregnancy [17, 29].

Existing studies have shown that early sexual debut is a significant risk factor for both unintended pregnancies and HIV infection due to various factors such as limited knowledge

about safe sex practices, lack of access to contraceptives, and potential engagement in risky sexual behaviours [38,39]. This may also be linked to adolescent mothers learning of their HIV status for the first time when they confirm that they are pregnant [30,40]. This review also highlights that adolescent mothers are more likely to be involved in relationships where they have poor relationship power, gender inequitable norms, intimate partner violence, and little HIV preventive communication with higher-risk partners [43,44, 41,42].

The association between lower education levels and higher repeat pregnancies among adolescents living with HIV, suggests that improving educational opportunities and knowledge about reproductive health may play a crucial role in preventing repeated pregnancies in this vulnerable group [45,46]. Additionally, the role of poverty in early pregnancies is a common thread in many studies. Poverty often limits access to contraceptives, healthcare, and comprehensive reproductive health education [46,47]. Targeted interventions that address economic disparities are imperative. The prevention of adolescent pregnancies and HIV goes beyond healthcare; it involves broader socio-economic and educational support. The findings of this review emphasize the importance of holistic interventions that encompass not only clinical care but also social and economic empowerment.

Mental health challenges among adolescent mothers living with HIV are a shared concern, as evidenced by the high prevalence of probable mental health disorders in this population [4,48]. Stigma, lack of social support, poverty, and the timing of HIV diagnosis may contribute to these mental health challenges [48].

Hindrances in relation to access, utilization, and the quality of services are common themes in this review. Issues such as service availability in rural areas and shortages of well-trained

personnel are shared concerns. Geographic disparities and socio-economic barriers also hinder access to essential care [13,27]. While the South African government provides free services such as counselling, child support grants, food parcels, housing and social grants to adolescent mothers living with HIV, these services are often not accessible to those who stay in rural communities and are far from social welfare offices [13]. Smaller rural clinics, which are often more accessible to these adolescents, also have to deal with the shortage of well-trained personnel and insufficient staffing as a hindrance to achieving effective adolescent care [27].

The included studies underscore notable negligence in meeting the reproductive health needs of individuals transitioning from childhood to adulthood. Often, practical solutions for sexually active adolescents, or those intending to be sexually active, are lacking. Emphasis tends to revolve around advocating abstinence from sexual intercourse without providing comprehensive and practical guidance or support [30, 31].

RECOMMENDATIONS

This study suggests that larger facilities, such as referral hospitals, may be better equipped to offer adolescent-friendly PMTCT services that emphasize confidentiality, address stigma related to HIV infection and adolescent pregnancy, and integrate models of increasing social support. Contact with antenatal and postnatal care services and HIV care services for adolescents living with HIV also provide an opportunity to merge differing branches of health service through providing screening, monitoring, and referral for mental health.

The results from this study also recommend the need for integrated mental health services within reproductive health care programs tailored to the unique needs of adolescent mothers in SSA. Challenges faced by adolescent mothers living with HIV, including accelerated disease

371 progression and heightened mental health issues, were highlighted in this review. Research has
 372 repeatedly identified stigma, lack of social support, parenting stress, and knowledge gaps as
 373 factors contributing to mental health challenges [4,48]. This highlights the necessity for a more
 374 holistic approach, one that goes beyond advocating abstinence and includes comprehensive
 375 reproductive health education, access to contraceptives, support for safer sexual practices, and
 376 tailored interventions for adolescents living with HIV who are navigating parenthood.
 377 Addressing these gaps is vital to ensure the well-being and reproductive health empowerment of
 378 adolescent mothers living with HIV.

Funding: “This research received no external funding.”

Acknowledgement: This work was supported by the Nigerian Institute of Medical Research, Lead City Univeristy, and Fogarty-IeDEA Mentorship Program (FIMP), West Africa.

Data Availability Statement: The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest: “The authors declare no conflict of interest.”

References

1. Sully, Elizabeth. 2020. “Adding It Up: Investing in Sexual and Reproductive Health 2019.” Guttmacher Institute. July 10, 2020. <https://www.guttmacher.org/report/adding-it-up-investing-in-sexual-reproductive-health-2019>.
2. Maharaj, Niren Ray. 2022. “Adolescent Pregnancy in Sub-Saharan Africa – a Cause for Concern.” *Frontiers in Reproductive Health* 4 (December). <https://doi.org/10.3389/frph.2022.984303>.
3. Kassa, Getachew Mullu, A. O. Arowojolu, A. A. Odukogbe, and Alemayehu Worku Yalew. 2018. “Prevalence and Determinants of Adolescent Pregnancy in Africa: A Systematic Review and Meta-Analysis.” *Reproductive Health* 15 (1). <https://doi.org/10.1186/s12978-018-0640-2>.
4. Roberts, Kathryn J., Colette Smith, Lucie Cluver, Elona Toska, Siyanai Zhou, Mark Boyes, and Lorraine Sherr. 2021. “Adolescent Motherhood and HIV in South Africa: Examining Prevalence of Common Mental Disorder.” *AIDS and Behavior* 26 (September). <https://doi.org/10.1007/s10461-021-03474-8>.
5. Unicef. 2022. “Adolescent HIV Prevention - UNICEF DATA.” UNICEF DATA. 2022. <https://data.unicef.org/topic/hivaids/adolescents-young-people/>.
6. Calabrese, Samantha, Sonia Lee, Sonia Lee, Susannah Allison, Gina Brown, Patrick Jean-Philippe, Nahida Chakhtoura, Jack Moye, and Bill G Kapogiannis. 2023. “Adolescent and Young Adult Research across the HIV Prevention and Care Continuum: An International Programme Analysis and Targeted Review.” *Journal of the International AIDS Society* 26 (3). <https://doi.org/10.1002/jia2.26065>.
7. Toska, Elona, Christina A. Laurenzi, Kathryn J. Roberts, Lucie Cluver, and Lorraine Sherr. 2020. “Adolescent Mothers Affected by HIV and Their Children: A Scoping

- Review of Evidence and Experiences from Sub-Saharan Africa.” *Global Public Health* 15 (11): 1655–73. <https://doi.org/10.1080/17441692.2020.1775867>.
8. Arksey H, O’Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005;8(1): 19-32.
9. Tricco AC, Lillie E, Zarin W, O’Brien KK, Colquhoun H, Levac D et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Int Med*, 2018;169(7):467-73)
10. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015; 4(1):1.
11. Elsevier. “Clinical Questions: PICO and PEO Research | Author Services Blog.” Elsevier Author Services - Articles. Elsevier, 2021. <https://scientific-publishing.webshop.elsevier.com/research-process/clinical-questions-pico-and-peo-research/>.
12. Desiree Govender, Saloshni Naidoo and Myra Taylor. “Prevalence and Risk Factors of Repeat Pregnancy among South African Adolescent Females” *Afr J Reprod Health* 2019; 23[1]: 73-87.
13. MphatsweW, et al, Prevalence of repeat pregnancies and associated factors among teenagers in KwaZulu-Natal, South Africa, *Int J Gynecol Obstet* (2016), <http://dx.doi.org/10.1016/j.ijgo.2015.09.028>
14. Ilze Slabbert, Tasneemah Cornelissen-Nordien and Nomfundo Thokozile Singwayo. The needs of HIV-positive teenage mothers within a South African socio-cultural context Experiences of social service providers

15. Groves, Allison K., Luwam T. Gebrekristos, Patrick D. Smith, Kirsten Stoebe, Marie C. Stoner, Wole Ameyan, and Alex C. Ezech. 2022. "Adolescent Mothers in Eastern and Southern Africa: An Overlooked and Uniquely Vulnerable Subpopulation in the Fight against HIV." *Journal of Adolescent Health* 70 (6): 895–901. <https://doi.org/10.1016/j.jadohealth.2021.12.012>.
16. Anderson, K, T Mutemaringa, K-G Technau, L F Johnson, K Braithwaite, E Mokotoane, A Boule, and M-A Davies, on behalf of IeDEA-SA Pediatrics. 2021. "The next Generation: Pregnancy in Adolescents and Women Living with Perinatally Acquired HIV in South Africa." *South African Medical Journal* 111 (3): 260. <https://doi.org/10.7196/samj.2021.v111i3.14987>.
17. Norman, Makunika. 2021. "Pregnancy and Childbearing among Adolescents and Young Persons Living with HIV: The Case of Chitungwiza, Harare." *International Journal of Sexual and Reproductive Health Care*, 067–72. <https://doi.org/10.17352/ijsrhc.000026>.
18. Bossard, Claire, Menard Chihana, Sarala Nicholas, Damian Mauambeta, Dina Weinstein, Nolwenn Conan, Elena Nicco, et al. 2022. "HIV, Sexual Violence, and Termination of Pregnancy among Adolescent and Adult Female Sex Workers in Malawi: A Respondent-Driven Sampling Study." *PLoS ONE* 17 (12): e0279692–92. <https://doi.org/10.1371/journal.pone.0279692>.
19. Silas Onyango Awuor. 2021. "Prevalence of Pregnancy among Adolescent Living with HIV in Muhoroni Sub County Kisumu County, Kenya." *Annals of Obstetrics and Gynecology* 9 (1): 7. <https://doi.org/10.11648/j.ijacm.20210901.12>.
20. Napyo, Agnes, Victoria Nankabirwa, David Mukunya, Josephine Tumuhamy, Grace Ndeezi, Anna Agnes Ojok Arach, Beatrice Odongkara, Paul Waako, Thorkild Tylleskär,

- and James K. Tumwine. 2020. "Prevalence and Predictors for Unintended Pregnancy among HIV-Infected Pregnant Women in Lira, Northern Uganda: A Cross-Sectional Study." *Scientific Reports* 10 (1). <https://doi.org/10.1038/s41598-020-73490-6>.
21. Birungi, Harriet, Francis Obare, Anke van der Kwaak, and Jane Harriet Namwebya. 2011. "Maternal Health Care Utilization among HIV-Positive Female Adolescents in Kenya." *International Perspectives on Sexual and Reproductive Health* 37 (03): 143–49. <https://doi.org/10.1363/3714311>.
22. Carbone, Nicole B., Joseph Njala, Debra J. Jackson, Michael T. Eliya, Chileshe Chilangwa, Jennifer Tseka, Tasila Zulu, et al. 2019. "'I Would Love If There Was a Young Woman to Encourage Us, to Ease Our Anxiety Which We Would Have If We Were Alone': Adapting the Mothers2Mothers Mentor Mother Model for Adolescent Mothers Living with HIV in Malawi." Edited by Mellissa H Withers. *PLOS ONE* 14 (6): e0217693. <https://doi.org/10.1371/journal.pone.0217693>.
23. Christofides, Nicola J, Rachel K Jewkes, Kristin L Dunkle, Mzikazi Nduna, Nwabisa Jama Shai, and Claire Sterk. 2014. "Early Adolescent Pregnancy Increases Risk of Incident HIV Infection in the Eastern Cape, South Africa: A Longitudinal Study." *Journal of the International AIDS Society* 17 (1): 18585. <https://doi.org/10.7448/ias.17.1.18585>.
24. Fatti, Geoffrey, Najma Shaikh, Brian Eley, Debra Jackson, and Ashraf Grimwood. 2014. "Adolescent and Young Pregnant Women at Increased Risk of Mother-To-Child Transmission of HIV and Poorer Maternal and Infant Health Outcomes: A Cohort Study at Public Facilities in the Nelson Mandela Bay Metropolitan District, Eastern Cape, South Africa." *South African Medical Journal* 104 (12): 874. <https://doi.org/10.7196/samj.8207>.

25. Jama, Ngcwalisa Amanda, Aurene Wilford, Lyn Haskins, Anna Coutsooudis, Lenore Spies, and Christiane Horwood. 2018. "Autonomy and Infant Feeding Decision-Making among Teenage Mothers in a Rural and Urban Setting in KwaZulu-Natal, South Africa." BMC Pregnancy and Childbirth 18 (1). <https://doi.org/10.1186/s12884-018-1675-7>.
26. Josephine, Adeagbo Morolake. 2019. "'Motherhood Is Hard': Exploring the Complexities of Unplanned Motherhood among HIV-Positive Adolescents in South Africa." SAGE Open 9 (2): 215824401984880. <https://doi.org/10.1177/2158244019848802>.
27. Roux, Karl le, Joan Christodoulou, Linnea Stansert-Katzen, Elaine Dippenaar, Christina Laurenzi, Ingrid M. le Roux, Mark Tomlinson, and Mary Jane Rotheram-Borus. 2019. "A Longitudinal Cohort Study of Rural Adolescent vs Adult South African Mothers and Their Children from Birth to 24 Months." BMC Pregnancy and Childbirth 19 (1). <https://doi.org/10.1186/s12884-018-2164-8>.
28. Luseno, Winnie K., Bonita J. Iritani, Suzanne Maman, Isabella Mbai, Barrack Ongili, Florence Anyango Otieno, and Denise Dion Hallfors. 2019. "'If the Mother Does Not Know, There Is No Way She Can Tell the Adolescent to Go for Drugs': Challenges in Promoting Health and Preventing Transmission among Pregnant and Parenting Kenyan Adolescents Living with HIV." Children and Youth Services Review 103 (August): 100–106. <https://doi.org/10.1016/j.childyouth.2019.05.036>.
29. Machekano, Rhoderick, Appolinaire Tiam, Seble Kassaye, Vincent Tukei, Michelle Gill, Florence Mohai, Masepeli Nchepe, et al. 2018. "HIV Incidence among Pregnant and Postpartum Women in a High Prevalence Setting." Edited by Javier R. Lama. PLOS ONE 13 (12): e0209782. <https://doi.org/10.1371/journal.pone.0209782>.

30. Mchunu, G, K Peltzer, B Tutshana, and L Seutlwadi. 2013. “Adolescent Pregnancy and Associated Factors in South African Youth.” *African Health Sciences* 12 (4). <https://doi.org/10.4314/ahs.v12i4.5>.
31. Nuwagaba-Biribonwoha, Harriet, Agnes N Kiragga, Constantin T Yiannoutsos, Beverly S Musick, Kara K Wools-Kaloustian, Samuel Ayaya, Hilary Wolf, et al. 2018. “Adolescent Pregnancy at Antiretroviral Therapy (ART) Initiation: A Critical Barrier to Retention on ART.” *Journal of the International AIDS Society* 21 (9): e25178. <https://doi.org/10.1002/jia2.25178>.
32. Obare, Francis, Anke van der Kwaak, and Harriet Birungi. 2012. “Factors Associated with Unintended Pregnancy, Poor Birth Outcomes and Post-Partum Contraceptive Use among HIV-Positive Female Adolescents in Kenya.” *BMC Women’s Health* 12 (1). <https://doi.org/10.1186/1472-6874-12-34>.
33. Ramraj, Trisha, Debra Jackson, Thu-Ha Dinh, Steve Olorunju, Carl Lombard, Gayle Sherman, Adrian Puren, et al. 2018. “Adolescent Access to Care and Risk of Early Mother-To-Child HIV Transmission.” *Journal of Adolescent Health* 62 (4): 434–43. <https://doi.org/10.1016/j.jadohealth.2017.10.007>.
34. Omoro, Tereza, Simone C. Gray, George Otieno, Calvin Mbeda, Penelope A. Phillips-Howard, Tameka Hayes, Fredrick Otieno, and Deborah A. Gust. 2017. “Teen Pregnancy in Rural Western Kenya: A Public Health Issue.” *International Journal of Adolescence and Youth*, December, 1–10. <https://doi.org/10.1080/02673843.2017.1402794>.

35. Ronen, Keshet, Christine J. McGrath, Agnes C. Langat, John Kinuthia, Danvers Omolo, Benson Singa, Abraham K. Katana, Lucy W. Ngang'A, and Grace John-Stewart. 2017. "Gaps in Adolescent Engagement in Antenatal Care and Prevention of Mother-To-Child HIV Transmission Services in Kenya." JAIDS Journal of Acquired Immune Deficiency Syndromes 74 (1): 30–37. <https://doi.org/10.1097/qai.0000000000001176>
36. Mandiwa, Chrispin, Bernadetta Namondwe, and Mtondera Munthali. 2021. "Prevalence and Correlates of Comprehensive HIV/AIDS Knowledge among Adolescent Girls and Young Women Aged 15–24 Years in Malawi: Evidence from the 2015–16 Malawi Demographic and Health Survey." BMC Public Health 21 (1). <https://doi.org/10.1186/s12889-021-11564-4>.
37. Mabaso, Musawenkosi, Goitseone Maseko, Ronel Sewpaul, Inbarani Naidoo, Sean Jooste, Sinovuyo Takatshana, Tarylee Reddy, Khangelani Zuma, and Nompumelelo Zungu. 2021. "Trends and Correlates of HIV Prevalence among Adolescents in South Africa: Evidence from the 2008, 2012 and 2017 South African National HIV Prevalence, Incidence and Behaviour Surveys." AIDS Research and Therapy 18 (1). <https://doi.org/10.1186/s12981-021-00422-3>.
38. Dine, Roseline Dzekem, Oluwaseun Adebayo Bamodu, and Joseph Ntaganira. 2021. "Youth Health Risk Behavior: Effects of Early Sexual Debut on HIV Incidence among Rwandan Youth." Journal of Public Health, June. <https://doi.org/10.1007/s10389-021-01617-8>.
39. Ningpuanyeh, Wilson C, and A Sathiya Susuman. 2016. "Correlates of Early Sexual Debut and Its Associated STI/HIV Risk Factors among Sexually Active Youths in

Malawi.” *Journal of Asian and African Studies* 52 (8): 1213–24.
<https://doi.org/10.1177/0021909616654300>.

40. Madiba, Sphiwe, and Mamorapeli Putsoane. 2020. “Testing Positive and Disclosing in Pregnancy: A Phenomenological Study of the Experiences of Adolescents and Young Women in Maseru, Lesotho.” *AIDS Research and Treatment* 2020 (February): 1–8.
<https://doi.org/10.1155/2020/6126210>.

41. Bermudez, Laura Gauer, Drosin Mulenga, Maurice Musheke, and Sanyukta Mathur. 2021. “Intersections of Financial Agency, Gender Dynamics, and HIV Risk: A Qualitative Study with Adolescent Girls and Young Women in Zambia.” *Global Public Health*, July, 1–14. <https://doi.org/10.1080/17441692.2021.1951800>.

42. Crankshaw, Tamaryn L., Samantha Chareka, Pemberai Zambezi, and Nana K. Poku. 2021. “Age Matters: Determinants of Sexual and Reproductive Health Vulnerabilities amongst Young Women Who Sell Sex (16–24 Years) in Zimbabwe.” *Social Science & Medicine* 270 (February): 113597. <https://doi.org/10.1016/j.socscimed.2020.113597>.

43. Bhushan, Nivedita L., Marie C.D. Stoner, Allison K. Groves, Kathleen Kahn, and Audrey E. Pettifor. 2022. “Partnership Dynamics and HIV-Related Sexual Behaviors among Adolescent Mothers in South Africa: A Longitudinal Analysis of HIV Prevention Trials Network 068 Data.” *Journal of Adolescent Health*, March.
<https://doi.org/10.1016/j.jadohealth.2022.02.003>.

44. Groves, Allison K., Luwam T. Gebrekristos, Luz McNaughton Reyes, Dhayendre Moodley, and Suzanne Maman. 2020. “Describing Relationship Characteristics and

Postpartum HIV Risk among Adolescent, Young Adult, and Adult Women in South Africa.” *Journal of Adolescent Health* 67 (1): 123–26.
<https://doi.org/10.1016/j.jadohealth.2019.12.008>.

45. Marphatia, Akanksha A., Naomi M. Saville, Gabriel S. Amable, Dharma S. Manandhar, Mario Cortina-Borja, Jonathan C. Wells, and Alice M. Reid. 2020. “How Much Education Is Needed to Delay Women’s Age at Marriage and First Pregnancy?” *Frontiers in Public Health* 7 (January). <https://doi.org/10.3389/fpubh.2019.00396>.

46. Akombi-Inyang, Blessing Jaka, Emma Woolley, Chinonyerem Ogadi Iheanacho, Khulan Bayaraa, and Pramesh Raj Ghimire. 2022. “Regional Trends and Socioeconomic Predictors of Adolescent Pregnancy in Nigeria: A Nationwide Study.” *International Journal of Environmental Research and Public Health* 19 (13): 8222.
<https://doi.org/10.3390/ijerph19138222>.

47. Worku, Misganaw Gebrie, Zemenu Tadesse Tessema, Achamyeleh Birhanu Teshale, Getayeneh Antehunegn Tesema, and Yigizie Yeshaw. 2021. “Prevalence and Associated Factors of Adolescent Pregnancy (15–19 Years) in East Africa: A Multilevel Analysis.” *BMC Pregnancy and Childbirth* 21 (1). <https://doi.org/10.1186/s12884-021-03713-9>.

48. Dessauvagie, A. S., A. Jörens-Presentati, A.-K. Napp, D. J. Stein, D. Jonker, E. Breet, W. Charles, et al. 2020. “The Prevalence of Mental Health Problems in Sub-Saharan Adolescents Living with HIV: A Systematic Review.” *Global Mental Health* 7.
<https://doi.org/10.1017/gmh.2020.18>.

588 **Appendix: Search Terms**

589 **PubMed- 59**

	Keywords	Search Strings
1	Prevalence	((TS=(Prevalence)) OR TS=(Epidemiology)) OR TS=(Frequency) and Preprint Citation Index (Exclude – Database)
2	Adolescent pregnancy	((((((((((TS=(Adolescence, Pregnancy)) OR TS=(Teen Pregnancy)) OR TS=(Pregnancy, Teen)) OR TS=(Adolescent Pregnancy)) OR TS=(Pregnancy, Adolescent)) OR TS=(Pregnancy, Teenage)) OR TS=(Teenage Pregnancies)) OR TS=(Preteen Pregnancy)) OR TS=(Pregnancy, Preteen)) OR TS=(Child Pregnancy)) OR TS=(Girls pregnancy)) OR TS=(Adolescent mothers)) OR TS=(Pregnancy unplanned)) OR TS=(Pregnancy unwanted) and Preprint Citation Index (Exclude – Database)
3	HIV	((((((((((TS=(Human Immunodeficiency Virus)) OR TS=(Immunodeficiency Virus, Human)) OR TS=(Immunodeficiency Viruses, Human)) OR TS=(Virus, Human Immunodeficiency)) OR TS=(Viruses, Human Immunodeficiency)) OR TS=(Human Immunodeficiency

		Viruses)) OR TS=(Human T Cell Lymphotropic Virus Type III)) OR TS=(LAV-HTLV-III)) OR TS=(Lymphadenopathy-Associated Virus)) OR TS=(Human T Lymphotropic Virus Type III)) OR TS=(AIDS Virus)) OR TS=(Acquired Immune Deficiency Syndrome Virus)) OR TS=(HTLV-III) and Preprint Citation Index (Exclude – Database)

590 **CINAHL= 0**

	Keywords	Search Strings
1	Prevalence	TI (prevalence or incidence or epidemiology or frequency or occurrence or statistics) OR TI prevalence rates
2	Adolescent Pregnancy	TI adolescent pregnancy or teenage pregnancy or teen pregnancy
3	HIV	TI HIV or aids or acquired human immunodeficiency syndrome or human immunodeficiency virus

591

592 **WEB OF SCIENCE**

593 **Web of science (Number)**

S/N	Keywords	
1	Pregnancy	((TS=(Pregnancy)) OR TS=(Pregnancies)) OR TS=(Gestation) and Preprint Citation Index (Exclude – Database)
2	adolescents	((((((((((((((TS=(adolescents)) OR TS=(Adolescence)) OR TS=(Teens)) OR TS=(Teen)) OR TS=(Teenagers)) OR TS=(Teenager)) OR TS=(Youth)) OR TS=(Youths)) OR TS=(Adolescents, Female)) OR TS=(Adolescent, Female)) OR TS=(Female Adolescent)) OR TS=(Female Adolescents)) OR TS=(Adolescents, Male)) OR TS=(Adolescent, Male)) OR TS=(Male Adolescent)) OR TS=(Male Adolescents) and Preprint Citation Index (Exclude – Database)
3	HIV	((((((((((((((((((((((TS=(HIV)) OR TS=(Human Immunodeficiency Virus)) OR TS=(Immunodeficiency Virus, Human)) OR TS=(Immunodeficiency Viruses, Human)) OR TS=(Virus, Human Immunodeficiency)) OR TS=(Viruses, Human Immunodeficiency)) OR TS=(Human Immunodeficiency Viruses)) OR TS=(Human T Cell Lymphotropic Virus Type III)) OR TS=(Human T-Cell Lymphotropic Virus Type III)) OR TS=(Human T-Cell Leukemia Virus Type III)) OR TS=(Human T Cell Leukemia Virus Type III)) OR TS=(LAV-HTLV-III)) OR TS=(Lymphadenopathy-Associated Virus)) OR

		<p>TS=(Lymphadenopathy Associated Virus)) OR</p> <p>TS=(Lymphadenopathy-Associated Viruses)) OR TS=(Virus, Lymphadenopathy-Associated)) OR TS=(Viruses, Lymphadenopathy-Associated)) OR TS=(Human T Lymphotropic Virus Type III)) OR TS=(Human T-Lymphotropic Virus Type III)) OR TS=(AIDS Virus)) OR TS=(AIDS Viruses)) OR TS=(Virus, AIDS)) OR TS=(Viruses, AIDS)) OR TS=(Acquired Immune Deficiency Syndrome Virus)) OR TS=(Acquired Immunodeficiency Syndrome Virus)) OR TS=(HTLV-III) and Preprint Citation Index (Exclude – Database)</p>
4	Sub-Sahara Africa	<p>((TS=(Sub-Sahara Africa)) OR TS=(Sub-Saharan Africa)) OR TS=(Subsaharan Africa)) OR TS=(Africa, Sub-Saharan) and Preprint Citation Index (Exclude – Database)</p>

594

595 **SCOPUS = 462**

596 (TITLE-ABS-KEY(prevalence) AND TITLE-ABS-KEY(adolescent) AND TITLE-ABS-KEY(Pregnancy) AND TITLE-
597 ABS-KEY(HIV) AND TITLE-ABS-KEY(Africa)) AND PUBYEAR > 2001 AND PUBYEAR < 2023 AND (LIMIT-TO (DOCTYPE,"ar")