1	Functional brain connectivity in early adolescence after hypothermia-treated
2	neonatal hypoxic-ischemic encephalopathy
3	
4	Gustaf Håkansson <sup>1,2</sup> *; Katarina Robertsson Grossmann <sup>1,2</sup> ; Ulrika Ådén <sup>1,3</sup> ; Mats Blennow <sup>1,2</sup> ;
5	Peter Fransson <sup>4</sup>
6	
7	<sup>1</sup> Department of pediatrics, Karolinska University Hospital, Stockholm, Sweden.
8	<sup>2</sup> Department of Clinical Intervention and Technology, Karolinska Institutet, Stockholm, Sweden
9	<sup>3</sup> Department of Women's and Children's health, Karolinska Institutet, Stockholm, Sweden
10	<sup>4</sup> Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden
11	
12	*Correspondence to Karolinska University Hospital, K59 Huddinge, 141 86 Stockholm. Email:
13	gustaf.hakansson@ki.se
14	ORCID-id: Gustaf Håkansson 0000-0002-0608-9742
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	

# 1 Abstract

2	Aim: Neonatal hypoxic-ischemic encephalopathy (HIE) injures the infant brain during the basic
3	formation of the developing functional connectome. This study aimed to investigate long-term
4	changes in the functional connectivity (FC) networks of the adolescent brain following neonatal
5	HIE treated with therapeutic hypothermia (TH).
6	Method: This prospective, population-based cohort study included all infants (n=66) with TH-
7	treated neonatal HIE in Stockholm during 2007-2009 and a control group (n=43) of children with
8	normal neonatal course. Assessment with resting-state functional magnetic resonance imaging
9	(fMRI) was performed at Karolinska Institutet, Stockholm at age 9-12 years.
10	Results: fMRI data met quality criteria for 35 children in the HIE-cohort (mean [SD] age at MRI:
11	11.2 [0.74] years, 46% male) and 30 children in the control group (mean [SD] age at MRI: 10.1
12	[0.78] years, 53% male). Clinical neurologic symptoms were present in 40% of children in the
13	HIE-cohort. Non-parametric statistical analysis failed to detect any significant (p<0.001) alterations
14	of FC networks in the HIE-cohort, nor between children in the HIE-cohort with or without
15	neurological sequelae.
16	Interpretation: HT-treated HIE could not be associated with persistent alteration of the functional
17	connectome. This indicates a notable resilience of the functional architecture of the brain against
18	this type of early brain injury.
19	
20	
21	
22	
22	
23	
24	

## 1 Introduction

2 Neonatal hypoxic-ischemic encephalopathy (HIE) affects the infant brain during the time of basic 3 formation of the developing functional connectome. The functional connectivity (FC) networks that 4 comprise the connectome emerge as rudimentary networks already in the fetal brain (1). By full 5 term, primary sensory areas have formed more mature networks, while higher-order networks are 6 present in an immature state (1,2). During child development, these networks need to undergo 7 dramatic changes in order to fully integrate to a mature functional connectome (1). Resting-state 8 functional magnetic resonance imaging (rs fMRI) provides an opportunity to investigate networks 9 by utilizing the slow fluctuations in the blood oxygen level dependent (BOLD) signal to calculate 10 temporal correlations between brain regions (3).

11

12 HIE affects approximately 1-3 per 1000 live born children in countries with developed health care 13 and is a common cause of severe neurological disability (4). The acute event evolves into a neuro-14 detrimental process that can persist long after the initial insult (5). HIE can also negatively impact 15 neuroplasticity, which is essential for both normal brain development and recovery following brain 16 injury (6). The severity of encephalopathy is often assessed using a three-grade clinical score, such 17 as the Sarnat score (7). Treatment with therapeutic hypothermia (TH) reduces the risk for disability 18 and has been implemented as Standard of care for moderate to severe HIE in most high resource 19 settings (8). However, even with TH-treatment, most affected children exhibit some neurologic 20 abnormality when assessed at school age (8,9).

21

A few studies have identified aberrant FC networks in infants with neonatal HIE (10–14), and one
study have investigated effects at age 6-8 years (15). By early adolescence, FC network

24 development has typically reached a stable modular organization after which no significant

network reorganization is expected (1). The aim of our study was to examine the effects of TH treated neonatal HIE on the FC networks of the brain during early adolescence.

3

4 Method

### 5 **Participants**

6 All term or near-term ( $\geq$ 34 weeks gestational age) infants in Stockholm from January 2007 to 7 December 2009 who met clinical criteria for neonatal HIE and received TH-treatment were 8 included in the cohort. Children with genetic and/or metabolic disorders known to have 9 neurological effects, were excluded. This cohort and the clinical outcomes have been described 10 previously by Robertsson Grossmann et al (2022) (16). From the originally 66 included children, 11 one was excluded due to a genetic disorder and an additional 13 were ineligible for this follow-up 12 study due to being either deceased (n=8) or moved abroad (n=5) (see Figure S1 for flowchart). The 13 remaining 52 children were invited to participate. A control group of 43 children participated in the 14 same follow up protocol. The controls were singleton, term-born in Stockholm Sweden with an 15 uneventful neonatal period (5-min Apgar score > 3) that originally were recruited through random 16 identification from the Swedish Medical Birth Registry for a separate study (17). Our study was 17 approved by the Ethical Review Board in Stockholm, Sweden (2009/735-31/4, 2010/850-31/1, 18 2012/617-32, 2016/1921-32, 2019-01447, 2020-03318) and conformed to the Declaration of 19 Helsinki. Written, informed consent was obtained from all care givers and children gave assent to 20 participation.

21

#### 22 Data acquisition

23 Data collection was performed at Karolinska Institute and Karolinska University Hospital,

24 Stockholm, Sweden with neonatal and medical information collected prospectively. Follow up of

the HIE cohort was performed from January 2018 to December 2019 at age 10-12 years. Medical

1 assessment was performed by an experienced pediatric neurologist. Intelligence was assessed using 2 Wechsler Intelligence Scale for Children, fifth edition (18) (WISC-V, Swedish version), performed 3 by either an experienced child psychologist or a psychologist under supervision. Clinical diagnoses 4 were retrieved from the participants' electronic medical records. 5 MR images of the brain were acquired using a Sigma 3T MR Scanner (GE Healthcare) at the MR-6 Centrum, Karolinska Institute. No sedation was used. Participants were prepared for the scanning 7 with an optional information movie and familiarization using a mock scanner. At least one parent 8 was present with the child before scanning and observing from the monitor room. During 9 anatomical MRI acquisition all participants were offered to watch a movie or listen to music of 10 their choice. For the resting state fMRI session, the participants were instructed to remain still and 11 keeping their eyes open. T1-weighted two-dimensional spin-echo images (TE = 2,7 ms, flip angle 12 12 degrees, slice thickness 1.0 mm) were obtained in three planes with a 64-channel head coil. For 13 the resting state fMRI session, 300 T2\*-weighted whole-brain echo-planar image (EPI) volumes 14 were acquired over 10 minutes (TR = 2.0 s, TE = 30 ms, flip angle 70 degrees, 3x3x3 mm voxel 15 size).

16

### 17 Image Preprocessing and Denoising

Both functional and anatomical images were minimally preprocessed using fMRIPrep (19) 20.2.4.
Anatomical T1-weighted images were corrected for intensity non-uniform signal intensity with
N4BiasFieldCorrection (20). After skull-stripping with antsBrainExtraction, the T1-weighted
references were normalized to a pediatric template (7.5 to 13.5y age range [TemplateFlow ID:
MNIPediatricAsym:cohort-4] (21)) using volume-based spatial normalization through nonlinear
registration with antsRegistration (22) 2.3.3.
A visual quality control was performed and image quality metrics for resting-state fMRI data were

25 extracted using MRIQC (23) 0.16.1. For most subjects, motion artifacts were pronounced at the end

1 of the rs-fMRI sequence. We therefore decided to exclude the last minute of the data recorded, 2 resulting in a 9-minute BOLD sequence (270 volumes) for all subjects. Criteria for data exclusion 3 related to excessive subject head motion was set to frame-wise displacement (FD) > 0.5 mm on a 4 maximum of 20% of image volumes or DVARS > 1.5 SD. 5 6 In fMRIPrep, a functional reference volume and its skull-stripped version was generated using a 7 median of a motion corrected subset of volumes. The functional reference volume was co-8 registered to the T1-weighted image volume with FLIRT (24) (FSL 5.0.9) using a boundary-based 9 registration cost-function (25). The co-registration was performed using nine degrees of freedom to 10 account for remaining spatial distortions in the functional reference volume. Functional images 11 were resampled to native space using a fixed-body model (three translational and thee rotational 12 movement regressors) that was estimated from the reference volume using MCFLIRT (26) (FSL 13 5.0.9). After slice-time correction to 0.976s using 3dTshift from AFNI, the resampled functional 14 images were further resampled into a pediatric standard space (MNIPediatricAsym:cohort-4) (21). 15 16 Signal denoising of the BOLD signal time-series with respect to non-neuronal signal sources was 17 performed using principal component analysis-based noise regression. For this, we used CompCor 18 (27) to generate three probabilistic masks (cerebrospinal fluid (CSF), white matter (WM) and 19 combined CSF+WM) in native anatomical space. This was done using principal component analysis such that the retained components' time series are sufficient to explain 50% of variance 20 21 across the mask. Preprocessed BOLD time-series in MNI space were spatially smoothed with an 22 isotropic, Gaussian kernel of 6 mm FWHM (full-width half-maximum). The denoising regression 23 was performed in one single regression model to avoid reintroduction of artifacts (28) with the 24 following variables: six rigid body realignment parameters, top five anatomical CompCor

1 decompositions, motion outliers (frames exceeding 0.5 mm FD), FD and cosine filter (128s cut-of). 2 The data was low pass filtered at 0.1 Hz, no high pass filter was used aside from the cosine filter. 3 Group level spatial brain FC maps (with the HIE cohort and control group treated as a single 4 dataset) were calculated in Nilearn using canonical ICA (29) with 20 fixed components. ICA was 5 chosen rather than a static, parcellation-based delineation of the brain to increase methodological 6 flexibility for any cortical reorganization and/or recruitment of cortical areas outside the standard 7 location. Canonical ICA improves the ICA-analysis by adding a canonical correlation analysis to 8 the ICA-based pattern extraction for identification of a common data subspace and thresholding the 9 independent components based on the absolute value of voxel intensity (29). 10 11 Statistical analysis 12 Comparison of FC between children with HIE and controls was performed using dual regression 13 where the spatial maps derived from the canonical ICA were used to generate subject-specific 14 versions of both spatial maps and associated timeseries in a two-step regression analysis (30). 15 Testing for group-differences was performed using non-parametric testing with 5000 permutations 16 with correction for multiple comparisons using the threshold-free cluster enhanced (TFCE) 17 technique (31,32). 18 19 As noted in Table 1, age at MRI differed significantly between the groups. To address a possible 20 introduction of bias, we therefore performed an additional dual regression analysis in which we 21 included a covariate pertaining to group differences for age at scan in the dual regression model. 22 Similar to the previous model, significance of group differences was performed using non-

23 parametric testing (5000 permutations, multiple comparison correction using TFCE). Results were

24 thresholded for a minimal cluster size of 10 voxels and p-values of all significant results were

25 further corrected for multiple comparison of all FC components using Bonferroni correction.

- 44

2	A subgroup analysis was performed in which we divided the HIE-cohort into two groups based on
3	clinical outcome to examine if children with neurologic sequelae hade more alterations than normal
4	development, or if evidence for network reorganization affecting the functional outcome could be
5	detected in the group of HIE exposed children with normal development. Subjects fulfilling any of
6	the following criteria were classified as "unfavorable" outcome: IQ < 85, Cerebral Palsy (CP),
7	epilepsy, Central Visual Impairment (CVI), cerebral hearing impairment, Autism Spectrum
8	Disorder (ASD), Attention Deficit Hyperactivity Disorder (ADHD) and Developmental
9	Coordination Disorder (DCD). Remaining children with HIE were subsequently classified as
10	"favorable" outcome. Testing for subgroup-differences was done on the two subgroups with non-
11	parametric testing similar to the main group analysis.
12	

13 Results

### 14 Study sample

15 From the eligible HIE-cohort (n=52), four children declined all participation in the study for 16 personal reasons. An additional eight children specifically declined assessment with MRI, five of 17 these due to worry about being in the scanner, two due to severe CP and one due to not having 18 time. In the control group one participant did not complete the whole MRI protocol and two 19 participants were excluded due to the detection of cerebral abnormalities (one cerebellar arachnoid 20 cyst and one cortical cavernoma). An additional 15 participants were excluded based on set criteria 21 for excessive head motion during rs-fMRI, five from the HIE-cohort and ten from the control group 22 (see Figure S1 for an overview flowchart and Table S3 for sensitivity analysis). Characteristics of 23 the final study sample (n=65), consisting of 35 children in the HIE-cohort and 30 children in the 24 control group, is presented in Table 1. In the HIE-cohort, clinical neurologic diagnoses before the 25 follow-up were: 8 children (23%) with neurodevelopmental diagnose (i.e. ADHD, ASD and/or

- 1 DCD), 3 children (9%) with CP, 1 child (3%) with neurologic hearing impairment and 1 child (3%)
- 2 with CVI (Table S2). HIE grading was moderate (grade II) in 33 (94%) and severe (grade III) in 2
- 3 (6%) of the exposed children (Table S3).
- 4
- 5 **Table 1.** Characteristics of study population stratified by exposure to hypothermia-treated neonatal hypoxic-ischemic
- 6 encephalopathy.

Characteristic	HIE (n=35)	Control (n=30)	P value
Female, No (%)	19 (54.3)	14 (46.7)	
Male, No (%)	16 (45.7)	16 (53.3)	0.55
Gestational age at birth,			
median (IQR), wk	40.6 (39.2 – 41.2)	40.1 (39.4 – 41.1)	0.80
Birth weight, median (IQR), g	3500 (3326 - 3984)	3590 (3329 - 3738)	0.88
Age at WISC-V, median (IQR), y	11.3 (10.5 – 12.8)	12.1 (11.9 – 12.5)	< .001
Age at MRI, median (IQR), y	11.4 (10.5 – 11.8)	10.0 (9.6 – 10.6)	< .001
Framewise Displacement			
during MRI, mean (SD) mm	0.19 (0.13)	0.15 (0.08)	0.14
DVARS (standardized)			
during MRI, mean (SD) mm	1.14 (0.068)	1.16 (0.097)	0.32

Abbreviations: IQR, interquartile range; SD, Standard Deviation; WISC-V, Wechsler Intelligence Scale for Children
 version 5; MRI, Magnetic Resonance Imaging.

9

## 10 **Group analysis of functional connectivity**

11 20 fixed independent components were extracted from the combined bold dataset, of which five

12 were categorized as representing noise or artifacts. The remaining 15 FC networks were in general

- 13 agreement with known brain networks from previous literature (33) hierarchically representing nine
- 14 well-established networks: Occipital (visual) Network, Cerebellar Network,

Hippocampal/Brainstem Network, Pericentral (somatomotor) Network, Pericentral (auditory)
 Network, Lateral Frontoparietal (control) Network, Dorsal Frontoparietal (attention) Network,
 Medial Frontoparietal (Default Mode) Network, Midcingulo-Insular (salience) Network (Figure
 S2).

5

6 The non-parametric permutation tests revealed a total of 25 significant clusters residing in gray 7 matter (p<0.05, uncorrected for multiple statistical tests) in four of the 15 FC networks with a 8 majority residing in the cerebellar network (Table S1, Figure S3A). After subsequent additional 9 correction for multiple comparisons across individual networks (15 two-sided tests, corrected 10 significance threshold: p<0.00167), the medial visual network showed a decrease of FC in the right 11 dorsolateral prefrontal cortex (DLPFC) for the HIE cohort compared to the control group (Figure 12 S3B). Excluding the participant with CVI from analysis did not alter this result. However, 13 including age at scan as a co-variate of no-interest in our dual regression model resulted in this 14 finding no longer being significant. From the 25 significant clusters in the original analysis (before 15 correction for multiple comparisons), only 11 findings in two functional networks were significant 16 after inclusion of age as a covariate. Using AtlasReader (34), all clusters were anatomically labeled 17 by three different atlases (Automated Anatomical Labelling (AAL), Harvard-Oxford and Desikan 18 Kiliany). Combined with visual inspection, we deemed four of these clusters to be mainly localized 19 in cerebral white matter. The remaining 7 cluster differences consisted of decreased FC in the HIE 20 cohort in the left lateral Frontoparietal Network and the Cerebellar Network (Table 2, Figure 1). 21 None of these remained significant after correction for multiple comparisons.





Fig.1. Differences in functional connectivity between groups with and without exposure for hypothermia-treated neonatal hypoxicischemic encephalopathy corrected for age difference. Results from non-parametric testing (p<0.05, age corrected, voxelwise threshold free cluster enhancement corrected (TFCE), minimal cluster size = 10) on whole brain dual regression of the 15 ICA-derived networks (see Figure S2) detected clusters of decreased FC in children with hypothermia-treated HIE (N=35) compared to the control group (N=30) in the left Lateral Frontoparietal network (A) and the Cerebellar network (B). All significant changes (before correction for multiple comparisons) are colored red with seed networks are colored blue (transparent).

9

10 **Table 2.** Age corrected functional connectivity differences between groups with and without exposure for

11 hypothermia-treated neonatal hypoxic-ischemic encephalopathy.

		Hemisphere	Volume <sup>a</sup>		Peak	Peak	Peak
Network	Location		(mm <sup>3</sup> )	P value <sup>b</sup>	x <sup>c</sup>	y <sup>c</sup>	z <sup>c</sup>
Left lateral FPN	Superior Temporal	Right	2684	0.01	51.5	-36.5	11.5
Left lateral FPN	Superior Temporal	Right	8	0.0498	59.5	-22.5	11.5

CN	Frontal Pole	Right	3144	0.013	23.5	59.5	27.5
CN	Cerebellum	Left	872	0.0224	-30.5	-78.5	-48.5
CN	Superior Temporal	Right	712	0.0434	49.5	-26.5	5.5
CN	Hippocampus	Right	232	0.0432	27.5	-22.5	-10.5
CN	Middle Temporal	Right	16	0.0492	39.5	-54.5	19.5

1 <sup>a</sup> Thresholded for minimal cluster size of 10 voxels

2 <sup>b</sup>TFCE-corrected

3 <sup>c</sup> MNI coordinates

4 Abbreviations: FPN, Frontoparietal Network; CN, Cerebellar Network; VN, Visual Network; TFCE, Threshold Free

5 Cluster Enhancement; MNI, Montreal Neurological Institute and Hospital.

6

#### 7 Subgroup analysis

8 All 35 children in the HIE-cohort and a total of 22 children in the control group completed 9 cognitive assessment with WISC-V. Mean full scale IQ was 100 (SD 16,9) in the HIE-cohort and 10 112 (SD 12,5) in the control group. In the HIE-cohort, 14 (40%) of the children met the set criteria 11 for unfavorable outcome and the remaining 21 (60%) had favorable outcome (Table S2). The two 12 groups differed regarding gestational age at birth, with the Unfavorable outcome-group being born 13 significantly earlier (Table 3). Non-parametric testing for differences between the favorable and 14 unfavorable subgroups (TFCE-corrected) detected increased FC for the Favorable outcome-group 15 between the right lateral frontoparietal network and the cerebellum bilaterally and between the 16 dorsal frontoparietal network and the bilateral anterior cingulate and postcentral cortex (Table 4, 17 Figure 2). None of these findings did however survive additional correction for multiple 18 comparisons (threshold p<0.00167).

19

- 1 **Table 3.** Characteristics of the HIE-cohort stratified by "favorable" or "unfavorable" outcome as determined by
- 2 received clinical neurologic and/or neurodevelopmental diagnosis and/or IQ < 1,5 SD.

Characteristic	Favorable Outcome (n=21)	Unfavorable Outcome (n=14)	P value
Female, No (%)	13 (61.9)	6 (42.9)	
Male, No (%)	8 (38.1)	8 (57.1)	0.27
Gestational age at birth,			
median (IQR), wk	40.6 (40.3 - 41.3)	39.0 (38.1 - 40.1)	<.001
Birth weight, median (IQR), g	3706 (3380 - 3990)	3458 (3118 - 3838)	0.25
Age at WISC-V, median (IQR), y	11.2 (10.4 – 11.8)	11.2 (10.8 – 11.5)	0.95
Age at MRI, median (IQR), y	11.2 (10.4 – 11.8)	11.2 (10.8 – 11.7)	0.75
Framewise Displacement			
during MRI, mean (SD)	0.19 (0.14)	0.19 (0.12)	0.97
DVARS (standardized)			
during MRI, mean (SD)	1.13 (0.063)	1.15 (0.075)	0.34

3

4 **Table 4.** Differences in functional connectivity between children exposed to hypothermia-treated neonatal hypoxic-

## 5 ischemic encephalopathy with and without neurologic sequelae

Network	Location	Hemisphere	Volume <sup>a</sup>	P value <sup>b</sup>	Peak <sup>c</sup>	Peak <sup>c</sup>	Peak <sup>c</sup>
			(mm <sup>3</sup> )		х	у	Z

Decreased Connectivity in Group with Unfavorable Outcome

Right lateral	Cerebellum		176	0.017	-6.5	-40.5	-20.5
Right lateral	Cerebellum		112	0.031	5.5	-42.5	-22.5
Increased Connec	ctivity in group with Un	favorable Outcome					
Dorsal FPN	Anterior Cingulate	Left	616	0.027	-2.5	7.5	47.5
Dorsal FPN	Anterior Cingulate	Right	592	0.020	9.5	7.5	33.5
Dorsal FPN	Postcentral	Right	240	0.023	41.5	-14.5	35.5
Dorsal FPN	Postcentral	Left	112	0.036	-24.5	-14.5	49.5

6

7 <sup>a</sup>Thresholded for minimal cluster size of 10 voxels

8 <sup>b</sup>TFCE-corrected

- 1 <sup>c</sup> MNI coordinates
- 2 Abbreviations: FPN, Fronto-Parietal Network; TFCE, Threshold Free Cluster Enhancement; MNI, Montreal
- 3 Neurological Institute and Hospital.
- 4





Fig.2. Differences in functional connectivity between children exposed to hypothermia-treated neonatal hypoxicischemic encephalopathy with and without neurologic sequelae. Results from non-parametric test (p<0.05, TFCE,</li>
minimal cluster size = 10) of a dual regression analysis showing clusters of increased FC in the favorable subgroup
compared to the unfavorable subgroup between the right Lateral Frontoparietal Network to the cerebellum bilaterally
(A) and increased FC in the unfavorable subgroup compared to the favorable subgroup between the Dorsal
Frontoparietal Network to bilateral anterior cingulate and postcentral cortex (B). All significant changes are colored
red, seed networks are colored blue.

13

5

## 14 Discussion

In this study, we examined the resting-state FC networks in early adolescent children exposed to
HT-treated neonatal HIE. In our first model (i.e. without including scan at age as a co-variate in the

1	model), no alterations were found in a majority of the FC networks. But in four of the networks,
2	decreased FC in the HIE-cohort were observed and most prominently in the cerebellar network to
3	several areas located across the cortex. Early injury of the cerebellum has been linked to impaired
4	neurocognitive development (35). It has recently been recognized as underestimated and resulting
5	in volume reductions after neonatal HIE (36,37). However, after correction for multiple
6	comparisons across all resting-state networks, only one of these findings remained significant,
7	showing decreased FC in the HIE-cohort between the medial visual network and a cluster localized
8	to the right DLPFC which is a hub in the Lateral Frontoparietal (Control) Network but also partly
9	involved in the Dorsal Frontoparietal (Attention) Network (33).

10

11 Importantly, when including age as a co-variate of no-interest in the dual regression model, no 12 changes were found. This null result is in line with the only other study of long term FC effects 13 after neonatal HIE (Spencer et al, 2022, preprint) who compared FC in 22 children exposed to HT-14 treated neonatal HIE at age 6-8 with healthy controls (N=20) (15). Taken together, the absence of 15 significant major alterations in FC after HT-treated neonatal HIE suggests an overall high resilience 16 of the developing human connectome against this condition. Interestingly though, Spencer et al 17 reports case-control differences in FC between the Attention/Cognitive Control Network and the 18 Visual Network being significant before correcting for multiple comparisons. The common 19 minimal findings of altered FC between the Visual Network to Networks important for attention 20 and cognitive control indicates a possible common group effect that might be concealed due to 21 underpower of the two studies. The dorsal stream of the visual system integrates cognitive brain 22 functions like working memory and attention to the visual function (42,43). Specific vulnerability 23 of the dorsal stream has been described after conditions such as perinatal brain injury (44) and can 24 give rise to a cognitive subtype of cerebral visual impairment (CVI) with disturbed visual

perception and integration that is especially common after HIE (45). This is also highlighted by
 findings of a reduction in visuo-spatial processing ability as well as impaired structural connectivity
 in TH-treated HIE (46,47). A possible connection between this condition and exposure to neonatal
 HIE should be further investigated by combining fMRI with testing of visual function.

5

6 Earlier studies have reported aberrant FC in the early phase of neonatal HIE. Tusor et al (2014) 7 compared the FC (acquired within five weeks form birth) of four pre-selected FC networks in 8 infants (N=15) with neonatal HIE and healthy controls (N=15) (11). Jiang et al (2022) specifically 9 selected motor FC networks for comparison between infants with TH-treated neonatal HIE (N=16) 10 and neurologically intact controls (N=11) within 11 days from birth (10). Both studies reported 11 significant disruptions in multiple FC networks. Li et al compared atlas-based acquired network 12 properties of infants with mild (N=12) and severe (N=12) neonatal HIE and found reduced local 13 efficiency and clustering coefficient in the severe group within the first month of life. Finally, in 14 two retrospective studies Boerwinkle et al (2022, 2024) correlated FC from the acute phase with 15 clinical data at 6 months (12) and up to 42 months (mean 28.2 months) (13) in children with acute 16 neonatal brain injury including 24 children with neonatal HIE (of which 22 were exposed to 17 moderate to severe HIE according to Sarnat criteria). In comparison of typical versus atypical FC in 18 four selected networks they found several associations to assessment of general development, 19 cognition, motor function as well as mortality and epilepsy. The results from these earlier studies 20 should however be interpreted with caution mainly due to the small sample sizes employed but 21 taken together with our results we suggest that many of the FC changes seen in the acute or 22 subacute stage are either reversible or recovered in later development. After early focal injury, such 23 as stroke in the neonatal period, remarkable cortical reorganization of functions such as motor and 24 speech has been observed using fMRI (48). Our sub-group analysis compared the FC of

participants with or without neurologic sequelae but failed to identify any further significant
 changes or alterations suggestive of functional cortical reorganization in any of the networks.

4 The heterogenous impact on brain from neonatal HIE, as reflected by the broad spectrum of 5 neurological outcome of this condition, is another possible explanation for the absence of 6 significant findings in our study. The cortical injury pattern of neonatal HIE in the acute phase (if 7 not total) is commonly divided into either mainly Central/Basal Ganglia - Thalamus or Watershed 8 (border zone) injury, which has been shown to provide some prognostic information of neurologic 9 outcome (49). These patterns arise from specific brain regions (i.e., the basal ganglia, hippocampus 10 and primary motor cortex) being more susceptible to injury as a consequence of high metabolic 11 activity or containment of glutamate and/or from localization to the boundaries between blood 12 supply (e.g., border zone areas in middle-frontal and parietal-occipital cortex), depending on the 13 course of HIE (50). The patterns of injury are however commonly overlapping, whilst at the same 14 time many exposed children present a normal clinical MRI of the brain (51). Furthermore, the 15 secondary (and tertiary) phases of HIE involving inflammatory processes are not detectable on 16 MRI. It is therefore possible that individual injury patterns, along with insufficient effect size, led 17 to a dilution of the effects on a group level. In future studies, correlating the brain FC with 18 performance from clinical testing and increasing the study population could provide more 19 information on function specific network alterations in these children.

20

## 21 Limitations

There are several limitations to this study. First, although the current work is the largest study yet of FC in this rare condition, the sample size still limits its statistical power, especially for the

1	subgroup analysis. We also note a significant drop-out rate from the original cohort of 66 children.
2	The final analysis only included 40% of the original cohort of surviving children with HIE grade III
3	and 69% of the surviving children with HIE grade II (Table S3). Furthermore, some of the children
4	were unable to participate in the MRI acquisition only due to symptom severity (i.e. cerebral palsy)
5	and so the cohort of children included in the final analysis is not representing the full clinical
6	spectrum of HIE. Our results should therefore only be generalized to children with TH-treated
7	neonatal HIE without severe CP and mainly to neonatal HIE grade II.
8	Second, head motion during fMRI scanning is a substantial source for noise especially at younger
9	age (52). Exposure to neonatal HIE has been associated with unsuccessful MRI in early school age
10	(53) and diagnoses such as ADHD, autism spectrum disorder (ASD) and epilepsy also tend to be
11	associated with more movement (54). The control group in this study was scanned at a significantly
12	younger age (Table 1). The mean difference in age between groups was comparatively small (but
13	significant) and effected the results when included as a co-variate of no-interest in our dual
14	regression model. We have therefore chosen to only present the results with age included as
15	covariate in the main article, but due to the similarity in results with respect to earlier investigations
16	of this condition, we include no-age corrected results as supporting material.
17	
18	Acknowledgements
19	We thank the children and their parents for participating.
20	This study was financially supported by Stiftelsen Promobilia (F20501), Stiftlesen Sunnerdahls
21	Handikappfond (F18/20), Drottning Silvias Jubileumsfond, Lilla Barnets Fond, Jerringfonden,
22	Stiftelsen Kempe-Carlgrenska Fonden, HKH Kronprinsessan Lovisas Förening för Barnasjukvård,
23	Norrbacka-Eugeniastiftelsen (866/20), Neurofonden (F2020-0009) and grants provided by Region
24	Stockholm (ALF project and Centre for innovative medicine).

1	Abbreviations: ADHD, Attention Deficit Hyperactivity Disorder; ASD, Autism Spectrum Disorder;
2	BOLD, Blood Oxygen Level Dependent; CP, Cerebral Palsy; DCD. Developmental Coordination
3	Disorder; FD, Framewise Displacement; FC, Functional Connectivity; fMRI, functional Magnetic
4	Resonance Imaging; FWHM, full-width half-maximum; HIE, Hypoxic-Ischemic Encephalopathy;
5	IQ, Intelligence Quote; MRI, Magnetic Resonance Imaging; TFCE, Threshold Free Cluster
6	Enhancement; WISC, Wechsler Intelligence Scale for Children
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	

# 1 References

- Edde M, Leroux G, Altena E, Chanraud S. Functional brain connectivity changes across the
   human life span: From fetal development to old age. J Neurosci Res. 2021 Jan;99(1):236–62.
- Fransson P, Åden U, Blennow M, Lagercrantz H. The Functional Architecture of the Infant
   Brain as Revealed by Resting-State fMRI. Cereb Cortex. 2011 Jan;21(1):145–54.
- Fox MD, Raichle ME. Spontaneous fluctuations in brain activity observed with functional
   magnetic resonance imaging. Nat Rev Neurosci. 2007 Sep;8(9):700–11.
- Marlow N, Shankaran S, Rogers EE, Maitre NL, Smyser CD. Neurological and developmental outcomes following neonatal encephalopathy treated with therapeutic hypothermia. Semin
   Fetal Neonatal Med. 2021 Oct;26(5):101274.
- Davidson JO, Gonzalez F, Gressens P, Gunn AJ. Update on mechanisms of the pathophysiology
   of neonatal encephalopathy. Semin Fetal Neonatal Med. 2021 Oct;26(5):101267.
- Rocha-Ferreira E, Hristova M. Plasticity in the Neonatal Brain following Hypoxic-Ischaemic
   Injury. Neural Plast. 2016;2016:1–16.
- Sarnat HB, Sarnat MS. Neonatal Encephalopathy Following Fetal Distress: A Clinical and
   Electroencephalographic Study. Arch Neurol. 1976;33:696–705.
- Azzopardi D, Strohm B, Marlow N, Brocklehurst P, Deierl A, Eddama O, et al. Effects of
   Hypothermia for Perinatal Asphyxia on Childhood Outcomes. N Engl J Med. 2014 Jul
   10;371(2):140–9.
- Shankaran S, Pappas A, McDonald SA, Vohr BR, Hintz SR, Yolton K, et al. Childhood Outcomes
   after Hypothermia for Neonatal Encephalopathy. N Engl J Med. 2012 May 31;366(22):2085–
   92.
- 10. Jiang L, El-Metwally D, Sours Rhodes C, Zhuo J, Almardawi R, Medina AE, et al. Alterations in
   motor functional connectivity in Neonatal Hypoxic Ischemic Encephalopathy. Brain Inj. 2022
   Feb 3;1–8.
- 11. Smyser CD, Wheelock MD, Limbrick DD, Neil JJ. Neonatal brain injury and aberrant
   connectivity. NeuroImage. 2019 Jan;185:609–23.
- 12. Boerwinkle VL, Sussman BL, Manjón I, Mirea L, Suleman S, Wyckoff SN, et al. Association of
   network connectivity via resting state functional MRI with consciousness, mortality, and
   outcomes in neonatal acute brain injury. NeuroImage Clin. 2022;34:102962.
- Boerwinkle VL, Manjón I, Sussman BL, McGary A, Mirea L, Gillette K, et al. Resting-State
   Functional Magnetic Resonance Imaging Network Association With Mortality, Epilepsy,
   Cognition, and Motor Two-Year Outcomes in Suspected Severe Neonatal Acute Brain Injury.
- 34 Pediatr Neurol. 2024 Mar;152:41–55.

- 14. Li HX, Yu M, Zheng AB, Zhang QF, Hua GW, Tu WJ, et al. Resting-state network complexity and
   magnitude changes in neonates with severe hypoxic ischemic encephalopathy. Neural Regen
   Res. 2019;14(4):642.
- 4 15. Spencer APC, Goodfellow M, Chakkarapani E, Brooks JCW. Resting-state functional
   5 connectivity in children cooled for neonatal encephalopathy [Internet]. 2022 [cited 2024 Apr
- 6 22]. Available from: http://medrxiv.org/lookup/doi/10.1101/2022.12.17.22283576
- 16. Grossmann KR, Westblad ME, Blennow M, Lindström K. Outcome at early school age and
   adolescence after hypothermia-treated hypoxic-ischaemic encephalopathy: an observational,
   population-based study. Arch Dis Child Fetal Neonatal Ed.
- 10 17. Serenius F, Kallen K, Blennow M, Ewald U, Fellman V, Holmstrom G, et al.
- Neurodevelopmental Outcome in Extremely Preterm Infants at 2.5 Years After Active
   Perinatal Care in Sweden.
- 13 18. Wechsler D. Wechsler Intelligence Scale for Children Fifth Edition, Swedish Version. Pearson;
   2015.
- 15 19. Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, et al. FMRIPrep: a robust
   preprocessing pipeline for functional MRI. 2019;30.
- 17 20. Tustison NJ, Avants BB, Cook PA, Yuanjie Zheng, Egan A, Yushkevich PA, et al. N4ITK: Improved
   N3 Bias Correction. IEEE Trans Med Imaging. 2010 Jun;29(6):1310–20.
- 19 21. Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL. Unbiased average age appropriate atlases for pediatric studies. NeuroImage. 2011 Jan;54(1):313–27.
- 21 22. Avants B, Epstein C, Grossman M, Gee J. Symmetric diffeomorphic image registration with
   22 cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. Med
   23 Image Anal. 2008 Feb;12(1):26–41.
- 24 23. Esteban O, Birman D, Schaer M, Koyejo OO, Poldrack RA, Gorgolewski KJ. MRIQC: Advancing
   25 the automatic prediction of image quality in MRI from unseen sites. :21.
- 26 24. Jenkinson M, Smith S. A global optimisation method for robust affine registration of brain
   27 images. Med Image Anal. 2001 Jun;5(2):143–56.
- 28 25. Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based
   29 registration. NeuroImage. 2009 Oct;48(1):63–72.
- 26. Jenkinson M, Bannister P, Brady M, Smith S. Improved Optimization for the Robust and
   Accurate Linear Registration and Motion Correction of Brain Images. NeuroImage. 2002
   Oct;17(2):825-41.
- 27. Behzadi Y, Restom K, Liau J, Liu TT. A component based noise correction method (CompCor)
   for BOLD and perfusion based fMRI. NeuroImage. 2007 Aug;37(1):90–101.
- 28. Lindquist MA, Geuter S, Wager TD, Caffo BS. Modular preprocessing pipelines can reintroduce
   artifacts into fMRI data. Hum Brain Mapp. 2019 Jun;40(8):2358–76.

- Varoquaux G, Sadaghiani S, Pinel P, Kleinschmidt A, Poline JB, Thirion B. A group model for
   stable multi-subject ICA on fMRI datasets. NeuroImage. 2010 May;51(1):288–99.
- 30. Beckmann C, Mackay C, Filippini N, Smith S. Group comparison of resting-state FMRI data
   using multi-subject ICA and dual regression. NeuroImage. 2009 Jul;47:S148.
- 5 31. Smith S, Nichols T. Threshold-free cluster enhancement: Addressing problems of smoothing,
   6 threshold dependence and localisation in cluster inference. NeuroImage. 2009 Jan 1;44(1):83–
   7 98.
- 8 32. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the
   9 general linear model. NeuroImage. 2014 May;92:381–97.
- 33. Uddin LQ, Yeo BTT, Spreng RN. Towards a Universal Taxonomy of Macro-scale Functional
   Human Brain Networks. Brain Topogr. 2019 Nov;32(6):926–42.
- 34. Notter M P, Gale D, Herholz P, Markello R D, Notter-Bielser M-L, Whitaker K. AtlasReader: A
   Python package to generate coordinate tables, region labels, and informative figures from
   statistical MRI images. Ournal Open Source Softw. 2019 Feb 24;4(34):1257.
- 15 35. Stoodley CJ. The Cerebellum and Neurodevelopmental Disorders. The Cerebellum. 2016
   Feb;15(1):34–7.
- 36. Wu CQ, Cowan FM, Jary S, Thoresen M, Chakkarapani E, Spencer APC. Cerebellar growth,
  volume and diffusivity in children cooled for neonatal encephalopathy without cerebral palsy.
  Sci Rep. 2023 Sep 8;13(1):14869.
- 37. Annink KV, de Vries LS, Groenendaal F, van den Heuvel MP, van Haren NEM, Swaab H, et al.
   The long-term effect of perinatal asphyxia on hippocampal volumes. Pediatr Res. 2019
   Jan;85(1):43–9.
- 38. Nejati V, Movahed Alavi M, Nitsche MA. The Impact of Attention Deficit-hyperactivity Disorder
   Symptom Severity on the Effectiveness of Transcranial Direct Current Stimulation (tDCS) on
   Inhibitory Control. Neuroscience. 2021 Jul;466:248–57.
- 39. Biotteau M, Chaix Y, Blais M, Tallet J, Péran P, Albaret JM. Neural Signature of DCD: A Critical
  Review of MRI Neuroimaging Studies. Front Neurol [Internet]. 2016 Dec 16 [cited 2022 Aug
  8];7. Available from: http://journal.frontiersin.org/article/10.3389/fneur.2016.00227/full
- 40. Willcutt EG, Doyle AE, Nigg JT, Faraone SV, Pennington BF. Validity of the Executive Function
   Theory of Attention-Deficit/Hyperactivity Disorder: A Meta-Analytic Review. Biol Psychiatry.
   2005 Jun 1;57(11):1336–46.
- 41. Lachambre C, Proteau-Lemieux M, Lepage JF, Bussières EL, Lippé S. Attentional and executive
   functions in children and adolescents with developmental coordination disorder and the
   influence of comorbid disorders: A systematic review of the literature. Rodríguez C, editor.
   PLOS ONE. 2021 Jun 4;16(6):e0252043.

- 42. Ungerleider LG, Courtney SM, Haxby JV. A neural system for human visual working memory.
   Proc Natl Acad Sci. 1998 Feb 3;95(3):883–90.
- 43. Parks EL, Madden DJ. Brain Connectivity and Visual Attention. Brain Connect. 2013
   Aug;3(4):317–38.
- 44. Atkinson J. The Davida Teller Award Lecture, 2016: Visual Brain Development: A review of
  "Dorsal Stream Vulnerability"—motion, mathematics, amblyopia, actions, and attention. J Vis.
  2017 Mar 31;17(3):26.
- 45. Fazzi E, Signorini SG, Bova SM, La Piana R, Ondei P, Bertone C, et al. Spectrum of Visual
  Disorders in Children With Cerebral Visual Impairment. J Child Neurol. 2007 Mar;22(3):294–
  301.
- 46. Tonks J, Cloke G, Lee-Kelland R, Jary S, Thoresen M, Cowan FM, et al. Attention and visuospatial function in children without cerebral palsy who were cooled for neonatal
  encephalopathy: a case-control study. Brain Inj. 2019 Jun 7;33(7):894–8.
- 47. Spencer APC, Brooks JCW, Masuda N, Byrne H, Lee-Kelland R, Jary S, et al. Motor function and
  white matter connectivity in children cooled for neonatal encephalopathy. NeuroImage Clin.
  2021;32:102872.
- 17 48. Dunbar M, Kirton A. Perinatal Stroke. Semin Pediatr Neurol. 2019 Dec;32:100767.
- 49. Wisnowski JL, Wintermark P, Bonifacio SL, Smyser CD, Barkovich AJ, Edwards AD, et al.
  Neuroimaging in the term newborn with neonatal encephalopathy. Semin Fetal Neonatal
  Med. 2021 Oct;26(5):101304.
- 50. Miller SP, Ramaswamy V, Michelson D, Barkovich AJ, Holshouser B, Wycliffe N, et al. Patterns
   of brain injury in term neonatal encephalopathy. J Pediatr. 2005 Apr 1;146(4):453–60.
- 51. Weeke LC, Groenendaal F, Mudigonda K, Blennow M, Lequin MH, Meiners LC, et al. A Novel
   Magnetic Resonance Imaging Score Predicts Neurodevelopmental Outcome After Perinatal
   Asphyxia and Therapeutic Hypothermia. J Pediatr. 2018 Jan;192:33-40.e2.
- S2. Yan CG, Cheung B, Kelly C, Colcombe S, Craddock RC, Di Martino A, et al. A comprehensive
   assessment of regional variation in the impact of head micromovements on functional
   connectomics. NeuroImage. 2013 Aug;76:183–201.
- S3. Woodward K, Spencer APC, Jary S, Chakkarapani E. Factors associated with MRI success in
  children cooled for neonatal encephalopathy and controls. Pediatr Res. 2023 Mar;93(4):1017–
  23.
- 54. Yerys BE, Jankowski KF, Shook D, Rosenberger LR, Barnes KA, Berl MM, et al. The fMRI success
   rate of children and adolescents: Typical development, epilepsy, attention
   deficit/hyperactivity disorder, and autism spectrum disorders. Hum Brain Mapp. 2009
   Oct;30(10):3426–35.
- 36