

Facilitating the Virtual Exercise Games for Youth with T1D (ExerT1D) Peer Intervention: Protocol Development and Feasibility

Garrett I. Ash^{1,2}, Soohyun Nam³, Matthew Stults-Kolehmainen^{4,5}, Adrian D. Haughton^{1,6}, Carolyn Turek⁷, Annette Chmielewski¹, Michael Shelper¹, Julien S. Baker⁸, Stuart A. Weinzierl^{3,9}, Laura M. Nally⁹

¹Department of Internal Medicine, Yale School of Medicine, New Haven, CT, United States

²Center for Pain, Research, Informatics, Medical Comorbidities and Education Center (PRIME), VA Connecticut Healthcare System, West Haven, CT, United States

³Yale School of Nursing, Orange, CT, United States

⁴Center for Weight Management, Yale New Haven Hospital, North Haven, CT, United States

⁵Department of Biobehavioral Sciences, Teachers College – Columbia University, New York, NY, United States

⁶Department of Health and Movement Sciences Programs, Southern Connecticut State University, New Haven, CT, United States

⁷Child Study Center, Yale School of Medicine, New Haven, CT, United States

⁸Centre for Health and Exercise Science Research, Population Health and Medical Informatics, Hong Kong Baptist University, Kowloon Tong, Hong Kong SAR, China.

⁹Department of Pediatrics, Yale School of Medicine, New Haven, CT, United States

Correspondence:

Garrett I. Ash, PhD

Yale School of Medicine, Section of General Internal Medicine

Building 35A

950 Campbell Avenue

West Haven, CT 06516

Phone: 203-444-3079

Fax: 203-937-3829

Email: garrett.ash@yale.edu

ABSTRACT

Background

Barriers to moderate-to-vigorous physical activity (MVPA) for adolescents with type 1 diabetes (T1D) include physiology, transition to autonomy, and diabetes-specific stigma. Opportunities for T1D peer activities with T1D role model support are limited. To address this need, our single-arm pilot study tested the Home-based Virtual Activity Program for Youth with T1D (HAP-V-T1D) for feasibility.

Methods

Participants (n=15) were mean age 15.6 [SD 1.5] years, 7 non-Hispanic white, 6 female, 2 non-binary, mean A1c 8.9%±2.2%. The program included an MVPA videogame, physician-led education regarding managing T1D around MVPA, objective habitual MVPA goal-setting , and T1D management skills guided by young adult instructors living with T1D.

Results

For feasibility, 13/15 participants attended 10/12 sessions. Participants' perceptions of the program, comfort, instructors, and group cohesion were rated high/very high (4.2±0.5 to 4.8±0.3 out of 5). Motivation for the videogame was also high (4.1±0.4 out of 5). Instructor-adolescent interactions related to building T1D management skills were rated as excellent for 78% of sessions. Similarly, sharing knowledge and experiences were rated as excellent for 68% of sessions. However, adolescent-adolescent interactions were poor (communication 29% excellent, peer interactions 8% excellent). The most reported barriers to participation were negative mood and oversleeping. No participants experienced diabetic ketoacidosis, severe hypoglycemia, or injuries during the study period. Compared to baseline, glycemic metrics appeared to decrease during and post intervention (d= -0.72, -1.12).

Conclusion

HAP-V-T1D facilitated unprecedented T1D peer support achievements by engaging diverse youth with T1D in an MVPA program led by T1D role models. Larger studies are needed to assess if this intervention can improve glycemic measures and reduce diabetes-specific stigma.

Abstract **word** **count:** 264 (limit 300)

INTRODUCTION

The adolescent years present unique challenges to youth with type 1 diabetes (T1D). While it is estimated that only 20% of youth meet the recommended A1c level below 7%, it is well-known that intensive management of T1D is necessary for long-term health.^{1,2} Interventions aimed at improving health outcomes need to be introduced during the transition period of adolescence when youth are developing the skills they will need as adults to manage T1D. Navigating this transitional time presents challenges for adolescents with diabetes, creating a vital need for interventions to address T1D self-management support for adolescents.

Moderate-to-vigorous intensity physical activity is an essential complement to insulin therapy in T1D that has been demonstrated to improve overall physical (body composition, muscle strength, and cardiopulmonary fitness) and mental health, as well as diabetes-specific risks, including improving glycemic management, reducing insulin resistance, and potentially attenuating risk of long-term cardiovascular complications.³ Yet, MVPA is largely unaddressed by T1D support interventions for individuals with T1D at any age. Predicting an individual's glycemic response to MVPA is incredibly complex and requires more frequent glucose monitoring with subsequent adjustments to diet and insulin dosing. Further, managing MVPA can be especially challenging due to puberty-specific physiologic factors like increased insulin resistance.³⁻⁶ Moreover, for adolescents with T1D, glycemic variability encountered during MVPA may impede adopting healthy, active behaviors. Barriers to physical activity reported by adolescents include the psychological fear of glycemic destabilization⁷ and social stigma associated with disclosing an individual's diabetes diagnosis when engaging in MVPA. Apart from psychosocial factors, adolescents report that the logistical complications of glycemic dysregulation also impede their MVPA (e.g., needing to interrupt for carbohydrate supplementation).⁷ Adolescents also report teachers and coaches have limited T1D knowledge⁸ and their parents discourage MVPA.⁷ It is therefore no surprise that adolescents with T1D engage in less MVPA than the general adolescent population,⁹ of which less than 10% meets the MVPA recommendations of the Centers for Disease Control and Prevention (60min per day).¹⁰ Thus, it is warranted to pursue

53 multifaceted interventions targeted at addressing these barriers.

54 Current pediatric T1D MVPA guidelines detail the basic factors to consider during MVPA, but
 55 most providers do not actively prescribe them¹¹ and there is a striking lack of data regarding feasibility,
 56 acceptability, safety, and efficacy of protocols supporting their implementation for those with low
 57 baseline MVPA.³ Active videogames, defined as videogames that are controlled by gross motor body
 58 movements, have been leveraged as an alternative strategy to promote MVPA among youth,¹² aligned
 59 with new consensus guidelines that screen time should be tailored to achieve health benefits rather
 60 than minimized.¹³ Group activities incorporating active videogame technology with T1D mentors would
 61 provide an exciting new avenue to deliver all the components of a successful T1D self-management
 62 intervention:¹⁴ 1) behavior modification (goal setting, self-monitoring); 2) psychosocial support
 63 (interactions with T1D peers and role models); and 3) medical guidance (MVPA safety monitoring and
 64 advice). The purpose of the present study was to deliver a 6-week alpha version of a virtual home
 65 intervention to promote MVPA among adolescents with T1D, evaluate its feasibility, acceptability,
 66 safety, and describe its final curriculum for purposes of a future larger study. We hypothesized that
 67 metrics of feasibility and acceptability would meet our set criteria for warranting a future definitive trial.

68

69

METHODS

Overview

71 We present a single-arm prospective feasibility study of youth with T1D who underwent a 6-week
 72 intervention aimed at promoting MVPA. The study was approved by the Institutional Review Board
 73 (Yale-HIC#2000030105) in accordance with the Declaration of Helsinki and registered on
 74 clinicaltrials.gov (NCT05163912).

Participants and Enrollment

76 Participants were eligible if they were between the ages of 14 and 19 years (inclusive), had diagnosis
 77 of T1D for at least 6 months, were not achieving recommended physical activity targets (i.e.,
 78 <4days/wk with 60+min MVPA according to self-report at baseline), already wearing or willing to wear

a continuous glucose monitor (CGM) for the duration of the study, and did not have medical conditions that would preclude participation in group MVPA (cerebral palsy, current pregnancy, or others at investigator's discretion).

Participants were recruited from the Yale Children's Diabetes Center, clinicaltrials.gov, and diabetes social media groups (Facebook) between December 23, 2021 and June 15, 2022. In addition, recruitment materials were distributed publicly in collaboration with Children with Diabetes (T1-Today, Inc., Columbus, OH), a non-profit dedicated to providing education and support to families living with T1D.

First, potential participants completed a telephone screening to see if they met inclusion criteria for the study. The screening included questions about current MVPA by the Prochaska Physical Activity Assessment Question (specificity 61%-80%, test-retest ICC=0.6-0.8),¹⁵ as well as the Physical Activity Readiness Questionnaire.¹⁶ For any health conditions presenting special considerations for physical activity, additional documentation was requested from the primary care physician or sub-specialist for MVPA clearance. Each qualifying participant and one parent completed an electronic informed permission/assent and consent process prior to any study-related activities. Following consent/assent, two additional descriptive surveys were completed by the participant (recreational screen time,¹⁷ pubertal development status¹⁸) and a demographics form was completed by the parent. The research team shipped supplies to the participant and held a 1.5-hour HIPAA compliant Zoom (Zoom Video Communications, San Jose, CA) tele-video call to guide setup.

Equipment setup

Participants fitted their home television monitor with a Nintendo Switch Entertainment System (Nintendo of America, Inc., Redmond, WA) including Ring Fit Adventure cartridge and equipment. Ring Fit Adventure alternates low- (stationary walking or jogging) with higher-intensity exercises that are dynamic (e.g., knee-lifts) or resistance-based (e.g., overhead press, squat, abdominal press). The game uses a leg-strap accelerometer and handheld ring with gyroscopic/accelerometer sensors to measure distance covered and successful repetitions of each exercise. Success with exercises leads

to unlocking of more advanced adaptations (e.g., abdominal press upgraded to Russian twist). It has been previously demonstrated to stimulate activity intensity reaching the MVPA threshold among children.¹⁹ Participants were also provided with a metal tripod and Bluetooth headphones, so they could position their smartphone to capture the game screen while communicating over Zoom during the group MVPA. They were also provided a yoga mat, poster with Borg 6-20 rating of perceived exertion scale, Precision Xtra Blood Ketone meter and strips (Abbott Diabetes Care, Alameda, CA), and Fitbit Inspire 2 (San Francisco, CA) to track goals for weekly steps and MVPA heart rate. Fitbit accuracy studies have yielded variable results but a systematic review with quantitative synthesis reported the majority found overestimation of steps (~5%) and underestimation of heart rate (~3%).²⁰

Intervention

The intervention was based upon Social Cognitive Theory which predicts that an individual health behavior such as MVPA is influenced by individual experiences, environmental factors, and the actions of others.²¹ We therefore integrated peer and role model interactions into home-based MVPA: livestreamed active videogame sessions and tele-video discussions about experiences self-monitoring MVPA and glucose.

We involved three role models ages 20-39yr living with T1D and participating in nationally visible athletics, amateur athletics, and a non-performance-based physically active lifestyle respectively. Additionally, two of them were nationally visible advocates for T1D and the third was a certified diabetes care and education specialist. They agreed to work in a rotation to attend classes. The sessions were also monitored by a board-certified pediatric endocrinologist living with T1D and an exercise physiologist. Each of these two monitors attended all but two classes respectively, for which they designed the lesson plan and remained available by phone in the event of clinical concerns. Planning discussions and two pilot sessions were held with input from the above leaders, a pediatric exercise physiologist, an exercise telemedicine clinician, a pediatric psychologist seeing patients with T1D, and senior diabetes investigators in behavioral science and technology respectively.

Participants were recruited in 4 sequential cohorts for a 6-week intervention with meetings every


131 Saturday and Wednesday (Table 1) over HIPAA compliant Zoom tele-video. Sessions began with an
 132 icebreaker question that was either T1D-related (e.g., “what is your favorite low blood sugar
 133 treatment?”) or not (e.g., “if you could bring one thing with you on a deserted island, what would it
 134 be?”). At the first session in each cohort, the pediatric endocrinologist gave an overview of safe
 135 exercise practices related to T1D management based on national and international guidelines current
 136 at the time of the study.^{4,5,22} At each session, before and after exercise teens reported verbally or
 137 through the Zoom chat function to the clinician their sensor glucose value and trend arrow, insulin
 138 dosing (last bolus and/or insulin-on-board), and results of a ketone test if sensor glucose exceeded
 139 250mg/dL (13.9 mmol/L). The clinician then provided recommendations for exercise based on
 140 published guidelines.^{4,5,22} Clinical guidance was given when ketones were present (≥ 0.6 mmol/L). For
 141 safety, the instructors also confirmed that the participants had fast-acting carbohydrates accessible
 142 (e.g., orange juice, glucose tablets), proper exercise shoes and clothing, and an 8'-by-8' unobstructed
 143 space for exercise. On a weekly basis, the endocrinologist reviewed data from CGM, ketone tests, and
 144 clinical events, identified any safety concerns related to exercise, and communicated these concerns to
 145 the participants and their parents.

146 The segment devoted to exercise was 32min the first Saturday, 40min subsequent Saturdays,
 147 and 20min on Wednesdays. Participants were required to remain “present” for the full session, with
 148 either their image or the television display of their avatar reflecting their exercise in Zoom camera view
 149 (Figure 1). They communicated to the instructor if they needed to leave the session for any reason. For
 150 the purposes of coaching exercise, we tracked the proportion of each session spent in active
 151 movement, repetitions of each exercise, Borg 6-20 rating of perceived exertion of each exercise, and
 152 average Fitbit heart rate.

153 The remaining 20-30min of each session was spent on discussion activities over Zoom. Week 1
 154 sessions were dedicated to creating exercise goals. Instructors and participants discussed and agreed
 155 upon individual fitness-related goals. Goal-setting followed the SMART principle (Specific, Measurable,
 156 Attainable, Relevant, Timely). In addition, participants in each cohort formulated a group goal of

157 cumulative achievements (e.g., group steps) to promote a cooperative rather than competitive
 158 approach.²³ Goal progress was checked and discussed each week, including the provision of MVPA
 159 and T1D management information that helped support the specific goals. At the week 6 sessions, the
 160 instructors led a discussion about the relevance of the intervention's MVPA to building and sustaining
 161 an overall active lifestyle.

162 At the week 2-5 sessions, instructors engaged teens to create role-playing skits that involved
 163 educational points related to T1D management. Skit planning and acting occurred through verbal
 164 dialogue on Zoom. It included choosing a plot based on suggestions from instructors and the teens,
 165 developing a list of roles, choosing and assigning roles, developing a script with a combination of set
 166 and improvised lines, and acting. The instructors and endocrinologist guided the planning so that the
 167 skit integrated educational points about T1D.



168
 169
 170
 171
 172 **Figure 1.** Virtual exercise session livestreamed over Zoom.

173

Table 1. Structure of intervention sessions.

Component	Saturdays Time of Day ¹	Wednesdays Time of Day
Icebreaker question	10:00-10:05	19:00-19:05
Sensor glucose check and guidance ²	10:05-10:20	19:05-19:20 ³
Exercise	10:20-11:00	19:20-19:40 ³
Sensor glucose check and guidance	11:00-11:05	19:40-19:45 ³
Discussion Activities	11:05-11:30	19:45-20:00
• MVPA goals (week 1: set goals; weeks 2-5: check progress and provide MVPA and T1D management information supporting goals; week 6: reflection on relationship to overall active lifestyle) • Role-playing skits (weeks 2-5: planning and acting)		

¹USA eastern time zone, which was the local time of all participants except 1 who was USA mountain time zone (2 hours earlier)

²Extended to 30 minutes at first session to give general overview of guidelines. This session had reduced exercise by 8 minutes and discussion activities by 5-10 minutes.

³Cohort #1 only had these components on Saturdays (i.e., Wednesday session only met 19:00-19:20). Added these components to Wednesdays per expressed interest of cohort #2.

174

175

176 Assessments

177 Fidelity

178 Session recordings were double-coded by two exercise physiologists (one not involved with
 179 intervention). They identified missed opportunities for practicing each core competency of a role model
 180 and peer intervention:²⁴⁻²⁸ building skills, sharing knowledge and experiences, enriching
 181 communication, and facilitating peer interactions. Each competency was then coded as “excellent”
 182 (competency evident with 0 missed opportunities), “fair” (competency evident with ≥1 missed
 183 opportunities), or “poor” (competency not evident). The identified missed opportunities were discussed
 184 with instructors at weekly meetings as areas to iteratively improve the intervention. To generate rich
 185 discussion, each missed opportunity identified by either rater was included provided agreement by the
 186 other rater.

187 Follow-Up Retrospective Survey of Acceptability

188 The Motivation for Exergame Play Inventory²⁹ in our sample had adequate reliability for the full scale
189 ($\alpha=0.79$, 22 items) and the sensory immersion/flow subscale ($\alpha=0.79$, 8 items). The other subscales are
190 typically considered too short to establish meaningful reliability (3-4 items) but nonetheless reported.²⁹

191 The perceived cohesion scale had adequate reliability for the morale (e.g., “I am happy to be
192 part of this group”) ($\alpha=0.78$) and belonging subscales (e.g., “I see myself as part of this group”)
193 ($\alpha=0.83$).³⁰

194 We also issued a satisfaction survey taken from our past work²⁵ for participants to evaluate
195 program components/strategies (5 items, $\alpha=0.72$), personal comfort level (6 items, $\alpha=0.81$), and
196 interactions with instructors (12 items, $\alpha=0.95$). Participants also rated their top-3 of these factors by
197 order of importance.

198 Finally, participants and leaders each completed a semi-structured satisfaction interview which
199 will be reported separately. Leaders consented verbally after reviewing an information sheet.

200 **Ancillary Analysis of Glycemic Outcomes**

201 HbA1c was unlikely to change during the short timeframe of the intervention. To explore
202 probable magnitude of glycemic effects to inform future trials, we calculated the glucose management
203 indicator (GMI) for each 14-day block of the 6-week intervention. GMI uses CGM to project HbA1c
204 using an equation from cohort data.³¹ We also estimated maintenance of glycemic effects using follow-
205 up HbA1c values 39 (SD=15) weeks later, collected at screening for an addendum study (September 1,
206 2022 through September 19, 2023).

207 **Data Analysis**

208 Criteria for warranting a future definitive trial were recruitment uptake ($\geq 35\%$),³² attendance ($\geq 75\%$
209 attending $\geq 75\%$ of sessions),³³ biosensor wear-time ($\geq 70\%$),³⁴ survey completion ($\geq 85\%$), fidelity ratings
210 ($\geq 70\%$ excellent, $\geq 90\%$ at least fair), and psychosocial perceptions of acceptability (≥ 3.0 out of 5).

211 Metrics were tested for normality using the Shapiro-Wilks test, then reported by mean and standard
212 deviation, median and interquartile range, or frequencies as appropriate. We refined the curriculum
213 based on data after each cohort, ending the trial when no further refinements were found. Probable

214 magnitude of glycemic effect from baseline was calculated non-parametrically due to the small sample
215 size, by calculating the Wilcoxon Signed Rank test statistic (SPSS v28, Chicago, IL) then using Fritz's
216 formulas³⁵ to calculate biserial r and convert to Cohen's d (≥ 0.20 is small, ≥ 0.50 moderate, ≥ 0.80 large).

217 **RESULTS**

218 **Recruitment**

219 Clinic recruitment yielded 26 candidates referred from providers. Upon phone call by the study team, 5
220 were disqualified due to exceeding the baseline physical activity criteria, 6 declined to participate (2
221 due to lack of energy for physical activity, 2 due to time conflicts, 2 without giving a reason). The
222 remaining 15 scheduled a consenting visit, of whom 8 withdrew prior to starting the intervention (4
223 without giving a reason, 1 due to technical difficulties sharing CGM, 1 due to time conflicts, 1 due to
224 illness, and 1 due to change in housing conditions). The remaining 7 enrolled. One of them required
225 clearance for a medical condition from a specialist (arthritis).

226 Children with Diabetes posted the advertisement in three issues of their weekly newsletter
227 (February 9th–February 23rd, 2022) as well as their Facebook page on February 15th. The Facebook
228 post received 12 likes, 10 shares, and 23 email inquiries. After the study team replied with a full
229 description of the study, 17 completed eligibility screening. Among them, 9 were disqualified due to
230 exceeding the baseline MVPA criteria, and the other 8 qualified, consented, and started the
231 intervention.

232 Thus overall, 23/43 of those screened were eligible and scheduled a consent visit, among
233 whom 15 enrolled (35%). All 15 completed follow-up assessments.

234 **Demographics**

235 Eight states were represented: 8 participants from CT, and 1 each from DE, FL, GA, MD, NJ, UT, VA.
236 Participants represented diverse demographics: 8 people of color, 6 female, 2 non-binary, 4 public
237 insurance, 2 household income below poverty line, and 7 household income $< \$60,000$. Notably, most
238 of the participants had A1c levels above American Diabetes Association targets: 12 with A1c $> 7.0\%$
239 (> 53 mmol/mol), 5 with A1c $> 10.0\%$ (> 86 mmol/mol). Diverse participants appeared to stem from clinic

240 recruitment rather than social media (Table 2).

241

242

Table 2. Demographics (n=15, mean±SD).

	Recruited from Clinic	Recruited from Social Media	Total
N	7	8	15
Age (yr) ¹	15.8±1.1	15.5±1.9	15.6±1.5
Gender	3 female (43%) 2 non-binary (29%) 2 male (29%)	3 female (38%) 0 non-binary (0%) 5 male (62%)	6 female (40%) 2 non-binary (13%) 7 male (47%)
Pubertal Stage	0 Early (0%) 0 Mid (0%) 5 Late (71%) 2 Post (29%)	1 Early (13%) 3 Mid (38%) 3 Late (38%) 1 Post (13%)	1 Early (7%) 3 Mid (20%) 8 Late (53%) 3 Post (20%)
Race/Ethnicity	1 non-Hispanic black (14%) 4 Hispanic white or other (57%) 0 more than one race 2 non-Hispanic white (29%)	2 non-Hispanic black (25%) 0 Hispanic white or other (0%) 1 more than one race (13%) 5 non-Hispanic white (63%)	3 non-Hispanic black (20%) 4 Hispanic white (27%) 1 more than one race (7%) 7 non-Hispanic white (47%)
A1c (%)	10.1±2.1 7 over 7.0% (100%) 4 over 10.0% (57%)	7.9±1.8 5 over 7.0% (63%) 1 over 10.0% (13%)	8.9±2.2 12 over 7.0% (67%) 5 over 10.0% (33%)
A1c (mmol/mol)	87±34 7 over 53 (100%) 4 over 86 (57%)	63±20 5 over 53 (63%) 1 over 86 (13%)	74±24 12 over 53 (67%) 5 over 86 (33%)
Duration of type 1 diabetes (yr)	8.0±4.1	8.4±3.8	8.2±3.8
Household income annual	2 <\$20,000 (29%) 3 \$40,000-\$59,999 (43%) 1 >\$100,000 (14%) 1 declined to respond	0 <\$20,000 (0%) 2 \$40,000-\$59,999 (25%) 3 \$80,000-\$99,999 (38%) 3 >\$100,000 (38%) 0 declined to respond	2 <\$20,000 (13%) 5 \$40,000-\$59,999 (33%) 3 \$80,000-\$99,999 (20%) 4 >\$100,000 (27%) 1 declined to respond (7%)
Insurance (N public, %)	3 (43%)	1 (13%)	4 (27%)
Therapy (N pump, %)	6 (86%)	8 (100%)	14 (94%)
Total daily insulin dose (U/kg)	0.8±0.2	0.9±0.3	0.9±0.3
Body mass index (%ile) ¹	79.0±12.4	55.8±35.7	66.6±29.1
Physical activity (days/wk with 60+ min MVPA)	1.1±1.1	1.8±1.0	1.5±1.1
Screen time (hours/day)	9.3±5.2	8.3±4.4	8.8±4.6

¹Non-normally distributed (age right-skewed, body mass index left-skewed).
Bolded figures indicate apparent difference from social media group.

243

244 Intervention Feasibility

245 Attendance

246 Overall, 13 participants attended 10/12 sessions. Ten participants attended 11 sessions. Three
 247 participants attended 10 sessions. Among absences by these participants, 7 were due to family
 248 activities, 1 was due to unavailability of television set or computer screen, 2 were due to school
 249 extracurricular activities, and 2 were unexplained. The remaining two participants attended 6 sessions.
 250 One of these participants reported the main cause of non-attendance was negative mood, and the
 251 other reported it was oversleeping. Finally, all 15 participants completed the follow-up survey and
 252 interview, the latter lasting 18-48min (median 20 [IQR 19,31]). All 5 staff also completed the interview
 253 (31-48min).

254 The last cohort took one week longer than expected to recruit the minimum 3 participants. In
 255 fairness to the 2 participants who signed up on the preset schedule, we offered them 2 sessions during
 256 the week of delay which were credited as extra sessions.

257 Technical Barriers

258 One participant had technical problems that they rated as causing a moderate or higher disruption,
 259 which included 1 day of dead controllers that took 85% of the MVPA session to recharge, and 2 days
 260 when they conversed by text chat due to phone audio interruption. In addition, 2 participants had Zoom
 261 connectivity interruptions on 1-2 occasions, which they reported caused mild disruption.

262 Fidelity

263 Instructor competencies were rated highest for building T1D and MVPA management skills followed by
 264 sharing T1D and MVPA knowledge and experiences, enriching communication at the group sessions,
 265 and facilitating peer interactions at the group sessions (Table 3). Most competencies trended upward
 266 for each successive cohort, except #3 to #4 where they trended downward. Cohort #4 was tied for
 267 lowest size and had lowest attendance among those enrolled. Cohorts had different instructors, but all
 268 the instructors achieved similar scores on each competency.

269 Intervention Acceptability

270 Participant satisfaction ratings were highest for interactions with instructors, followed by perceived

271 group cohesion, program components and strategies, comfort (Table 3), and lastly Ring Fit game
 272 motivation (Table 4). The instructor component was also most frequently selected as a top-1 or top-3
 273 important component (80%, 60%). Nonetheless, a few selections related to comfort (13%, 23%) or
 274 program structure (7%, 16%).

Table 3. Attendance, Fidelity, Satisfaction, and Perceived Cohesion by Group

Cohort	Size (n)	Attendance (Mean %)	Instructor Competencies (% of sessions rated excellent) ¹				Psychosocial Perceptions mean±SD (% rated >=4 out of 5)			
			Building T1D and MVPA Management Skills	Sharing T1D and MVPA Knowledge and Experiences	Enriching Communication at the Group Sessions	Facilitating Peer Interactions at the Group Sessions	Interactions with Instructors	Group Cohesion (Perceived)	Program Components and Strategies	Comfort
1	3	75	75	58	25	0	4.6±0.7 (67%)	4.2±0.9 (67%)	4.1±0.2 (100%)	3.9±0.9 (67%)
2	4	92	100	90	46	27	4.9±0.0 (100%)	4.6±0.4 (100%)	4.5±0.3 (100%)	4.5±0.1 (100%)
3	5	95	100	100	50	0	4.9±0.2 (100%)	4.2±0.4 (80%)	4.3±0.6 (60%)	4.2±0.4 (80%)
4	3	72	38	25	0	0	4.8±0.3 (67%)	4.7±0.2 (100%)	4.3±0.7 (33%)	4.3±0.6 (33%)
Total	15	76	78	68	29	8	4.8±0.3 (93%)	4.4±0.5 (87%)	4.3±0.5 (73%)	4.2±0.5 (80%)

¹All other sessions were scored "fair" (i.e., none were scored "poor").

T1D, type 1 diabetes.

MVPA, moderate-to-vigorous physical activity.

Table 4. Exergaming Motivation (n=15)

Component	Mean±SD (% rated ≥4 out of 5)
Goals of the Game	4.6±0.4 (93%)
Performance Feedback	4.3±0.5 (87%)
User Control	4.3±0.6 (73%)
Sensor Immersion/Flow	4.0±0.6 (53%)
Game Challenge and Difficulty	3.2±0.6 (13%)
Total	4.1±0.4 (60%)

279 **Intervention Safety and Monitoring**

280 No participants experienced diabetic ketoacidosis, severe hypoglycemia, or other study-related adverse
 281 events during the study period. One participant developed elevated ketone levels prior to exercise (1.1
 282 mmol/L), that resolved after insulin administration prior to exercise. One participant deferred exercise
 283 due to feeling unwell as a result of having a high sensor glucose level (400mg/dL, 22.2 mmol/L) without
 284 elevated ketones. No participants reported a glucose level less than 70mg/dL (3.9 mmol/L) during the
 285 sessions, although there were 3 participants who reported hypoglycemia symptoms on 1 occasion that
 286 resolved upon ingesting 10g-15g of carbohydrates. As well, CGM review indicated 3 participants had
 287 sensor glucose levels less than 70mg/dL (3.9 mmol/L) less than 1 hour after 1 of the sessions. These
 288 instances resolved without guidance from staff.

289 All participants were using a CGM in their clinical care before the study (12 Dexcom G6, 3
 290 Abbott Libre 2). Weekly CGM audits found 75.3% of the possible readings were captured across all
 291 study participants. Ten participants met the recommended $\geq 70\%$ completeness. Primary reasons for
 292 not wearing CGM at least 70% of the time included scanning the Libre 2 sensor less frequently than its
 293 8hr memory duration (n=2), cloud syncing errors (n=1), changing the Dexcom G6 sensor less
 294 frequently than its 10-day life (n=1), and periodic psychological discomfort with being monitored (n=1).
 295 Study coordinators checked CGM wear-time twice weekly at the group sessions and addressed
 296 barriers with participants outside of each session.

297 Weekly CGM physician reviews resulted in 4 participants being referred to their providers for
 298 insulin dose adjustments to address elevated percentages of low sensor glucose values. In 3 cases the
 299 issue was resolved following adjustment by the diabetes provider, and in 1 case it was resolved
 300 following a small adjustment by the parent (change in insulin correction factor by 20%) and checking in
 301 with the teen about accurate carbohydrate counting. The physician also noted 3 instances among 3
 302 participants where sensor glucose had overnight lows after evening exercise. Trends were addressed
 303 with the participant directly. Topics related to avoiding hypoglycemia after exercise were also discussed
 304 in the group lesson.

305 Fitbit heart rates were captured for 96/117 person-sessions (82%), with 14 having at least one
306 session. Among them, 1 averaged in the vigorous range of 70%-90% age-predicted maximum, 12
307 averaged in the moderate range of 55%-70%, and 1 averaged below the moderate range. The average
308 participant heart rate across sessions was 61.4% (SD=5.2%) of maximum.

309 **Intervention Activities**

310 Participants set individual goals related to achieving Fitbit step counts, playing a sport, playing Ring Fit
311 outside of group sessions, or managing diabetes (Table 5). The success rate was 33% overall and
312 similar across categories. One participant declined to wear the Fitbit due to wrist joint discomfort, so
313 instead quantified a step goal as walking from school to home. Group goals related to sleep duration (1
314 group), playing Ring Fit outside of class (2 groups), and average Fitbit distance (1 group). The last two
315 of the four were successful.

Table 5. Individual Goals (n=15)

Goal Category	Number of participants	Specific Value	Achieved Goal
Steps	6	School to home, 3d/week 4000 per day 7000 per day 8500 per day 10000 per day 12500 per day	2 / 6 (33%)
Sport	5	Find a soccer group Bike 26.2 miles Run 8:00 mile Run 6:35 mile Run 7:30 pace for 5 miles	2 / 5 (40%)
Ring Fit	3	20 minutes per day 500 reps of 1 specific movement Reach world 9	1 / 3 (33%)
Diabetes management	1	7.0% estimated A1c from CGM	0 / 1 (0%)
Total	15		5 / 15 (33%)

316 Fitbit wear time on average met literature standards (70.5%, or 16.9 hours per day suggesting
317 an average of 10-15 hours during waketime)³⁴. Cohort 1 participants stopped wearing the Fitbit with 2-4
318 weeks left in the program, so we instilled a biweekly check of wear-time and troubleshooting after which
319 this pattern did not repeat. Average steps for those with Fitbit step goals were 8274 (SD=3734) per day
320 and for the total sample were 7178 (SD=4013) per day.

Each skit successfully integrated 3-5 educational points (Table 6).

Table 6. Structure of topics, roles, and competencies taught during skits for each cohort.

Cohort	Topic	Roles ¹	Educational Points and How Integrated
1	Team sports and type 1 diabetes (T1D)	-Athlete with newly diagnosed with T1D -Teammate living with T1D for several years -Continuous glucose monitor with alerts -Coach	-Speaking to coaches who are not familiar with T1D management around MVPA -Peer mentoring -Competitive sports and glycemic management
2 & 3	Zombie Apocalypse	-Zombie -Zombie-killing engineer -Teen newly diagnosed with T1D and knows nothing about what they need for T1D management during an apocalypse -Teen with longer duration of T1D -Dr. Rescue -News host -Continuous glucose monitor with alerts	-Essential supplies for T1D -Prioritizing aspects of T1D management -Operating continuous glucose monitors -Impact of MVPA on glycemic management -Impact of stress on glycemic management
4	Wilderness Excursion	-Lilo (Hawaiian girl) -Stitch (helpful alien) -Evil singing crab -Battler of evil singing crab -Diabuddy (diabetes buddy) -Angel -Narrator	-Same as Zombie Apocalypse

¹Self-chosen by interested participants, remaining ones assigned.
MVPA, moderate-to-vigorous physical activity

Ancillary Analysis of Glycemic Outcomes

Compared to baseline HbA1c, GMI during the intervention was moderately lower and HbA1c at maintenance follow-up was largely lower (Table 7). From baseline to maintenance follow-up, change in body mass index percentile was negligible (67 ± 29 to 66 ± 30 %ile, $d = -0.10$, 95% CI $[-0.37, 0.17]$) and in total daily insulin dose was small (0.88 ± 0.27 to 0.83 ± 0.24 U/kg, $d = -0.35$, 95% CI $[-0.62, -0.07]$). Seven participants upgraded their insulin device to a closed-loop system, but device upgrades did not drive HbA1c changes; i.e., the average HbA1c change among these 7 participants was the same as the full cohort.

Table 7. Glycemic outcomes (n=15)

	Mean $\pm$ SD (%)	Mean $\pm$ SD (mmol/mol)	# of participants having decrease from baseline	# of participants having increase from baseline	# of participants missing*	Effect size (d) vs baseline [95% CI]
Baseline HbA1c †	8.9 $\pm$ 2.2	74 $\pm$ 24	N/A	N/A	0 missing HbA1c (6 missing CGM)	N/A
Weeks 1-2 GMI	8.7 $\pm$ 2.0	72 $\pm$ 31	6	4	5 missing CGM	-0.35 [-0.62, -0.07]
Weeks 3-4 GMI	8.4 $\pm$ 1.5	68 $\pm$ 17	6	6	3 missing CGM	-0.45 [-0.73, -0.17]
Weeks 5-6 GMI	8.3 $\pm$ 1.6	67 $\pm$ 18	7	5	3 missing CGM	-0.72 [-1.01, -0.43]
Long-term HbA1c ‡	8.3 $\pm$ 1.8	67 $\pm$ 20	10	3	2 missing HbA1c	-1.12 [-0.81, -1.44]

GMI, glucose management indicator which projects HbA1c based on continuous glucose monitor (CGM) data.

*Missing values reflect partially complete CGM data (<70%) that was excluded from analysis and handled by intention-to-treat (last observation carried forward) for calculation of mean values and effect size. Every participant had at least one non-imputed value following baseline.

†Taken an average of 7 weeks (range 1-14) before intervention start

‡Taken an average of 45 weeks (range 19-77) after intervention start

DISCUSSION

Adolescents with T1D are vulnerable to inactivity, challenges with diabetes self-management, and social isolation. While many interventions have targeted this problem, the incorporation of peers and role models is in nascent stages^{26,36} and mostly focused on T1D-specific behaviors rather than those important for general health such as MVPA. A major strength of the study was the enrollment of a diverse sample of adolescents in greatest need of support for MVPA and T1D self-management (lower

income, people of color, elevated HbA1c).³⁷ This study had no A1c upper limit, thus incorporating a group of youth with T1D at high risk of dysglycemia that has been largely excluded from research studies. Further supporting generalizability, instructors from a diversity of backgrounds were able to facilitate with similar competencies. Intervention attendance was excellent and overcame logistical barriers related to alignment of group schedules, availability of MVPA supplies, timing MVPA with insulin and diet, and provision of real-time guidance. Given the focus on feasible uptake of technology and inclusion of youth with all HbA1c levels, we enrolled participants even if they were not consistently wearing CGM at enrollment, which meant glycemic outcomes could only be assessed by combining HbA1c and GMI values in an ancillary analysis. HbA1c and GMI can be discordant due to the timeframe they reflect and non-glycemic factors that can impact HbA1c.³¹ Nonetheless, glycemic reductions were seen during and after the intervention. These findings were encouraging, especially in light of the elevated HbA1c levels in the cohort at baseline, as these youth are largely excluded from research.

Prior studies supporting MVPA for adolescents with T1D provided self-management education and decision guidance^{32,38-40} and only one included personal support, which came from key family members such as a parent.³² In real-world contexts, 91% of adolescents with T1D report that their parents discourage MVPA.⁷ Support from peers and peer mentors, however, was rated as important by adolescents with T1D^{41,42} and associated with increased diabetes self-care.⁴³ Peer support for adolescents with T1D should be applied to MVPA and other general health metrics, whereas to date it has only been applied to T1D-specific behaviors.^{26-28,36} Such restriction of peer support to T1D-specific behaviors has been negatively associated with diabetes self-care,⁴³ similar to the “nagging” perceived from parents.⁴⁴ It may also be less engaging, as these prior peer interventions have been almost entirely restricted to asynchronous communications without physical activities.^{27,28} When synchronous chats have been added, just 39%-48% of participants attended one or more sessions.^{28,36} Overall, the present study shows the promise of peer group MVPA as a novel strategy both for supporting MVPA and generally engaging adolescents with T1D.

365 The primary finding of the study was that the virtual, compared to our previous in-person
 366 protocol,²⁴ greatly increased feasibility while maintaining excellent safety standards. Compared to our
 367 in-person group, we doubled recruitment uptake proportion from 16% to 35%, which is comparable to
 368 37% seen in personalized MVPA prescription for T1D that did not require fitting a group schedule.³² We
 369 also expanded from clinic to social media venues. In addition, we improved the attendance from 56% to
 370 85%-90%, while also doubling the frequency of sessions from once to twice per week. A glucose
 371 monitoring protocol was followed with no adverse events and occurrence of MVPA-induced
 372 hypoglycemia no more frequently than typical daily living. The program was also highly cost-efficient.
 373 Social media advertisements utilized community postings rather than paid advertisements, and virtual
 374 technology was all within the scope of devices normally owned by diverse families of teens with T1D.
 375 Videogame technology is popular among teens and substantially more so among those from lower-
 376 income families,⁴⁵ the smartwatch was the most economy-grade on the market (retail \$80 USD), and
 377 T1D was monitored by standard technology within the scope of the standard of care (CGM, fingerstick
 378 glucose and ketone meters).

379 The high acceptability of the in-person protocol²⁵ was also maintained based on mean
 380 satisfaction scores, though unbalanced across subscales. Intervention delivery competencies that could
 381 be directly delivered from young adult role model instructors – sharing T1D and MVPA knowledge and
 382 experiences and building T1D and MVPA self-management skills – achieved an excellent rating at
 383 68%-78% of sessions. By contrast, those that depended upon teen-to-teen interactions were achieved
 384 excellently much less often. Similarly, instructor interactions were the most highly rated and most
 385 important component of the program to teens. Perceived group cohesion was also rated highly by the
 386 teens, reflecting some combination of perceived cohesion to the instructors and other teens. Program
 387 structure and comfort were rated less highly than instructor interactions and perceived group cohesion,
 388 though nonetheless received ratings of 4+ out of 5 and were likely essential to the program's feasibility.

389 The technological aspects of the virtual MVPA experience received more modest ratings.
 390 Participants generally reported that the Ring Fit game functioned well in terms of clear goals, feedback,

391 and control. Concordantly, we observed that they achieved heart rate in the MVPA range as previously
 392 reported for other videogames that elicit full body movements and engage large muscle groups (i.e.,
 393 active videogames).¹⁹ However, participants were more neutral about the components that reflect
 394 enjoyment of the game which include sensory experience, challenge, and difficulty. In summary, the
 395 advantages of our program were effectiveness at connecting teens to young adult role models in a way
 396 that was logistically feasible, time efficiency for the role models since they could interact with a group all
 397 at once, and sustainment of favorable MVPA and clinical metrics. The relative limitations were the
 398 capability to facilitate peer interactions and have an enjoyable MVPA experience. It may therefore be
 399 beneficial to integrate our curriculum into more immersive technology such as virtual reality headsets
 400 allowing interactions through avatars.

401 Guidelines for safe MVPA with T1D provide general rules for maintaining glycemic stabilization
 402 through the adjustment of factors like diet, insulin, and type of MVPA.³⁻⁵ However, they presently lack
 403 adequate real-time support for their implementation. They acknowledge that they need a trial-and-error
 404 refinement process for each individual; the existing knowledge does not provide a one-size-fits-all
 405 approach nor can the needed refinements to it be accomplished within the scope of clinical
 406 appointments. Therefore, youth with T1D need to be given support during the time of MVPA in order to
 407 interpret and overcome the glycemic and psychosocial challenges that arise with the sometimes
 408 unpredictable glycemic responses to MVPA and the opportunity to build problem-solving skills “in the
 409 moment”.

410 Thus, there is a need for interventions that not only facilitate MVPA, but do so with a
 411 combination of behavior-change theory and practical considerations. First, from a practical standpoint,
 412 teens with T1D could benefit from regular interactions with their clinicians so they can ask questions
 413 about glycemic changes related to specific MVPA experiences, but most clinicians lack the time and
 414 resources to make this commitment. In the present study, the time efficiency of this process was
 415 quadrupled due to the group setting. Second, basic resources such as neighborhoods with safe play
 416 space are often inequitably accessed across socioeconomic groups.³⁷ In the present study, resources

417 were accessible by a diverse group of participants and socioeconomic levels. Third, advanced
 418 resources including peer support and role model mentoring are typically even less available than the
 419 basic guidance to teens with T1D due to the rarity of the condition, and guidance given in schools and
 420 through sports is not T1D-specific. Fourth, in our study, the activities and group process were driven by
 421 participants who could individually and collaboratively choose MVPA goals, giving them more flexibility
 422 and individualization than a prescribed curriculum.

423 The study had some limitations. First, it was a single-group study evaluating the feasibility and
 424 acceptability of the intervention with planned outcome metrics pertaining to safety and the most
 425 promising program components but not efficacy. Second, there are concerns about Fitbit heart rate
 426 measurements as two studies have reported they were less accurate for those with darker skin tone.⁴⁶
 427 However, Fitbit was mainly used for step-counting which does not have this limitation, and heart rate
 428 was only used for an assessment of convergent validity with the validated videogame.¹⁹ Third, while
 429 participants perceived high group cohesion, they had minimal verbal interactions with each other.
 430 Instructors were trained in group facilitation strategies derived from our expert planning group
 431 discussions and pilot sessions, but for future studies we will utilize an accredited group facilitation
 432 course. Peer interactions might also be enhanced by interoperable virtual reality technology that
 433 creates a more immersive experience. Finally, the study was limited to those who were able to afford
 434 Internet access at home, those who had television/computer screens, and time to attend the sessions.

435 CONCLUSIONS

436 Overall, we demonstrated the feasibility of facilitating a virtual network for MVPA, self-
 437 management education and guidance, diabetes-specific management skills, and peer and role model
 438 support for adolescents with T1D. We combined a fitness videogame, wearable step-tracking, a
 439 telehealth glucose safety monitoring protocol, diabetes self-management education, and skits designed
 440 to address diabetes-specific management skills. Successes included attraction of a diverse group, high
 441 attendance at the sessions, T1D self-management guidance that promoted safe MVPA, and meaningful
 442 connections to role-model instructors including shared experiences and skills in T1D self-management

and coping. The challenge was promoting immersive interaction among the adolescents. Therefore, future trials testing efficacy and moderators warrant upgrades of delivery to include more immersive virtual reality technologies, and financial investment to integrate and interoperate the components of our curriculum: MVPA, discussion, skits, viewership of glucose, and insulin dosing.

DATA AVAILABILITY

The individual participant data, study protocol, statistical analysis plan, informed consent form, and analysis code used to support the findings of this study are available from the corresponding author upon request. The data will be shared with researchers who provide a methodologically sound proposal to achieve the aims of the approved proposal. To gain access, data requesters must sign a data access agreement which will be drafted by Yale University and include the data requester's proposal as an exhibit.

CONFLICTS OF INTEREST

G.I.A. has in the last 3 years received grant support (to his institution) from the Patterson Trust, American Heart Association, National Institutes of Health, and Veterans Health Administration. He has also received professional services from Fitscript and Calm, on projects separate from the present study. S.N. receives grant support (to her institution) from the National Institutes of Health. C.T. is a consultant for Calm. M.S. is employed as a senior product manager by Abbott Laboratories and receives restricted stock units. S.A.W. currently receives grant support (to his institution) from Abbott Laboratories and the National Institutes of Health. In the last 3 years he has received honoraria for serving as a speaker for Abbott and Dexcom and as a consultant for Zealand Pharma. The Yale Children's Diabetes Center receives free CGM supplies through the Abbott Laboratories clinical sample program, which are distributed to patients including some who participated in this study. L.M.N. receives funding for research from the National Institutes of Health and Medtronic Diabetes. She is also a consultant for Medtronic, WebMD, and Calm. The authors attest that the Patterson Trust, American Heart Association, National Institutes of Health, Veterans Health Administration, Fitscript, Calm, Abbott Laboratories, Dexcom, Zealand Pharma, Medtronic, and WebMD had no influence on the design of this

study or its outcomes. The authors conducted the research outside of their responsibilities and affiliations with these entities.

FUNDING STATEMENT

The study and G.I.A. were supported by American Heart Association Grant #852679 (G.I.A., 2021–2024) and a Robert E. Leet and Clara Guthrie Patterson Trust Mentored Research Award, Bank of America, N.A., Trustee. G.I.A. and L.M.N. were supported by the National Institute of Diabetes, Digestive, and Kidney Diseases of the National Institutes of Health under mentored research scientist development awards (K01DK129441, K23DK128560). The study was further supported by the National Institutes of Health under the Yale Diabetes Research Center (P30DK045735). None of these entities were involved in the manuscript writing, editing, approval, or decision to publish.

ACKNOWLEDGMENTS

The authors thank the families and participants that help made this research possible. We also acknowledge Sa’Ra Skipper and Juanita Montoya for their assistance with conducting the study. Study data were collected and managed using REDCap electronic data capture tools hosted at Yale University.

AUTHORSHIP CONFIRMATION/CONTRIBUTION STATEMENT

GIA: Conceptualization (lead), data curation (lead), formal analysis (lead), funding acquisition (lead), investigation (equal), methodology (equal), project administration (lead), resources (lead), software (support), supervision (lead), validation (equal), visualization (lead), writing – original draft (lead). **SN:** Conceptualization (equal); investigation (equal); methodology (equal); writing – review and editing (equal). **MSK:** Methodology (supporting); Writing – review and editing (equal). **ADH:** Investigation (equal); methodology (equal); project administration (equal); software (equal); writing, review, and editing (supporting). **CT:** Investigation (equal); methodology (equal); writing, review, and editing (equal). **AC:** Investigation (equal); methodology (equal); writing – review and editing (equal). **MS:** Investigation (equal); methodology (equal); writing, review, and editing (equal). **JSB:** Methodology (supporting); Writing – review and editing (equal). **SAW:** Conceptualization (equal); investigation (equal);

methodology (equal); writing – review and editing (equal); writing – review and editing (equal). **LMN:**
 Conceptualization (equal), formal analysis (equal), investigation (equal), methodology (lead), software
 (lead), visualization (equal), writing – review and editing (lead).

498

499

500

REFERENCES

1. Foster NC, Beck RW, Miller KM, et al. State of Type 1 Diabetes Management and Outcomes from the T1D Exchange in 2016-2018. *Diabetes technology & therapeutics*. 2019;21(2):66-72.
2. Nathan DM, Group DER. The diabetes control and complications trial/epidemiology of diabetes interventions and complications study at 30 years: overview. *Diabetes care*. 2014;37(1):9-16.
3. Adolfsson P, Taplin CE, Zaharieva DP, et al. ISPAD Clinical Practice Consensus Guidelines 2022: Exercise in children and adolescents with diabetes. *Pediatr Diabetes*. 2022;23(8):1341-1372.
4. Moser O, Riddell MC, Eckstein ML, et al. Glucose management for exercise using continuous glucose monitoring (CGM) and intermittently scanned CGM (isCGM) systems in type 1 diabetes: position statement of the European Association for the Study of Diabetes (EASD) and of the International Society for Pediatric and Adolescent Diabetes (ISPAD) endorsed by JDRF and supported by the American Diabetes Association (ADA). *Pediatric diabetes*. 2020;21(8):1375-1393.
5. Riddell MC, Gallen IW, Smart CE, et al. Exercise management in type 1 diabetes: a consensus statement. *The lancet Diabetes & endocrinology*. 2017;5(5):377-390.
6. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV. Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. *The New England journal of medicine*. 1986;315(4):215-219.
7. Jabbour G, Henderson M, Mathieu ME. Barriers to Active Lifestyles in Children with Type 1 Diabetes. *Canadian journal of diabetes*. 2016;40(2):170-172.
8. Ryninks K, Sutton E, Thomas E, Jago R, Shield JP, Burren CP. Attitudes to Exercise and Diabetes in Young People with Type 1 Diabetes Mellitus: A Qualitative Analysis. *PloS one*. 2015;10(10):e0137562.
9. Elmesmari R, Reilly JJ, Martin A, Paton JY. Accelerometer measured levels of moderate-to-vigorous intensity physical activity and sedentary time in children and adolescents with chronic disease: A systematic review and meta-analysis. *PloS one*. 2017;12(6):e0179429.
10. Tapia-Serrano MA, Sevil-Serrano J, Sanchez-Miguel PA, Lopez-Gil JF, Tremblay MS, Garcia-Hermoso A. Prevalence of meeting 24-Hour Movement Guidelines from pre-school to adolescence: A systematic review and meta-analysis including 387,437 participants and 23 countries. *J Sport Health Sci*. 2022;11(4):427-437.
11. Chinchilla P, Dovic K, Braune K, et al. Perceived Knowledge and Confidence for Providing Youth-Specific Type 1 Diabetes Exercise Recommendations amongst Pediatric Diabetes Healthcare Professionals: An International, Cross-Sectional, Online Survey. *Pediatric Diabetes*. 2023;2023:8462291.
12. Benzing V, Schmidt M. Exergaming for Children and Adolescents: Strengths, Weaknesses, Opportunities and Threats. *Journal of clinical medicine*. 2018;7(11):422. doi: 410.3390/jcm7110422.
13. Council On C, Media. Media Use in School-Aged Children and Adolescents. *Pediatrics*. 2016;138(5).
14. Akhter K, Turnbull T, Simmons D. A systematic review of parent/peer-based group interventions for adolescents with type 1 diabetes: interventions based on theoretical/therapeutic frameworks.

541

- 542 *Br J Diabetes*. 2018;18:51-65.
- 543 15. Prochaska JJ, Sallis JF, Long B. A physical activity screening measure for use with adolescents
544 in primary care. *Archives of Pediatrics & Adolescent Medicine*. 2001;155(5):554-559.
- 545 16. Warburton DE, Gledhill N, Jamnik VK, et al. Evidence-based risk assessment and
546 recommendations for physical activity clearance: Consensus Document 2011. *Applied
547 physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*.
548 2011;36 Suppl 1:S266-298.
- 549 17. Currie C IJ, Molcho M, Lenzi M, Veselska Z & Wild F ed *Health Behaviour in School-aged
550 Children (HBSC) Study Protocol: Background, Methodology and Mandatory items for the
551 2013/14 Survey*. St Andrew: CAHRU.
- 552 18. Petersen AC, Crockett L, Richards M, Boxer A. A self-report measure of pubertal status:
553 Reliability, validity, and initial norms. *Journal of youth and adolescence*. 1988;17(2):117-133.
- 554 19. Comeras-Chueca C, Villalba-Heredia L, Pérez-Llera M, et al. Assessment of Active Video
555 Games' Energy Expenditure in Children with Overweight and Obesity and Differences by
556 Gender. *International journal of environmental research and public health*. 2020;17(18):6714.
557 doi: 6710.3390/ijerph17186714.
- 558 20. Fuller D, Colwell E, Low J, et al. Reliability and Validity of Commercially Available Wearable
559 Devices for Measuring Steps, Energy Expenditure, and Heart Rate: Systematic Review. *JMIR
560 mHealth and uHealth*. 2020;8(9):e18694.
- 561 21. Allen NA. Social cognitive theory in diabetes exercise research: an integrative literature review.
562 *The Diabetes educator*. 2004;30(5):805-819.
- 563 22. Adolfsson P, Riddell MC, Taplin CE, et al. ISPAD Clinical Practice Consensus Guidelines 2018:
564 Exercise in children and adolescents with diabetes. *Pediatric diabetes*. 2018;19 Suppl 27:205-
565 226.
- 566 23. Staiano AE, Abraham AA, Calvert SL. Motivating effects of cooperative exergame play for
567 overweight and obese adolescents. *Journal of diabetes science and technology*. 2012;6(4):812-
568 819.
- 569 24. Ash GI, Joiner KL, Savoye M, et al. Feasibility and safety of a group physical activity program
570 for youth with type 1 diabetes. *Pediatric diabetes*. 2019.
- 571 25. DeJonckheere M, Joiner KL, Ash GI, et al. Youth and Parent Perspectives on the Acceptability
572 of a Group Physical Activity and Coping Intervention for Adolescents With Type 1 Diabetes. *Sci
573 Diabetes Self Manag Care*. 2021;47(5):367-381.
- 574 26. Titoria R, Amed S, Tang TS. Peer Support Interventions on Digital Platforms for Children With
575 Type 1 Diabetes and Their Caregivers. *Diabetes Spectr*. 2022;35(1):26-32.
- 576 27. Walker AF, Haller MJ, Gurka MJ, et al. Addressing health disparities in type 1 diabetes through
577 peer mentorship. *Pediatr Diabetes*. 2020;21(1):120-127.
- 578 28. Troncone A, Cascella C, Chianese A, et al. Psychological support for adolescents with type 1
579 diabetes provided by adolescents with type 1 diabetes: The chat line experience. *Pediatric
580 diabetes*. 2019;20(6):800-810.
- 581 29. Staiano AE, Adams MA, Norman GJ. Motivation for Exergame Play Inventory: Construct validity
582 and relationship to game play. *Cyberpsychology: Journal of Psychosocial Research on
583 Cyberspace*. 2019;13(3):Article 7.
- 584 30. Chin WW, Salisbury WD, Pearson AW, Stollak MJ. Perceived Cohesion in Small Groups:
585 Adapting and Testing the Perceived Cohesion Scale in a Small-Group Setting. *Small Group
586 Research*. 1999;30(6):751-766.
- 587 31. Piona C, Marigliano M, Mozzillo E, et al. Evaluation of HbA1c and glucose management
588 indicator discordance in a population of children and adolescents with type 1 diabetes. *Pediatr
589 Diabetes*. 2022;23(1):84-89.
- 590 32. Faulkner MS, Michaliszyn SF, Hepworth JT, Wheeler MD. Personalized exercise for
591 adolescents with diabetes or obesity. *Biological research for nursing*. 2014;16(1):46-54.
- 592 33. Quirk H, Blake H, Tennyson R, Randell TL, Glazebrook C. Physical activity interventions in

children and young people with Type 1 diabetes mellitus: a systematic review with meta-analysis. *Diabetic medicine : a journal of the British Diabetic Association*. 2014;31(10):1163-1173.

34. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Medicine and science in sports and exercise*. 2008;40(1):181-188.

35. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. *J Exp Psychol Gen*. 2012;141(1):2-18.

36. Bisno DI, Reid MW, Pyatak EA, et al. Virtual Peer Groups Reduce HbA1c and Increase Continuous Glucose Monitor Use in Adolescents and Young Adults with Type 1 Diabetes. *Diabetes Technol Ther*. 2023;25(9):589-601.

37. Armstrong S, Wong CA, Perrin E, Page S, Sibley L, Skinner A. Association of Physical Activity With Income, Race/Ethnicity, and Sex Among Adolescents and Young Adults in the United States: Findings From the National Health and Nutrition Examination Survey, 2007-2016. *JAMA pediatrics*. 2018.

38. Wong CH, Chiang YC, Wai JP, et al. Effects of a home-based aerobic exercise programme in children with type 1 diabetes mellitus. *Journal of Clinical Nursing*. 2011;20(5-6):681-691.

39. Marrero DG, Fremion AS, Golden MP. Improving compliance with exercise in adolescents with insulin-dependent diabetes mellitus: results of a self-motivated home exercise program. *Pediatrics*. 1988;81(4):519-525.

40. Shetty VB, Soon WHK, Roberts AG, et al. A Novel Mobile Health App to Educate and Empower Young People With Type 1 Diabetes to Exercise Safely: Prospective Single-Arm Mixed Methods Pilot Study. *JMIR Diabetes*. 2021;6(4):e29739.

41. Ye CY, Jeppson TC, Kleinmaus EM, Kliems HM, Schopp JM, Cox ED. Outcomes That Matter to Teens With Type 1 Diabetes. *The Diabetes educator*. 2017;43(3):251-259.

42. Xie LF, Housni A, Nakhla M, et al. Adaptation of an Adult Web Application for Type 1 Diabetes Self-management to Youth Using the Behavior Change Wheel to Tailor the Needs of Health Care Transition: Qualitative Interview Study. *JMIR Diabetes*. 2023;8:e42564.

43. Doe E. An analysis of the relationships between peer support and diabetes outcomes in adolescents with type 1 diabetes. *J Health Psychol*. 2018;23(10):1356-1366.

44. Luyckx K, Seiffge-Krenke I, Missotten L, Rassart J, Casteels K, Goethals E. Parent-adolescent conflict, treatment adherence and glycemic control in Type 1 diabetes: the importance of adolescent externalising symptoms. *Psychol Health*. 2013;28(9):1082-1097.

45. Priceonomics Data S. Gender, Income & Education: Who Plays Video Games? In. Vol 20212018.

46. Koerber D, Khan S, Shamsheri T, Kirubarajan A, Mehta S. Accuracy of Heart Rate Measurement with Wrist-Worn Wearable Devices in Various Skin Tones: a Systematic Review. *J Racial Ethn Health Disparities*. 2022:1-9.