medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Article 1

Gut microbiome and lichen sclerosus: a two-sample 2 bi-directional Mendelian randomization study 3

Jiayan Chen^{1,†}, Peiyan Wang^{2,†}, Changji Xiao³, Kalibinuer Kelaimu³, Xianshu Gao¹, Xiaomei Li¹, Jun 4 Hu^{3*} 5

6 Department of Radiation Oncology, Peking University First Hospital, Beijing, China; chenjiayan2019-ro@bjmu.edu.cn (J.-Y.C.);

7 8 doctorgaoxs@126.com (X.-S.G.); lixiaomei1973@bjmu.edu.cn (X.-M.L.);

School of Information, University of Michigan, Ann Arbor, MI, United States; peiyanw@umich.edu (P.-Y.W.);

9 ³ Department of Obstetrics and Gynecology, Peking University First Hospital, Beijing, China; xcji1988@163.com (C.-J.X.);

10 13400052009@163.com (K.K.)

11 Correspondence: hujunpkufh@bjmu.edu.cn (J.H.)

These authors contributed equally to this work. 12

Abstract: (1) Background: Recent studies suggest a potential link between gut microbiomes (GMs) and inflammatory diseases, but 13 14 the role of GMs in lichen sclerosus (LS) remains unclear. This study aims to investigate the causal relationship between GMs and LS, 15 focusing on key GM taxa. (2) Methods: We utilized GWAS summary statistics for 211 GM taxa and their association with 2,445 LS patients and 353,088 healthy controls, employing Mendelian randomization (MR). GWAS data for GM taxa came from the 16 17 MiBioGen consortium, and for LS from the FinnGen consortium. The primary analytical tools included the inverse-variance 18 weighted (IVW) method, weighted MR, simple mode, weighted median, and MR-Egger methods. Sensitivity analyses included 19 leave-one-out analysis, MR-Egger intercept test, MR-PRESSO global test, and Cochrane's O-test. A reverse MR analysis was 20 conducted on bacteria identified in the forward MR study. (3) Results: We identified one strong causal relationship: order 21 Burkholderiales [odds ratio (OR) = 0.420, 95% confidence interval (CI): 0.230 - 0.765, p = 0.005], and three nominally significant relationships: phylum Cyanobacteria (OR = 0.585, 95% CI: 0.373 - 0.919, p = 0.020), class Betaproteobacteria (OR = 0.403, 95% 22 23 CI: 0.189 - 0.857, p = 0.018), and genus Butyrivibrio (OR = 0.678, 95% CI: 0.507 - 0.907, p = 0.009). Moreover, this MR analysis 24 was not impacted by horizontal pleiotropy, according to the MR-Egger intercept test and MR-PRESSO global test (p > 0.05). 25 Remarkably, the reliability of our results was confirmed by leave-one-out analysis. Reverse MR analysis showed no significant 26 causal relationship between LS and GM. (4) Conclusions: This MR study identifies specific gut flora linked to a lower risk of LS, 27 offering new insights for disease treatment and prevention. Future research should incorporate metagenomics sequencing of extensive microbiome GWAS datasets. 28

29 Keywords: gut microbiome, lichen sclerosus, causal relationship, Mendelian randomization, genome-wide association study 30

31 1. Introduction

32 Lichen sclerosus (LS) is a chronic mucocutaneous immune-mediated disease that is characterized by skin atrophy 33 and hypopigmentation [1]. A majority of the LS patients, exceeding 80%, present with anogenital lesions. Common 34 Typical extragenital sites involve neck, shoulders, upper trunk, and thighs [2]. The typical clinical presentation consists of persistent white atrophic patches in the external genitalia, often accompanied by discomfort and itching. 35 Potentially, the lesions could worsen and cause the vaginal introitus to constrict and scar, which would impair 36 37 urogenital and sexual function. Furthermore, women with vulvar LS have a high risk of squamous cell carcinoma with 38 a lifetime incidence rate ranging from 0.4% to 6% [3-4]. The prevalence of genital LS was reported to be 1.7% in 39 gynecological practice [5]. However, the precise incidence of LS remains understated. LS can manifest at any age and 40 has an impact on both sexes, with a prevalence ratio of females to males ranging from 3:1 to 10:1 [6]. Postmenopausal women are most commonly affected, followed by men, prepubertal children, and adolescents [7]. 41

42 It is still difficult to fully comprehend the pathophysiology of LS. LS is characterized by a hereditary and familial propensity; however, chronic irritation, endocrine status, and autoimmune diseases may also contribute to the 43 44 pathogenesis [8-11]. The microbiota is a key immune system modulator that contributes significantly to homeostasis 45 maintenance. Over the past 20 years, numerous investigations have been carried out to examine the constitution and 46 the inflammatory function of intestinal bacterial flora in preclinical and clinical studies focusing on dermatologic 47 conditions, such as alopecia areata, rosacea, atopic dermatitis, and psoriasis [12-14]. Certain notable associations between microbiome signatures have been discovered. The relationship between LS and gut microbiota, however, is 48 49 poorly unversion and the second the second the second the second the second the second second the second se 50 genesis of LS.

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

In epidemiological research, the Mendelian randomization (MR) method, which employs genetic variants as an instrumental variable, is widely used to investigate, the potential causal effect of exposure to specific diseases. In the MR approach, the genetic alleles could be randomly distributed like a randomized controlled trial, which is less likely to be impacted by confounding or reverse causation. Hence, we conducted a bidirectional large-scale MR analysis to identify potentially influential GM taxa using summary data from GWAS in order to shed light on the prevention and treatment of LS.

57 2. Materials and Methods

58 2.1. Study design

Figure 1 shows the design of the two-sample MR analysis. We followed three important MR assumptions to determine the possible causal association between GM and LS risk: IVs must be correlated with GM, they must be independent of confounding variables, and IVs must only affect LS risk through GM [15].



62

63 **Figure 1.** Study design and workflow.

64 *2.2. Exposure data*

16S fecal microbiome rRNA gene sequencing profiles and host genotypes of 18,340 participants were examined
 in-depth by the MiBioGen consortium [16]. A comprehensive GWAS encompassing 211 GM taxa from genus to phylum
 revealed genetic variations linked to 9 phyla, 16 classes, 20 orders, 35 families, and 131 genera. The GM GWAS
 summary statistics can be accessed by downloading them from https://mibiogen.gcc.rug.nl/ [17–19].

69 2.3. Outcome data

We acquired LS-related GWAS summary statistics from the FinnGen consortium R9 release, encompassing 2,445 cases and 353,088 controls, available at https://r9.finngen.fi/. It is noteworthy that our study, relying on publicly available databases, did not require additional ethics approval or informed consent.

73 2.4. Instrumental variable (IV) selection

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

In this MR study, single-nucleotide polymorphism (SNP) that exhibited strong association with each GM taxon were utilized as IVs. For a more comprehensive analysis, IVs achieving locus-wide significance ($p < 1 \times 10^{-6}$) were selectively employed. Concurrently, the PLINK clumping method was applied to exclude SNPs in linkage disequilibrium ($r^2 < 0.001$, kb = 10,000). The degree of weak instrumental bias was then estimated using the F-statistic of IVs, where a value of >10 indicated the lack of bias resulting from weak IVs [20]. Palindromic SNPs and those not found in the outcome were finally eliminated from the instrumental variable group.

80 2.5. Statistical analysis

81 The IVW approach, an extension of the Wald ratio estimator based on meta-analysis principles, was the main 82 technique used in this MR investigation to infer causality [21]. Four more MR methods—MR-Egger, weighted median, 83 weighted mode, and simple mode—were then utilized to supplement the IVW results if the IVW technique showed a 84 causal association (p < 0.05) for each GM taxon [22–23]. At the p-value of less than 0.05, causal relationships were 85 shown as OR with 95% CI. False discovery rate (FDR) correction was used with a threshold of q < 0.05 to address 86 multiple tests. Exposure-outcome couples exhibiting a consistent direction were required to demonstrate consistency across all MR techniques, demonstrating a causal link. Many sensitivity studies were carried out in order to evaluate 87 88 the stability of causal associations. The horizontal pleiotropy was determined using the MR-Egger intercept test and the MR-PRESSO global test [24-25]. Moreover, result robustness was evaluated using a leave-one-out analysis. Reverse MR studies were conducted to examine if identified major bacterial genera were causally impacted by LS, 89 90 91 employing SNPs linked with LS as IVs. The R software's "TwoSampleMR" and "MR-PRESSO" packages (version 4.2.2) 92 served as the foundation for all studies.

93 3. Results

94 3.1. IVs Overview

At the phylum, class, order, family, and genus levels, we found a total of 16, 26, 26, 54, and 149 SNPs linked to GM, with a significance level of $p < 1 \times 10^{-6}$. Notably, every IV showed stronger correlations between exposure and result (P_{exposure} < P_{outcome}), and every F-value was greater than 10. Supplementary Table S1 has comprehensive information about the IVs.

99 3.2. MR Analysis

In our implementation of the Inverse Variance Weighted Fixed Effects (IVW-FE) technique to assess the causal relationship between LS and 211 GM taxa in five hierarchical levels, several taxa were identified as potentially influencing LS risk. Specifically, phylum *Cyanobacteria* (id: 1500), class *Betaproteobacteria* (id: 2867), order *Burkholderiales* (id: 2874), and genus *Butyrivibrio* (id: 1993) exhibited a decreased risk for LS (Figure 2). Following FDR correction, the protective effect of order *Burkholderiales* (id: 2874) [OR= 0.420 (0.230 - 0.765), p = 0.005, q = 0.036] against LS remained significant. Importantly, Cochran's Q test results indicated an absence of heterogeneity.

To further support the causal relationship between these GM taxa and LS, we utilized four more MR techniques: MR-Egger, weighted median, weighted mode, and simple mode (Figure 3). The robustness of our observed causal relationships was reinforced by the encouraging consistency of the results obtained from these approaches with the IVW findings.

Exposure	Cochran's Q_pval	Method	nSNP			OR (95% CI)	P-value
phylum Cyanobacteria	0.208	IVW (RE)	2			0.585 (0.332 - 1.033)	0.065
(id: 1500)		IVW (FE)	2	—	-	0.585 (0.373 - 0.919)	0.020
class Betaproteobacteria	0.497	IVW (RE)	2	H		0.403 (0.189 - 0.857)	0.018
(id: 2867)		IVW (FE)	2	H		0.403 (0.189 - 0.857)	0.018
order Burkholderiales	0.811	IVW (RE)	3	H		0.420 (0.230 - 0.765)	0.005
(id: 2874)		IVW (FE)	3	—		0.420 (0.230 - 0.765)	0.005
genus Butyrivibrio	0.640	IVW (RE)	2	-	4	0.678 (0.507 - 0.907)	0.009
(id: 1993)		IVW (FE)	2		4	0.678 (0.507 - 0.907)	0.009
				0 0.5	1 1.	5	

110

111 Figure 2. MR results of causal effects between GM and LS risk.

112

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Exposure	Method		OR (95% CI)	P-value
phylum Cyanobacteria	IVW (RE)		0.585 (0.332 - 1.033)	0.065
(id: 1500)	IVW (FE)		0.585 (0.373 - 0.919)	0.020
class Betaproteobacteria	IVW (RE)	⊢ − −−−1	0.403 (0.189 - 0.857)	0.018
(id: 2867)	IVW (FE)	• •• •	0.403 (0.189 - 0.857)	0.018
order Burkholderiales	IVW (RE)		0.420 (0.230 - 0.765)	0.005
(id: 2874)	IVW (FE)	⊢ ∎i	0.420 (0.230 - 0.765)	0.005
	MR Egger	-	→ 0.224 (0.007 - 7.324)	0.555
	Simple mode		0.448 (0.197 - 1.017)	0.195
	Weighted median	—	0.422 (0.206 - 0.865)	0.018
	Weighted mode		0.407 (0.165 - 1.005)	0.191
genus Butyrivibrio	IVW (RE)	—	0.678 (0.507 - 0.907)	0.009
(id: 1993)	IVW (FE)	⊢ ∎(0.678 (0.507 - 0.907)	0.009
	(0 0.5 1	1.5	

113

114 Figure 3. Various MR outcomes for four GM taxa with identified causal associations to LS.

115 *3.3.Sensitivity analysis*

The MR-Egger intercept test results revealed no horizontal pleiotropy (p > 0.05) within the IVs associated with order *Burkholderiales* (id: 2874) and LS (Supplementary Table S2). Furthermore, the leave-one-out analysis demonstrated the robustness of the MR results, as the removal of any particular IV did not change the overall findings for order *Burkholderiales* (id: 2874) (Supplementary Figure S1). It is noteworthy, however, that due to an insufficient number of SNPs, MR-Egger intercept tests and leave-one-out analyses could not be conducted for the remaining three gut microbiota taxa. Similarly, the MR-PRESSO global test was precluded for the same reason.

122 3.4. Reverse MR

Nine SNPs linked to LS were found to be eligible IVs after a thorough series of IV screening procedures (Supplementary Table S3). The findings of the reverse MR analysis, displayed in Supplementary Table S4, yielded no proof of a causal effect from LS to the determined bacterial taxa. The results of the MR-Egger intercept test and the MR-PRESSO global test both indicated that there was no horizontal pleiotropy among the remaining IVs. A comprehensive presentation of the sensitivity tests can be found in Supplementary Table S5, demonstrating the robustness of our findings in the face of various analytical challenges.

129 4. Discussion

130 The gut microbiota is one of the potential targets for regulating the host's immune responses. The correlation between the composition of the gut microbiome and the immunological system regulation has provided a novel 131 132 perspective on the pathogenesis of numerous chronic multifactorial diseases. According to multiple studies that link 133 gastrointestinal homeostasis to skin disorders, both the component and function of the GM are altered in patients with autoimmune and inflammatory skin diseases. Atopic dermatitis (AD) development and progression may be influenced 134 135 by alterations in the microbiota diversity. For instance, decreased variation of the GM is associated with more server 136 disease. Recent evidence suggests that microbiota could be involved in modifying itchiness in atopic dermatitis through the interactions between the gut, skin and brain. Damage to the epidermal barrier, brain sensitization of 137 pruritus-generating mechanisms, and modulation of histamine-independent itch are caused by the presence of 138 139 proinflammatory cytokines, microbial metabolites, and a compromised immune response [26]. Microbiota dysbiosis 140 was found to induce an irregular immune response in individuals with psoriasis, and the alterations of microbiota 141 composition were linked to the level of inflammation-related biomarkers that were abnormal in the patients with 142 psoriasis. For example, the relative abundances of *Phascolarctobacterium* and *Dialister* may serve as possible 143 indicators of the psoriasis progression, as the IL-2 receptor showed a positive correlation with the former and a 144 negative correlation with the latter [27].

145 However, only a few studies have shown a link between changes in GM and LS disease. Chattopadhyay et al. 146 conducted a pilot case-control study with prepubescent females, which suggested that the females with vulvar LS 147 exhibited an increased relative abundance of Escherichia coli, Bifidobacterium adolescentis, Akkermansia muciniphila, 148 Clostridiales spp., Paraprevotella spp., and Dialister spp in comparison to healthy controls, while the abundance of Roseburia faecis and Ruminococcus bromii was noticeably decreased. These findings showed a potential correlation 149 150 between intestinal dysbiosis and juvenile vulvar LS [28]. Nygaard et al.'s study revealed that women with LS had a higher relative abundance of phylum Euryarchaeota in their GM than controls [29]. However, the inconsistent 151 152 findings and small sample of these studies pose challenges to reaching generalizable conclusions. Additionally, due to variations in gender, race, age, and body region, the composition of the GM may also differ among studies. The 153 154 presence of these unclear factors impedes the ability to establish a causal relationship between GM and the risk of LS 155 disease.

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

156 Our work thoroughly evaluated the causative influence of 211 GM taxa (from phylum to genus level) on LS, 157 emphasizing the importance of GMs in LS. Four causal links in all were found, three of which were nominal and one of 158 which was strong. Our MR analysis has provided the first confirmation that class *Betaproteobacteria* (id. 2867), genus Butyrivibrio (id.1993), order Burkholderiales (id.2874), phylum Cyanobacteria (id.1500) have a protective effect LS. 159 160 Most of these bacteria belong to Phylum Firmicutes and Proteobacteria, which was consistent with our previous research with vulvar skin microbiota [30]. It indicates an interesting linkage between gut and genital skin microbiota. 161 162 As far as we know, this is the first MR investigation to look into the possibility of a causal relationship between gut microbiota and LS. In accordance with researches in other inflammatory diseases, our results suggest that gut 163 dysbiosis may play a role in the development of LS. 164

It was believed that the metabolic activity and immunological response of gut microbiome may affect skin 165 166 conditions [31]. The gut barrier integrity is primarily maintained by the microbial communities, through the conversion of polysaccharides and vitamins, as well as short-chain fatty acids. For example, butyrate reduces 167 168 interstitial barrier permeability and improves the integrity of the epithelial barrier. Genus Butyrivibrio, class 169 Betaproteobacteria, and order Burkholderiales are commensal bacteria that are well recognized for their capacity to 170 generate short-chain fatty acids, particularly butyrate, which could help maintain systemic immunity balance [32]. Butyrate serves as the main source of energy for the colonic epithelium and exerts anti-inflammatory effects in the 171 colonic mucosa by preventing the activation of NF-kB [33]. Several studies have shown that patients with Crohn's 172 173 disease and ulcerative colitis, respectively, have a decreased enrichment of Butyrivibrio in colon and saliva samples 174 than healthy controls, respectively [34-35]. Additionally, lower concentrations of stool butyrate have been observed in 175 those suffering from inflammatory bowel disease (IBD), providing more evidence to support the possibility that gut dysbiosis plays a causal role in inflammatory conditions like IBD and potentially LS. Nevertheless, the relationship 176 between skin disease and immunological activity triggered by the gut microbiome remains obscure, and further 177 178 investigation is needed.

179 Our work contributed substantial evidence to the current literature concerning the causal relationship between 180 gut microbiota and LS. The primary benefits of our research are: (1) We analyzed genetic data from a large sample group, which increases the reliability of the results in comparison to smaller observational research. (2) Through MR 181 182 analysis, confounding variables are eliminated from causal association, and the causal association identified in our 183 study may serve as potential candidates for future functional investigations. This study has some limitations that 184 should be mentioned. The primary limitation of our work is that the most of participants in our research were limited to people who had European origins. Although reducing bias caused by population heterogeneity would be achieved, 185 186 further research is necessary to explore whether the MR results can be general in other populations. Second, in order to obtain a greater number of SNPs, we relaxed the p threshold, which may heighten the likelihood of contravening the 187 188 basic MR design premise. However, all SNPs have F statistic greater than 10, which indicates that all weak SNPs were 189 excluded from the MR estimation. And rigorous FDR correction was applied to identify the significant results to 190 reduce the likelihood of false-positive findings. Third, we were unable to fully reduce pleiotropy due to the fact that the specific biological functions of the utilized SNPs are still unknown. However, it is encouraging to note that distinct MR 191 192 models generated consistent estimates, and the analyses of sensitivity analyses under various assumptions were 193 unable to identify any horizontal pleiotropy. Studies have indicated that LS patients frequently exhibit several 194 comorbidities, such as depression and behavioral disorders. The prevalence rates of these comorbidities may reach 195 37% and 63%, respectively [36-37]. Many of these comorbidities have been demonstrated to have an association with the GM [38]. Therefore, it is exceedingly probable that variations in GM exist between the patients with LS alone and 196 197 those with LS accompanied by other medical conditions. However, the original GWAS data failed to make the 198 distinction, thus additional research is required to solve this limitation.

199 5. Conclusions

Our study serves as a foundation for recognizing the causal association between the gut microbiota and LS. In addition, it was found that a number of gut flora may lower the incidence of LS. These findings hold promise for their prospective use in the prevention and management of LS. This would have an implication to clinicians that microbiome-targeted therapies might be promising preventive and therapeutic tools for LS. Future investigation is required to unravel the fundamental mechanism. Additional observational studies or lab-based research is warranted to substantiate these results.

- Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Figure S1: title;
 Table S1: title; Video S1: title.
- Author Contributions: Jiayan Chen, Peiyan Wang, and Jun Hu designed the research. Peiyan Wang collected and analyzed the
- 209 data. Xianshu Gao and Xiaomei Li offered technical advice. Jiayan Chen, Peiyan Wang, and Jun Hu drafted the manuscript. Jun Hu 210 supervised the study. Changji Xiao and Kalibinuer Kelaimu revised the manuscript. All authors approved the submitted version of 211 the article and participated in its writing.
- Funding: This research was funded by Peking University Youth Physician Research Funding (2015QN031) and National High Level Hospital Clinical Research Funding (Scientific Research Seed Fund of Peking University First Hospital 2023SF08).
- 214 Data Availability Statement: The summary data of MiBioGen can be downloaded from the website <u>https://mibiogen.gcc.rug.nl/</u>.
 215 The summary data of FINNGEN can be downloaded from the website <u>https://r9.finngen.fi/</u>. This published work and its
 216 supplementary information files contain other datasets that were generated and/or analyzed during the current study, which are
 217 accessible to the public.

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

- 218 Acknowledgments: The researchers of MiBioGen and FinnGen, as well as the members of all GWAS cohorts that were used in 219 this study, are appreciated by the authors for providing the GWAS summary statistics.
- 220 Conflicts of Interest: The authors declare no conflicts of interest.

221

medRxiv preprint doi: https://doi.org/10.1101/2024.07.23.24310907; this version posted July 24, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

222 References

- 223 224 225 226 227 228 229 230 Papini, M, Russo, A, Simonetti, O, Borghi, A, Corazza, M, Piaserico, S, et al. Mucous membrane disorders research group of 1. SIDeMaST. Diagnosis and management of cutaneous and anogenital lichen sclerosus: recommendations from the Italian Society of Dermatology (SIDeMaST). Ital J Dermatol Venerol. 2021,156:519–533.
 - Kaufman, RH, DiPaola, GR, Friedrich, EG Jr, Hewitt, J, and Woodruff, JD. New nomenclature for vulvar disease. J Cutan 2. Pathol. 1976,3:159-161.
 - Chamli A, Souissi A. Lichen Sclerosus. Treasure Island (FL): StatPearls; 2022. 3.
 - 4. Meani R, Howard A, Veysey E. Incidence of vulval squamous cell carcinoma in women with vulval lichen sclerosus in an Australian tertiary referral centre. The Australasian journal of dermatology. 2019,60(1):76-7
- 231 232 233 234 235 Bieber AK, Steuer AB, Melnick LE, Wong PW, Pomeranz MK, Autoimmune and Dermatologic Conditions Associated with 5. Lichen Sclerosus, Journal of the American Academy of Dermatology. 2020, 85(1):228-229.
 - Kirtschig, G, Becker, K, Günthert, A, Jasaitiene, D, Cooper, S, Chi, CC, et al. Evidence-based (S3) guideline on (anogenital) 6. lichen sclerosus. J Eur Acad Dermatol Venereol. 2015, 29(10):e1-43.
- 236 237 238 239 Krapf, JM, Mitchell, L, Holton, MA, and Goldstein, AT. Vulvar lichen Sclerosus: current perspectives. Int J Women's Health. 7. 2020,12:11-20.
 - 8. Etiology, Clinical Features, and Diagnosis of Vulvar Lichen Sclerosus: A Scoping Review. Obstet Gynecol Int. 2020,2020:7480754
 - Vulvar Lichen Sclerosus from Pathophysiology to Therapeutic Approaches: Evidence and Prospects. Biomedicines. 9. 2021,9(8):950.
 - Birenbaum DL, Young RC. High prevalence of thyroid disease in patients with lichen sclerosus, J Reprod Med. 2007,52:28-30 10. The prevalence of self-reported medical comorbidities in patients with vulvar lichen sclerosus: A single-center retrospective 11.
 - study. Int J Gynaecol Obstet. 2021,153(2):340-343.
- 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 255 255 257 258 259 260 Mahmud MR, Akter S, Tamanna SK, Mazumder L, Esti IZ, Banerjee S, Akter S, Hasan MR, Acharjee M, Hossain MS, Pirttilä 12. AM. Impact of gut microbiome on skin health: gut-skin axis observed through the lenses of therapeutics and skin diseases. Gut Microbes. 2022 Jan-Dec;14(1):2096995.
 - Moniaga CS, Tominaga M, Takamori K. An Altered Skin and Gut Microbiota Are Involved in the Modulation of Itch in Atopic 13. Dermatitis. Cells. 2022 Dec 5;11(23):3930.
 - Buhas MC, Gavrilas LI, Candrea R, Cătinean A, Mocan A, Miere D, Tătaru A. Gut Microbiota in Psoriasis. Nutrients. 2022 Jul 14. 20:14(14):2970.
 - Davies, N. M., Holmes, M. V., and Davey Smith, G. (2018). Reading Mendelian randomisation studies: a guide, glossary, and 15. checklist for clinicians. BMJ 362:k601. doi: 10.1136/bmj.k601
 - Kurilshikov A, Medina-Gomez C, Bacigalupe R, et al. Large-scale association analyses identify host factors influencing human 16. gut microbiome composition. Nat Genet. 2021 Feb;53(2):156-165. doi: 10.1038/s41588-020-00763-1.
 - Swertz MA, Jansen RC. Beyond standardization: dynamic software infrastructures for systems biology. Nat Rev Genet. 2007 17. Mar;8(3):235-43. doi: 10.1038/nrg2048.
 - 18. Swertz MA, Dijkstra M, Adamusiak T, et al. The MOLGENIS toolkit: rapid prototyping of biosoftware at the push of a button. BMC Bioinformatics. 2010 Dec 21;11 Suppl 12(Suppl 12):S12. doi: 10.1186/1471-2105-11-S12-S12.
 - van der Velde KJ, Imhann F, Charbon B, et al. MOLGENIS research: advanced bioinformatics data software for 19. non-bioinformaticians. Bioinformatics. 2019 Mar 15;35(6):1076-1078. doi: 10.1093/bioinformatics/bty742.
- 260 261 262 263 264 20. Palmer, T. M., Lawlor, D. A., Harbord, R. M., Sheehan, N. A., Tobias, J. H., Timpson, N. J., et al. (2012). Using multiple genetic variants as instrumental variables for modifiable risk factors. Stat. Methods Med. Res. 21, 223–242. doi: 10.1177/0962280210394459
- 265 Pagoni P, Dimou NL, Murphy N, Stergiakouli E. Using Mendelian randomisation to assess causality in observational studies. 21. Evid Based Ment Health. 2019 May;22(2):67-71. doi: 10.1136/ebmental-2019-300085.
 - 22. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. Genet Epidemiol. 2016 May;40(4):304-14. doi: 10.1002/gepi.21965.
 - Burgess S, Thompson SG. Interpreting findings from Mendelian randomization using the MR-Egger method. Eur J Epidemiol. 23. 2017 May;32(5):377-389. doi: 10.1007/s10654-017-0255-x.
 - Rees JMB, Wood AM, Burgess S. Extending the MR-Egger method for multivariable Mendelian randomization to correct for 24. both measured and unmeasured pleiotropy. Stat Med. 2017 Dec 20;36(29):4705-4718. doi: 10.1002/sim.7492.
 - Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from 25. Mendelian randomization between complex traits and diseases. Nat Genet. 2018 May;50(5):693-698. doi: 10.1038/s41588-018-0099-7
 - Moniaga CS, Tominaga M, Takamori K. An Altered Skin and Gut Microbiota Are Involved in the Modulation of Itch in Atopic 26. Dermatitis. Cells. 2022 Dec 5;11(23):3930
 - Zhang, X.; Shi, L.; Sun, T.; Guo, K.; Geng, S. Dysbiosis of Gut Microbiota and Its Correlation with Dysregulation of Cytokines in Psoriasis Patients. BMC Microbiol. 2021, 21, 78. 27.
 - 28. Chattopadhyay S, Arnold JD, Malayil L, Hittle L, Mongodin EF, Marathe KS, et al. (2021) Potential role of the skin and gut microbiota in premenarchal vulvar lichen sclerosus: A pilot case- control study. PLoS ONE 16(1): e0245243.
 - Nygaard S, Gerlif K, Bundgaard-Nielsen C, Saleh Media J, Leutscher P, Sørensen S, Brusen Villadsen A, Thomsen Schmidt 29. Arenholt L. The urinary, vaginal and gut microbiota in women with genital lichen sclerosus - A case-control study. Eur J Obstet Gynecol Reprod Biol. 2023 Oct;289:1-8.
- 2666 2677 2688 2699 2700 2711 2722 2733 2744 2755 2766 2777 2788 2799 2800 2811 2822 2833 2844 2855 2866 Liu X, Zhuo Y, Zhou Y, Hu J, Wen H and Xiao C (2022) Analysis of the Vulvar Skin Microbiota in Asymptomatic Women and 30. Patients With Vulvar Lichen Sclerosus Based on 16S rRNA Sequencing. Front. Cell Dev. Biol. 10:842031.
- 287 Salem I, Ramser A, Isham N, Ghannoum MA. The gut microbiome as a major regulator of the gut-skin axis. Front Microbiol. 31. 288 2018;9:1459.
- 289 32. Tamanai-Shacoori Z, Smida I, Bousarghin L, Loreal O, Meuric V, Fong SB, et al. Roseburia spp.: a marker of health? Future 290 Microbiol. 2017; 12:157-70.
- 291 Segain JP, Raingeard de la Ble tière D, Bourreille A, Leray V, Gervois N, Rosales C., et al. Butyrate inhibits inflammatory 33. 292 responses through NFkappaB inhibition: implications for Crohn's disease. Gut. 2000; 47:397-403

- Masoodi I, Alshanqeeti AS, Ahmad S, Alyamani EJ, Al-Lehibi AA, Qutub AN, ALSyari KN, Alomair AO.Microbial dysbiosis in 34. inflammatory bowel diseases: results of a metagenomic study in Saudi Arabia. Minerva Gastroenterol Dietol. 2019 Sep;65(3):177-186.
- 293 294 295 296 297 298 299 300 301 302 Gryaznova MV, Solodskikh SA, Panevina AV, Syromyatnikov MY, Dvoretskaya YD, Sviridova TN, Popov ES, Popov 35. VN.Heliyon. Study of microbiome changes in patients with ulcerative colitis in the Central European part of Russia. Helivon.2021 Mar 10;7(3):e06432.
- Bell SG, Kobernik EK, Haefner HK, Welch KC.Association Between Vulvar Squamous Intraepithelial Lesions and Psychiatric 36.
- IllnessJ Low Genit Tract Dis. 2021 1;25(1):53-56. Malandrone F, Bevilacqua F, Merola M, Gallio N, Ostacoli L, Carletto S, Benedetto C.The Impact of Vulvar Cancer on 37. Psychosocial and Sexual Functioning: A Literature ReviewCancers (Basel). 2021 Dec 23;14(1):63.
- 303 38. Socała, K., Doboszewska, U., Szopa, A., Serefko, A., Włodarczyk, M., Zielińska, A., et al. (2021). The role of 304 microbiota-gut-brain axis in neuropsychiatric and neurological disorders. Pharmacol. Res. 172:105840.
- 305 Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual
- 306 author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury
- 307 to people or property resulting from any ideas, methods, instructions or products referred to in the content.