

A machine learning approach to predict mortality and pulmonary hypertension severity in newborns with congenital diaphragmatic hernia

Mortality and pulmonary hypertension prediction in congenital diaphragmatic hernia

Luana Conte^{1,2†}, Ilaria Amodeo^{3†}, Giorgio De Nunzio^{2,4*}, Genny Raffaeli³, Irene Borzani⁵, Nicola Persico^{6,7}, Alice Griggio⁸, Giuseppe Como³, Mariaros Colnaghi³, Monica Fumagalli^{3,6}, Donato Cascio^{1†}, Giacomo Cavallaro^{3†}

¹Department of Physics and Chemistry, Università degli Studi di Palermo, Palermo, Italy

²Advanced Data Analysis in Medicine (ADAM), Laboratory of Interdisciplinary Research Applied to Medicine (DReAM), Local Health Authority (ASL) Lecce and Università del Salento, Lecce, Italy

³Neonatal Intensive Care Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

⁴Department of Mathematics and Physics "E. De Giorgi", Laboratory of Biomedical Physics and Environment, Università del Salento, Lecce, Italy

⁵Pediatric Radiology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

⁶Department of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy

⁷Department of Obstetrics and Gynecology, Fondazione IRCCS Ca' Granda, Ospedale Maggiore Policlinico, Milan, Italy

⁸Department of Obstetrics and Gynecology, ASST Fatebenefratelli Sacco, Ospedale Macedonio Melloni, Milan, Italy

***Corresponding Author.** Giorgio De Nunzio, giorgio.denunzio@unisalento.it

† Co-First Author: Luana Conte and Ilaria Amodeo

† Co-Last Author: Donato Cascio and Giacomo Cavallaro

Luana Conte: luana.conte@unipa.it, ORCID: <https://orcid.org/0000-0002-8741-3478>

Ilaria Amodeo: ilaria.amodeo@policlinico.mi.it, ORCID: <https://orcid.org/0000-0002-2650-0084>

Giorgio De Nunzio*: giorgio.denunzio@unisalento.it, ORCID: <https://orcid.org/0000-0002-1998-0286>

Genny Raffaeli: genny.raffaeli@policlinico.mi.it, ORCID: <https://orcid.org/0000-0001-9175-9394>

Irene Borzani: irene.borzani@policlinico.mi.it, ORCID: <https://orcid.org/0000-0003-0386-4712>

Nicola Persico: nicola.persico@unimi.it, ORCID: <https://orcid.org/0000-0001-9028-9597>

Alice Griggio: alicegriggio89@gmail.com, ORCID: <https://orcid.org/0000-0002-2947-4627>

Giuseppe Como: pinocomo@gmail.com, ORCID: <https://orcid.org/0009-0009-2120-0427>

Mariaros Colnaghi: mariaros.colnaghi@policlinico.mi.it, ORCID:

<https://orcid.org/0000-0002-0186-0710>

Donato Cascio: donato.cascio@unipa.it, ORCID: <https://orcid.org/0000-0001-6522-1259>

Giacomo Cavallaro: giacomo.cavallaro@policlinico.mi.it, ORCID: <https://orcid.org/0000-0002-4921-1437>

45 **ABSTRACT**

46 Prenatal prediction of postnatal outcomes in newborns with congenital diaphragmatic hernia (CDH)
 47 remains challenging, especially for mortality and neonatal persistent pulmonary hypertension
 48 (PPHN). Despite the increasing utilization of advanced artificial intelligence (AI) technologies in the
 49 neonatal field, this study is pioneering in exploring AI methodologies in the context of CDH. It
 50 represents an initial attempt to implement a Machine Learning (ML) system to predict postnatal
 51 mortality and PPHN severity, using prenatal and early postnatal data as input variables. We enrolled
 52 50 patients with isolated left-sided CDH from singleton pregnancies and retrospectively collected
 53 clinical and imaging variables from fetal ultrasound (US) and shape features extracted from magnetic
 54 resonance imaging (MRI), combined with gestational age and birth weight. A supervised ML model
 55 for predicting mortality and PPHN severity was developed, achieving good accuracy (88% for
 56 mortality prediction and 82% for PPHN) and sensitivity (95% for mortality and 85% for PPHN). The
 57 area under the curve (AUC) of the ROC curve was 0.88 for mortality and 0.82 for PPHN predictions.
 58 Our results may lead to novel AI applications in the neonatal field, focusing on predicting postnatal
 59 outcomes based on prenatal data, ultimately improving prognostic assessments and intervention
 60 strategies for such a complex disease.

61 **Clinical Trial Registration:** The trial was registered at ClinicalTrials.gov with Identifier
 62 NCT04609163

63

64 **Keywords:** Newborn, Congenital Diaphragmatic hernia, Neonatal Persistent Pulmonary
 65 Hypertension, Mortality, Machine Learning, Deep Learning.

66

67

68 **What is Known:** prenatal prediction of postnatal mortality and severity of pulmonary hypertension
 69 in CDH newborns remains challenging and largely based on imaging through the volumetric
 70 assessment of fetal lungs.

71

72 **What is New:** developing a ML system for predicting PPHN severity and mortality risk based on the
 73 integrated assessment of prenatal and early postnatal variables is feasible, with good accuracy.

74

INTRODUCTION

Congenital diaphragmatic hernia (CDH) is a rare congenital anomaly characterized by incomplete closure of the diaphragm and herniation of abdominal organs into the chest, resulting in pulmonary hypoplasia, neonatal persistent pulmonary hypertension, and cardiac dysfunction [1–3]. CDH occurs in nearly 1 in 2500 births. Several factors influence the prognosis, such as defect size and location, associated anomalies, presence of liver up in the thorax, and gestational age at birth [4,5]. Risk stratification is essential to identify patients who might benefit from specific interventions and to enable a risk-adjusted analysis of outcomes, healthcare costs, and management approaches. Prenatal and postnatal CDH predicting tools have largely increased and have been validated during the last years based on clinical and instrumental data [6,7,16–18,8–15]. However, a universal risk stratification method has not been identified yet, and an agreed-upon set of risk-specific management guidelines is still lacking [19].

In particular, predicting the severity of Neonatal Persistent Pulmonary Hypertension (PPHN) using conventional prenatal diagnostic methods remains challenging. As a result, there is growing interest in leveraging advanced technologies that favor a timely and accurate prognosis.

Artificial Intelligence (AI) is increasingly applied in the neonatal field to support medical data analysis. Predictive algorithms are being developed using traditional Machine Learning (ML) approaches as well as its more advanced Deep Learning (DL) extension. ML and DL can process and analyze medical data, including images from different sources such as ultrasound, magnetic resonance imaging (MRI), and X-ray. Integrating these algorithms into healthcare systems holds promise for enhancing diagnostic accuracy and disease pattern classification. These algorithms could help predict specific outcomes, guide interventions, and improve the overall quality of care [20,21,30,22–29].

However, to our knowledge, these methodologies still need to be successfully applied to newborns with CDH. The aim of our study was to provide a predictive algorithm for mortality and PPHN in CDH based on the integrated analysis of prenatal and early postnatal data.

101 MATERIALS AND METHODS

102 *Study design*

103 This study represents an exploratory secondary analysis of a retrospective cohort study performed at
 104 Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy (CLANNISH, Clinical
 105 Trials identification n°: NCT04609163) [29]. The study involved the following services: the Fetal
 106 Surgery Center, Pediatric Radiology Service, and Neonatal Intensive Care Unit (NICU). Moreover,
 107 the Department of Mathematics and Physics at the Università del Salento (Lecce, Italy) and the
 108 Department of Physics and Chemistry at the Università degli Studi di Palermo (Palermo, Italy)
 109 developed the AI algorithms.

110 The current study adhered to the principles of good clinical practice and followed the guidelines of
 111 the Helsinki Declaration. It received approval from the local ethics committee (Milan Area 2, Italy)
 112 with approval number/ID 800_2020bis. However, considering its retrospective design, the ethics
 113 committee waived the need for informed consent. The study was also registered on ClinicalTrials.gov
 114 with the identifier NCT04609163.

116 *Patients*

117 The study population, inclusion-exclusion criteria, and a comprehensive description of the primary
 118 study design were previously published and are briefly summarized here [29]. Inborn CDH patients
 119 born between 01/01/2016 and 30/04/2020 admitted to the NICU at birth were included. The take-
 120 charge of the mothers took place at our Fetal Surgery Center at a gestational age of 30+6 weeks or
 121 below. Non-isolated CDH and twin pregnancies were excluded. Only left-sided CDH were
 122 considered because of their larger numerosity, homogeneity, and variability in liver position, leaving
 123 out right-sided CDH.

124 A total of 50 patients were included in the final study population.

126 *Data Collection*

127 Clinical maternal and fetal prenatal variables were retrospectively collected using Astraia software
 128 (Astraia Software GmbH, Ismaning, Germany) and NeoCare software (GPI SpA, Trento, Italy). A
 129 prenatal ultrasound (US) performed between 25+0 and 30+6 weeks of gestation was considered for
 130 each patient. In the case of fetal endoscopic tracheal occlusion (FETO), the fetal US was performed
 131 before the fetal procedure. Additionally, native sequences from fetal MRI were gathered for 36 out
 132 of 50 cases, with separate acquisitions for the lung and liver. The imaging software employed for this
 133 study was Synapse PACS and Synapse 3D (FUJIFILM Medical Systems, Lexington, MA, US). Lung
 134 volumes were computed using T2 HASTE sequences, selecting the best-quality image plane without
 135 motion-induced artifacts [31]. On the other hand, liver volumes were calculated based on T1 VIBE
 136 sequences [32]. An experienced pediatric radiologist (IB) freehand delineated Regions Of Interest
 137 (ROIs) to define the areas of the left and right lungs and the liver, excluding the pulmonary hila and
 138 mediastinal structures for each slice. Organ volumes were calculated using the software.
 139 Subsequently, the DICOM (Digital Imaging and Communications in Medicine) files were
 140 anonymized and converted to the NIfTI (neuroimaging informatics technology initiative) format for
 141 easy manipulation.

142 *Clinical and Imaging variables*

143 A detailed summary of the clinical and imaging variables included is reported in Table 1.

144

Table 1. Clinical and imaging variables
<i>Maternal data</i> Mother age Ethnicity Number of gestations ending the outcome for fetuses with an Assisted Reproduction (MAR): yes/no Antenatal use of corticosteroids: yes/no Premature rupture of membranes (pPROM): yes/no Gestational age at pPROM Gestational age at CDH diagnosis
<i>Fetal ultrasound data</i> Gestational age Estimated Fetal Weight the (EFW) Amniotic Fluid (AF) Umbilical Artery Pulsatility Index

Pulmonary Flow Pulsatility Index Pulmonary flow Peak Systolic Velocity Peak early diastolic reversed flow Organ herniation of Bowel, Stomach, Liver: yes/no Observed/expected lung-to-head ratio (tracing method) CDH severity: mild, moderate, severe Fetal Endoscopic Tracheal Occlusion (FETO): yes/no Gestational age at balloon insertion and removal
<i>MRI data</i> Gestational age Apparent diffusion coefficient (ADC) Right and left lung volume Total fetal lung volume (TFLV) Observed/expected TFLV (O/E TFLV%) Total liver volume Herniated Liver volume Percentage of liver herniation (%LH) Mediastinal shift angle (MSA) Apparent diffusion coefficient (ADC) of the left and right lung
<i>Early postnatal data</i> Gestational age at birth Birth weight

Radiomics features

In addition to the clinical variables, standard 3D radiomics features were extracted from the segmented ROIs in the MRI using the freely available and open-source Pyradiomics v. 3.01 software tool [30,33]. Pyradiomics produces many variables, with and without preprocessing by various filters and optional reslicing, with different interpolators. Only features from the original images without preprocessing were considered in this work. Due to significant dissimilarity in the gray-level content of the MRI scans, only shape features were utilized (MeshVolume, VoxelVolume, SurfaceArea, SurfaceVolumeRatio, Sphericity, Maximum3DDiameter, MajorAxisLength, MinorAxisLength, LeastAxisLength, Elongation, and Flatness). The geometric meaning of each feature is detailed in Table 2.

Table 2. Geometric meaning of Pyradiomics shape features	
Feature	Geometric Meaning

MeshVolume	The shape volume, calculated using a mesh representation. It is the three-dimensional space enclosed by the surface of the object.
VoxelVolume	The volume calculated by counting the number of voxels (3D pixels) within the shape and multiplying by the volume of a single voxel. This represents the discretized volume of the object.
SurfaceArea	The total area of the surface of the shape. This quantifies the two-dimensional extent of the object's surface.
SurfaceVolumeRatio	The ratio of surface area to volume. This measure indicates how 'compact' an object is; lower values suggest a more compact shape.
Sphericity	A measure of how spherical (round) the object is. Perfect spheres have a sphericity of 1. Lower values indicate less spherical shapes.
Maximum3DDiameter	The largest distance between any two points on the surface of the shape. This is the maximum length of the object in any dimension.
MajorAxisLength	The length of the major axis of the shape, which is the longest dimension in the principal component analysis (PCA) of the object.
MinorAxisLength	The length of the minor axis, which is perpendicular to the major axis and is the second longest dimension in the PCA of the object.
LeastAxisLength	The shortest axis length from the PCA of the object. It's perpendicular to both the major and minor axes.
Elongation	The ratio of the minor axis length to the major axis length. It indicates how much longer the shape is in one direction compared to the other.
Flatness	The ratio of the least axis length to the major axis length. This measure indicates how 'flat' or 'elongated' an object is compared to being spherical.

157

158 Variables computed from the gray levels were discarded, avoiding additional image manipulation,
159 such as intensity standardization. A total of 80 features were considered: 56 prenatal variables, 2 very
160 early postnatal variables as gestational age and birth weight, and 22 MRI-extracted shape features (11
161 from the lungs, 11 from the liver).

162 To ensure fairness in the classification process, the features were normalized to the 0-1 range using
163 min-max normalization on the training set. The same normalization parameters were then applied to
164 the validation set samples. However, in some cases, the lack of MRI data resulted in missing values
165 in the features extracted by Pyradiomics. This was also observed for non-radiomics features based on

the patient's diagnostic pathway. Imputation by a weighted average was considered [34] to handle these missing values, as in Equation 1:

$$f_n(m) = \frac{\frac{\overline{f_n^{(1)}}}{\sigma_n^{(1)}} + \frac{\overline{f_n^{(2)}}}{\sigma_n^{(2)}}}{\frac{1}{\sigma_n^{(1)}} + \frac{1}{\sigma_n^{(2)}}} \quad (1)$$

where $f_n(m)$ is the value to be assigned to the (missing) n -th feature for the m -th sample, $\overline{f_n^{(1)}}$ and $\overline{f_n^{(2)}}$ are the average values of the n -th features for classes 1 and 2, respectively, and $\sigma_n^{(1)}$ and $\sigma_n^{(2)}$ are the corresponding standard deviations. This way, in the approximation of Gaussian distributions, a neutral value for the distributions of the two classes is used as the missing feature.

Target variables: neonatal persistent pulmonary hypertension and mortality

For each included patient, a neonatologist performed a systematic revision of the first available echocardiogram within 24 hours after birth, focusing on direct and indirect signs of pulmonary hypertension. Data collection was focused on the presence and characteristics of the shunts through patent ductus arteriosus and *foramen ovale*, the characteristics of the intraventricular sept, the estimation of the systolic pulmonary artery pressure through tricuspid valve regurgitation, the systemic pressures, and concomitant use of pulmonary vasodilators. Patients were then stratified according to the presence and severity of pulmonary hypertension into two categories: severe (over-systemic, considered as the positive class) vs moderate/mild (iso/under-systemic). According to mortality, the study population was divided into non-survivors (positive class) vs survivors.

Feature selection

We employed the Recursive Feature Elimination (RFE) technique with Cross-Validation (CV), specifically using the Leave One Patient Out CV (LOPO-CV) scheme. The RFE method starts with

the entire feature set and recursively removes the minor essential features based on a chosen metric (in this case, accuracy) until the desired number of features is reached. Typically, the final number of features to select is a parameter that needs to be specified. This parameter was determined dynamically by varying its value and calculating the corresponding accuracy in our approach. We then selected the parameter value that maximized accuracy. The Random Forest (RF) classifier was used to evaluate the various configurations.

Training

To exploit the available samples as much as possible, we used a LOPO-CV scheme. In this method, we selected one patient as the validation set while using the remaining patients for training. We trained several classifiers and obtained performance metrics such as the confusion matrix, sensitivity, specificity, area under the ROC curve (AUC), and the area under the P-R curve. All the optimization steps were based on maximizing accuracy.

Classifiers

Three classification algorithms were tested: eXtreme Gradient Boosting (XGBoost), Support Vector Machine (SVM), and K-Nearest Neighbors (KNN). The first classifier was used because it natively and effectively deals with missing clinical values. The choice of the other two classifiers was due to their ability to allow good performance in conditions of a limited number of available samples, as they are characterized by a reduced number of parameters to be tuned [35]. Hyperparameter tuning was performed to avoid overfitting and improve model performance.

RESULTS

The final study population consisted of 50 patients: 26 severe (52%) and 24 moderate/mild (48%) cases of PPHN. According to mortality, 37 survivors (74%) and 13 non-survivors (26%) were present.

As regards mortality analysis, the feature selection procedure led to the choice of 10 out of 80 features, in particular: maternal age, gestational age at CDH diagnosis, CDH severity (mild, moderate, severe), centile of estimated fetal weight (EFW), observed/expected lung to head ratio (o/e LHR) by tracing method, umbilical artery pulsatility index, left and right fetal lung volume (FLV) at MRI, observed/expected total fetal lung volume (o/e TFLV), gestational age at birth, and birth weight. Remarkably, no shape features were selected. Table 3 shows the classification results obtained for mortality prediction.

Table 3. Classification figures of merit for mortality, computed in a LOPO scheme				
<i>Classifier</i>	<i>AUC</i>	<i>Accuracy</i>	<i>Sensitivity</i>	<i>Specificity</i>
XGBoost	0.88	88%	95%	69%
SVM	0.78	80%	97%	31%
KNN	0.78	82%	95%	46%
LOPO: Leave One Patient Out; KNN: K-Nearest Neighbors; SVM: Support Vector Machine; XGBoost: eXtreme Gradient Boosting.				

The results of the different classification methods were comparable regarding AUC and accuracy, though XGboost had better performance. Figure 1 shows the ROC and P-R curves for mortality prediction obtained by XGboost with feature selection. The trained model correctly identified 88% of cases and achieved a sensitivity of 95% and a specificity of 69%. The AUC from the ROC curve was 0.87, while the P-R curve subtended an area of 0.95 (with the frequency of positive cases equal to 52%). From the P-R curve, precision drops after 50% sensitivity but remains more than 85% when sensitivity is 90%, and even if we require complete sensitivity, precision remains relatively high (around 80%).

Fig 1. Mortality prediction. Left: ROC curve, right: P-R curve, obtained with the XGBoost classifier on prenatal clinical variables. No shape features extracted from MRIs were used, as required by the feature selection procedure.

As far as PH is concerned, feature selection led to the identification of the 14 features, in particular: gestational age at CDH diagnosis, liver position, grading of stomach herniation, gestational age at US ultrasound, centile of EFW, umbilical artery pulsatility index, peak early diastolic reversed flow, o/e TFLV, apparent diffusion coefficient of left lung, original shape elongation, gestational age at birth, and birth weight.

In this case, both clinical and shape features were selected. Table 4 reports the classification results obtained for PPHN classification. The results produced by the different classification methods show that XGboost also performed better in this case.

The XGboost classifier demonstrated significantly superior classification capabilities to the other two classifiers (Tables 3 and 4).

Table 4. Classification figures of merit for PPHN, computed in a LOPO scheme				
Classifier	AUC	Accuracy	Sensitivity	Specificity
XGBoost	0.82	82%	85%	79
SVM	0.75	74	79	71
KNN	0.65	54	53	33
LOPO: Leave One Patient Out; KNN: K-Nearest Neighbors; PPHN: Neonatal Persistent Pulmonary Hypertension; SVM: Support Vector Machine; XGBoost: eXtreme Gradient Boosting.				

Figure 2 shows the ROC and P-R curves for PPHN classification obtained by XGboost with the features selection. The AUC from the test ROC curve was 0.82, while the P-R curve subtended an area of 0.75 (with the frequency of positive cases equal to 26%). The trained model correctly identified 82% of cases and achieved a sensitivity of 85% and a specificity of 79%. From the P-R

curve, we deduce that even at very high sensitivity values (about 83-84%), precision is more than 80%.

Fig 2. Left: the ROC curve; right: the Precision-Recall (P-R) curve for PPHN predictions with the XGBoost classifier on prenatal clinical variables and shape features extracted from MRIs.

DISCUSSION

Congenital diaphragmatic hernia (CDH) is a life-threatening anomaly requiring high-skilled and multidisciplinary team of experts for appropriate management since from antenatal diagnosis [36].

Despite advancements over time, morbidity and mortality remain significant (20–40%), even within high-volume tertiary referral centers [37–39]. An estimated quarter of survivors experience neurodevelopmental impairments across all domains, encompassing motor and sensory (hearing, visual) deficits as well as cognitive, language, and behavioral impairments [40].

CDH patients exhibit varying degrees of pulmonary hypoplasia and abnormal pulmonary vascular disease, resulting in varying extents of pulmonary hypertension. Up to 30–40% of newborns with CDH experience concomitant cardiac ventricular dysfunction [41,42]. PPHN is associated with adverse outcomes in CDH patients, underscoring the critical nature of its management in the care of these infants [43].

Various clinical and laboratory parameters and prognostic indices in the perinatal period have been subject to study to predict postnatal outcomes [39,44–47]. The identification of variables predictive of mortality is paramount for clinical decision-making and parental guidance. The o/e LHR and o/e TFLV serve as pivotal metrics in this regard. Each of these parameters evaluates the extent of pulmonary hypoplasia associated with CDH, a critical determinant of both survival and long-term prognosis.

The o/e LHR has been widely studied and utilized in the prediction of postnatal survival in cases of isolated CDH. Jani et al. highlighted the significance of the o/e LHR in predicting survival in fetuses

with isolated diaphragmatic hernia [48]. Snoek et al. further assessed the predictive value of the o/e LHR for survival and chronic lung disease (CLD) in survivors with left-sided CDH, reflecting its ongoing relevance in an era of standardized neonatal treatment [49]. Their multicenter study underscores the evolving understanding of o/e LHR in predicting outcomes for CDH patients.

On the other hand, the o/e TFLV exhibits a stronger correlation with postnatal outcomes than the absolute lung volume. Moreover, a growing body of evidence supports the superior accuracy of o/e TFLV in predicting survival compared to ultrasound-based estimations of lung size, which may not fully account for the ipsilateral lung and could thus underestimate the effective lung volume [50–55].

In cases of isolated CDH, o/e TFLV has demonstrated efficacy in distinguishing survival, with an o/e TFLV < 25% being associated with more severe forms and a reduced survival rate [54,56–60].

Furthermore, o/e TFLV has been shown to forecast the necessity for extracorporeal membrane oxygenation (ECMO) after birth, with the combined assessment of lung volumetry and o/e LHR proving more effective than ultrasound alone in predicting the need for ECMO [61–64].

Prenatal prediction of PPHN plays a crucial role in prenatal management, delivery planning and postnatal care. However, while both o/e LHR and o/e TFLV offer insights into the extent of CDH-associated pulmonary hypoplasia, their predictive value for PPHN necessitates careful consideration.

Our findings support the possibility of successfully developing a ML system for predicting PPHN severity and mortality risk based on the integrated assessment of prenatal and early postnatal variables.

To achieve our goal, we enrolled 50 left-sided CDH cases. The dataset was relatively balanced concerning PPHN, with 26 severe and 24 moderate/mild cases, whereas mortality classes included 37 survivors and 13 non-survivors. We combined prenatal clinical and imaging data with gestational age and weight at birth, which both play a key role in survival in neonatal patients, especially those in critical conditions. In addition, standard 3D radiomics features were extracted from the segmented ROIs using the freely available Pyradiomics software tool. This software package facilitated automatic reslicing with a selected interpolator and computed multiple radiomics variables. As the

302 MRIs exhibited significant variations in grayscales, which would have required some form of
303 intensity standardization to use features based on gray values, we only utilized shape features to avoid
304 additional image manipulation and discarded variables based on the gray levels.

305 A feature selection phase was executed for both postnatal target variables, mortality, and PPHN. The
306 RFE approach used RF classification to evaluate different configurations, which was appropriate for
307 several reasons. First, this approach provides features of relative importance during the training
308 process. Each time a decision tree is constructed, the model tracks how much each feature contributes
309 to reducing the cost function, usually Gini impurity or entropy. An importance score for each feature
310 is obtained by averaging this importance across all trees. Second, because of its "forest" nature, a RF
311 is robust and less prone to overfitting than individual decision trees. This means that the computed
312 importance of features is more reliable and less affected by noise in the data. Finally, RFs can handle
313 highly correlated features without special preprocessing. In the presence of correlations, this approach
314 can distribute importance among correlated features, providing a complete picture of each feature
315 contribution.

316 We conducted a comprehensive evaluation of three classification algorithms: XGBoost, SVM, and
317 KNN. Our models were trained using prenatal and early postnatal clinical variables, as well as
318 selected shape features extracted from MRI data. Interestingly, we discovered that XGBoost
319 outperformed the other models and emerged as the best classification model for both clinical targets.
320 The supervised ML models, designed to predict PPHN severity and neonatal mortality, showed
321 promising preliminary results. Our study suggests that predicting mortality and PPHN severity in the
322 prenatal and very early postnatal period can be feasible by ML applications, achieving accuracies of
323 88% for mortality and 82% for postnatal PPHN. With significant accuracy rates and reliable
324 sensitivity, this model has the potential to revolutionize prognostic assessment in CDH, eventually
325 improving patient outcomes. By implementing the algorithm, risk categories could be simulated
326 based on available prenatal data and assuming gestational age and estimated fetal weight at birth. The
327 algorithm could also be updated in real time at subsequent obstetric visits or based on the threat of

preterm delivery, as prematurity plays a significant role in survival, especially in infants with underlying disease. This would assist with parenting counseling, birth planning, and postnatal care. To the best of our knowledge, our studies are the first to explore the application of AI methods to CDH [29,30].

Despite being encouraging, some limitations must be considered. First, the restricted dataset deriving from the rarity of the condition represented the weakest point. An appropriate number of cases during training/validation and data interpretation is crucial for ML applications. Potential strategies may involve collaborating with other institutions and prospectively considering including future cases to augment the study population. Another critical aspect is data inhomogeneity, specifically the lack of a standard grayscale in the images. This would require a standardization procedure, after which gray-level-based features could be used to increase ML quality for classification purposes. Nonetheless, MRI standardization is a delicate process that involves profound changes in image gray levels, which might even make ML procedures less accurate. Consequently, we preferred to simply discard ML features based on the gray-level content of ROIs, only using shape features. Interestingly, no shape features were selected for the mortality target, whereas clinical and shape variables were chosen for the PPHN target. We can speculate that the information provided by the images is more closely related to the structure and architecture of the lung parenchyma, which directly impacts the disease's pathophysiology. On the other hand, mortality may be an indirect result of these structural alterations, influenced by many factors. Although a conclusive interpretation is not yet possible, this aspect deserves further investigation, and an increase in the study population and image optimization are crucial. Finally, the retrospective data collection is largely affected by missing or inaccurate data and may be time-consuming for the clinician. Standardized assessment and computerized data collection could improve the dataset quality.

CONCLUSIONS

354 Although with limitations, with reasonable accuracy, a ML approach for predicting mortality and
 355 PPHN severity of CDH newborns using prenatal and very early post-natal variables appears feasible.
 356 Our results could pave the way for new AI applications in the neonatal field. They would enable risk-
 357 adjusted analyses of outcomes, healthcare costs, and management strategies, ultimately improving
 358 the overall quality of care.

359

360 **Statements and Declarations**

361 **Competing Interests:** The authors have no relevant financial or non-financial interests to disclose.

362

363 **Funding:** This study was (partially) funded by the Italian Ministry of Health - Current Research
 364 IRCCS.

365

366 **Author Contributions:** L.C., I.A., G.D.N., G.R., I.B., D.C., and G.C. contributed to the study's
 367 conception and design; L.C., I.A., G.D.N., G.R., I.B., D.C., G.C. (Giuseppe Como), N.P., and G.C.
 368 (Giacomo Cavallaro) contributed to the study's methodology, investigation, and data curation; I.B.
 369 contributed to manual segmentation; L.C., G.D.N., L.C., and D.C. contributed to ML and DL
 370 analysis; L.C., G.D.N., and D.C. performed the statistical analysis; L.C., I.A., G.D.N., G.R., I.B.,
 371 D.C., and G.C. (Giacomo Cavallaro) wrote the initial draft preparation of the manuscript; L.C., I.A.,
 372 G.D.N., G.R., I.B., D.C., G.C. (Giuseppe Como), N.P., M.C., M.F., and G.C. (Giacomo Cavallaro)
 373 wrote, reviewed, and edited the manuscript; L.C., G.D.N., and D.C. contributed to designing the
 374 figures; G.D.N. and G.C. (Giacomo Cavallaro) contributed equally to the visualization of the
 375 manuscript; G.D.N. and G.C. (Giacomo Cavallaro) contributed to the supervision and project
 376 administration of the study. All authors have read and agreed to the published version of the
 377 manuscript.

378

379 **Ethics approval**

380 The present study was conducted using the principles of good clinical practice and the Helsinki
381 Declaration. It was approved by the local ethics committee (Milan Area 2, Italy) with approval
382 number 800_2020bis. However, due to the study's retrospective nature, the Ethics Committee waived
383 informed consent. The study was registered at ClinicalTrials.gov with the identifier NCT04609163.

384

385 **Consent to participate**

386 Written informed consent was obtained from the parents.

387

388 **Conflicts of Interest**

389 The authors declare that the research was conducted without any commercial or financial
390 relationships that could be construed as a potential conflict of interest.

391

392 **References**

393

- 394 1. Russo FM, De Coppi P, Allegaert K, Toelen J, van der Veecken L, Attilakos G, et al. Current
395 and future antenatal management of isolated congenital diaphragmatic hernia. *Semin Fetal*
396 *Neonatal Med.* 2017;22: 383–390. doi:10.1016/j.siny.2017.11.002
- 397 2. Keijzer R, Liu J, Deimling J, Tibboel D, Post M. Dual-Hit Hypothesis Explains Pulmonary
398 Hypoplasia in the Nitrofen Model of Congenital Diaphragmatic Hernia. *Am J Pathol.*
399 2000;156: 1299–1306. doi:10.1016/S0002-9440(10)65000-6
- 400 3. Pierro M, Thébaud B. Understanding and treating pulmonary hypertension in congenital
401 diaphragmatic hernia. *Semin Fetal Neonatal Med.* 2014;19: 357–363.
402 doi:10.1016/j.siny.2014.09.008
- 403 4. Snoek KG, Greenough A, van Rosmalen J, Capolupo I, Schaible T, Ali K, et al. Congenital
404 Diaphragmatic Hernia: 10-Year Evaluation of Survival, Extracorporeal Membrane
405 Oxygenation, and Foetoscopic Endotracheal Occlusion in Four High-Volume Centres.

Neonatology. 2018;113: 63–68. doi:10.1159/000480451

5. Harting MT, Lally KP. The Congenital Diaphragmatic Hernia Study Group registry update. Semin Fetal Neonatal Med. 2014;19: 370–375. doi:10.1016/j.siny.2014.09.004

6. Metkus AP, Filly RA, Stringer MD, Harrison MR, Adzick NS. Sonographic predictors of survival in fetal diaphragmatic hernia. J Pediatr Surg. 1996;31: 148–152. doi:10.1016/S0022-3468(96)90338-3

7. Jani J, Peralta CFA, Benachi A, Deprest J, Nicolaidis KH. Assessment of lung area in fetuses with congenital diaphragmatic hernia. Ultrasound Obstet Gynecol. 2007;30: 72–76. doi:10.1002/uog.4051

8. Mahieu-Caputo D, Sonigo P, Dommergues M, Fournet JC, Thalabard JC, Abarca C, et al. Fetal lung volume measurement by magnetic resonance imaging in congenital diaphragmatic hernia. BJOG An Int J Obstet Gynaecol. 2001;108: 863–868. doi:10.1111/j.1471-0528.2001.00184.x

9. Bultez T, Quibel T, Bouhanna P, Popowski T, Resche-Rigon M, Rozenberg P. Angle of fetal head progression measured using transperineal ultrasound as a predictive factor of vacuum extraction failure. Ultrasound Obstet Gynecol. 2016;48: 86–91. doi:10.1002/uog.14951

10. Büsing KA, Kilian AK, Schaible T, Dinter DJ, Neff KW. MR Lung Volume in Fetal Congenital Diaphragmatic Hernia: Logistic Regression Analysis—Mortality and Extracorporeal Membrane Oxygenation. Radiology. 2008;248: 233–239. doi:10.1148/radiol.2481070934

11. Ruano R, Lazar DA, Cass DL, Zamora JJ, Lee TC, Cassady CI, et al. Fetal lung volume and quantification of liver herniation by magnetic resonance imaging in isolated congenital diaphragmatic hernia. Ultrasound Obstet Gynecol. 2014;43: 662–669. doi:10.1002/uog.13223

12. Cannie M, Jani J, Meerschaert J, Allegaert K, Done' E, Marchal G, et al. Prenatal prediction of survival in isolated diaphragmatic hernia using observed to expected total fetal lung

volume determined by magnetic resonance imaging based on either gestational age or fetal body volume. *Ultrasound Obstet Gynecol.* 2008;32: 633–639. doi:10.1002/uog.6139

13. Lazar DA, Ruano R, Cass DL, Moise KJ, Johnson A, Lee TC, et al. Defining “liver-up”: does the volume of liver herniation predict outcome for fetuses with isolated left-sided congenital diaphragmatic hernia? *J Pediatr Surg.* 2012;47: 1058–1062. doi:10.1016/j.jpedsurg.2012.03.003

14. Cordier A -G., Jani JC, Cannie MM, Rodó C, Fabietti I, Persico N, et al. Stomach position in prediction of survival in left-sided congenital diaphragmatic hernia with or without fetoscopic endoluminal tracheal occlusion. *Ultrasound Obstet Gynecol.* 2015;46: 155–161. doi:10.1002/uog.14759

15. Estimating disease severity of congenital diaphragmatic hernia in the first 5 minutes of life. *J Pediatr Surg.* 2001;36: 141–145. doi:10.1053/jpsu.2001.20032

16. Brindle ME, Cook EF, Tibboel D, Lally PA, Lally KP. A Clinical Prediction Rule for the Severity of Congenital Diaphragmatic Hernias in Newborns. *Pediatrics.* 2014;134: e413–e419. doi:10.1542/peds.2013-3367

17. Schultz CM, DiGeronimo RJ, Yoder BA. Congenital diaphragmatic hernia: a simplified postnatal predictor of outcome. *J Pediatr Surg.* 2007;42: 510–516. doi:10.1016/j.jpedsurg.2006.10.043

18. Skarsgard ED, MacNab YC, Qiu Z, Little R, Lee SK. SNAP-II Predicts Mortality among Infants with Congenital Diaphragmatic Hernia. *J Perinatol.* 2005;25: 315–319. doi:10.1038/sj.jp.7211257

19. Jancelewicz T, Brindle ME. Prediction tools in congenital diaphragmatic hernia. *Semin Perinatol.* 2020;44: 151165. doi:10.1053/j.semperi.2019.07.004

20. Masino AJ, Harris MC, Forsyth D, Ostapenko S, Srinivasan L, Bonafide CP, et al. Machine learning models for early sepsis recognition in the neonatal intensive care unit using readily available electronic health record data. Juarez JM, editor. *PLoS One.* 2019;14: e0212665.

doi:10.1371/journal.pone.0212665

21. Hamilton EF, Dyachenko A, Ciampi A, Maurel K, Warrick PA, Garite TJ. Estimating risk of severe neonatal morbidity in preterm births under 32 weeks of gestation. *J Matern Neonatal Med.* 2020;33: 73–80. doi:10.1080/14767058.2018.1487395
22. Mani S, Ozdas A, Aliferis C, Varol HA, Chen Q, Carnevale R, et al. Medical decision support using machine learning for early detection of late-onset neonatal sepsis. *J Am Med Inform Assoc.* 2014;21: 326–36. doi:10.1136/amiajnl-2013-001854
23. Kumaresan D, Stephenson J, Doxey AC, Bandukwala H, Brooks E, Hillebrand-Voiculescu A, et al. Aerobic proteobacterial methylotrophs in Movile Cave: genomic and metagenomic analyses. *Microbiome.* 2018;6: 1. doi:10.1186/s40168-017-0383-2
24. Azevedo RSS, de Sousa JR, Araujo MTF, Martins Filho AJ, de Alcantara BN, Araujo FMC, et al. In situ immune response and mechanisms of cell damage in central nervous system of fatal cases microcephaly by Zika virus. *Sci Rep.* 2018;8: 1. doi:10.1038/s41598-017-17765-5
25. Cooper JN, Minneci PC, Deans KJ. Postoperative neonatal mortality prediction using superlearning. *J Surg Res.* 2018;221: 311–319. doi:10.1016/j.jss.2017.09.002
26. Irles C, González-Pérez G, Carrera Muiños S, Michel Macias C, Sánchez Gómez C, Martínez-Zepeda A, et al. Estimation of Neonatal Intestinal Perforation Associated with Necrotizing Enterocolitis by Machine Learning Reveals New Key Factors. *Int J Environ Res Public Health.* 2018;15. doi:10.3390/ijerph15112509
27. Masino AJ, Harris MC, Forsyth D, Ostapenko S, Srinivasan L, Bonafide CP, et al. Machine learning models for early sepsis recognition in the neonatal intensive care unit using readily available electronic health record data. *PLoS One.* 2019;14: e0212665. doi:10.1371/journal.pone.0212665
28. Hamilton EF, Dyachenko A, Ciampi A, Maurel K, Warrick PA, Garite TJ. Estimating risk of severe neonatal morbidity in preterm births under 32 weeks of gestation. *J Matern Fetal Neonatal Med.* 2020;33: 73–80. doi:10.1080/14767058.2018.1487395

29. Amodeo I, De Nunzio G, Raffaelli G, Borzani I, Griggio A, Conte L, et al. A maChine and deep Learning Approach to predict pulmoNary hypertenSion in newbornS with congenital diaphragmatic Hernia (CLANNISH): Protocol for a retrospective study. PLoS One. 2021;16: e0259724. doi:10.1371/journal.pone.0259724
30. Conte L, Amodeo I, De Nunzio G, Raffaelli G, Borzani I, Persico N, et al. Congenital diaphragmatic hernia: automatic lung and liver MRI segmentation with nnU-Net, reproducibility of pyradiomics features, and a machine learning application for the classification of liver herniation. Eur J Pediatr. 2024. doi:10.1007/s00431-024-05476-9
31. Rypens F, Metens T, Rocourt N, Sonigo P, Brunelle F, Quere MP, et al. Fetal Lung Volume: Estimation at MR Imaging—Initial Results. Radiology. 2001;219: 236–241. doi:10.1148/radiology.219.1.r01ap18236
32. Zizka J, Elias P, Hodik K, Tintera J, Juttnerova V, Belobradek Z, et al. Liver, meconium, haemorrhage: the value of T1-weighted images in fetal MRI. Pediatr Radiol. 2006;36: 792–801. doi:10.1007/s00247-006-0239-6
33. van Griethuysen JJM, Fedorov A, Parmar C, Hosny A, Aucoin N, Narayan V, et al. Computational Radiomics System to Decode the Radiographic Phenotype. Cancer Res. 2017;77: e104–e107. doi:10.1158/0008-5472.CAN-17-0339
34. Xiao Z, Gong K, Zou Y. A combined forecasting approach based on fuzzy soft sets. J Comput Appl Math. 2009;228: 326–333. doi:10.1016/j.cam.2008.09.033
35. Thanh Noi P, Kappas M. Comparison of Random Forest, k-Nearest Neighbor, and Support Vector Machine Classifiers for Land Cover Classification Using Sentinel-2 Imagery. Sensors. 2017;18: 18. doi:10.3390/s18010018
36. Doyle NM, Lally KP. The CDH study group and advances in the clinical care of the patient with congenital diaphragmatic hernia. Semin Perinatol. 2004;28: 174–184. doi:10.1053/j.semperi.2004.03.009
37. Long A-M, Bunch KJ, Knight M, Kurinczuk JJ, Losty PD. One-year outcomes of infants

- 510 born with congenital diaphragmatic hernia: a national population cohort study. Arch Dis
- 511 Child - Fetal Neonatal Ed. 2019;104: F643–F647. doi:10.1136/archdischild-2018-316396
- 512 38. Cochius-den Otter SCM, Erdem Ö, van Rosmalen J, Schaible T, Peters NCJ, Cohen-
- 513 Overbeek TE, et al. Validation of a Prediction Rule for Mortality in Congenital
- 514 Diaphragmatic Hernia. Pediatrics. 2020;145. doi:10.1542/peds.2019-2379
- 515 39. Kipfmüller F, Schroeder L, Melaku T, Geipel A, Berg C, Gembruch U, et al. Prediction of
- 516 ECMO and Mortality in Neonates with Congenital Diaphragmatic Hernia Using the SNAP-II
- 517 Score. Klin Pädiatrie. 2019;231: 297–303. doi:10.1055/a-1009-6671
- 518 40. Montalva L, Raffler G, Riccio A, Lauriti G, Zani A. Neurodevelopmental impairment in
- 519 children with congenital diaphragmatic hernia: Not an uncommon complication for
- 520 survivors. J Pediatr Surg. 2020;55: 625–634. doi:10.1016/j.jpedsurg.2019.05.021
- 521 41. Kinsella JP, Steinhorn RH, Mullen MP, Hopper RK, Keller RL, Ivy DD, et al. The Left
- 522 Ventricle in Congenital Diaphragmatic Hernia: Implications for the Management of
- 523 Pulmonary Hypertension. J Pediatr. 2018;197: 17–22. doi:10.1016/j.jpeds.2018.02.040
- 524 42. Massolo AC, Paria A, Hunter L, Finlay E, Davis CF, Patel N. Ventricular Dysfunction,
- 525 Interdependence, and Mechanical Dispersion in Newborn Infants with Congenital
- 526 Diaphragmatic Hernia. Neonatology. 2019;116: 68–75. doi:10.1159/000499347
- 527 43. Gien J, Kinsella JP. Management of pulmonary hypertension in infants with congenital
- 528 diaphragmatic hernia. J Perinatol. 2016;36: S28–S31. doi:10.1038/jp.2016.46
- 529 44. Mank A, Carrasco Carrasco C, Thio M, Clotet J, Pauws SC, DeKoninck P, et al. Tidal
- 530 volumes at birth as predictor for adverse outcome in congenital diaphragmatic hernia. Arch
- 531 Dis Child - Fetal Neonatal Ed. 2020;105: 248–252. doi:10.1136/archdischild-2018-316504
- 532 45. Oh C, Youn JK, Han J, Yang H, Lee S, Seo J, et al. Predicting Survival of Congenital
- 533 Diaphragmatic Hernia on the First Day of Life. World J Surg. 2019;43: 282–290.
- 534 doi:10.1007/s00268-018-4780-x
- 535 46. Brown BP, Clark MT, Wise RL, Timsina LR, Reher TA, Vandewalle RJ, et al. A

multifactorial severity score for left congenital diaphragmatic hernia in a high-risk population using fetal magnetic resonance imaging. *Pediatr Radiol.* 2019;49: 1718–1725. doi:10.1007/s00247-019-04478-2

47. Dassios T, Ali K, Makin E, Bhat R, Krokidis M, Greenough A. Prediction of Mortality in Newborn Infants With Severe Congenital Diaphragmatic Hernia Using the Chest Radiographic Thoracic Area*. *Pediatr Crit Care Med.* 2019;20: 534–539. doi:10.1097/PCC.0000000000001912

48. Jani J, Nicolaides KH, Keller RL, Benachi A, Peralta CFA, Favre R, et al. Observed to expected lung area to head circumference ratio in the prediction of survival in fetuses with isolated diaphragmatic hernia. *Ultrasound Obstet Gynecol.* 2007;30: 67–71. doi:10.1002/uog.4052

49. Snoek KG, Peters NCJ, van Rosmalen J, van Heijst AFJ, Eggink AJ, Sikkink E, et al. The validity of the observed-to-expected lung-to-head ratio in congenital diaphragmatic hernia in an era of standardized neonatal treatment; a multicenter study. *Prenat Diagn.* 2017;37: 658–665. doi:10.1002/pd.5062

50. Amodeo I, Borzani I, Raffaeli G, Persico N, Amelio GS, Gulden S, et al. The role of magnetic resonance imaging in the diagnosis and prognostic evaluation of fetuses with congenital diaphragmatic hernia. *Eur J Pediatr.* 2022;181: 3243–3257. doi:10.1007/s00431-022-04540-6

51. Victoria T, Bebbington MW, Danzer E, Flake AW, Johnson MP, Dinan D, et al. Use of magnetic resonance imaging in prenatal prognosis of the fetus with isolated left congenital diaphragmatic hernia. *Prenat Diagn.* 2012;32: 715–23. doi:10.1002/pd.3890

52. Dütemeyer V, Cordier A-G, Cannie MM, Bevilacqua E, Huynh V, Houfflin-Debarge V, et al. Prenatal prediction of postnatal survival in fetuses with congenital diaphragmatic hernia using MRI: lung volume measurement, signal intensity ratio, and effect of experience. *J Matern Fetal Neonatal Med.* 2022;35: 1036–1044. doi:10.1080/14767058.2020.1740982

53. Jani J, Cannie M, Done E, Van Mieghem T, Van Schoubroeck D, Gucciardo L, et al. Relationship between lung area at ultrasound examination and lung volume assessment with magnetic resonance imaging in isolated congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol.* 2007;30: 855–60. doi:10.1002/uog.5168
54. Jani J, Cannie M, Sonigo P, Robert Y, Moreno O, Benachi A, et al. Value of prenatal magnetic resonance imaging in the prediction of postnatal outcome in fetuses with diaphragmatic hernia. *Ultrasound Obstet Gynecol.* 2008;32: 793–9. doi:10.1002/uog.6234
55. Bebbington M, Victoria T, Danzer E, Moldenhauer J, Khalek N, Johnson M, et al. Comparison of ultrasound and magnetic resonance imaging parameters in predicting survival in isolated left-sided congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol.* 2014;43: 670–4. doi:10.1002/uog.13271
56. Oluyomi-Obi T, Kuret V, Puligandla P, Lodha A, Lee-Robertson H, Lee K, et al. Antenatal predictors of outcome in prenatally diagnosed congenital diaphragmatic hernia (CDH). *J Pediatr Surg.* 2017;52: 881–888. doi:10.1016/j.jpedsurg.2016.12.008
57. Petroze RT, Caminsky NG, Trebichavsky J, Bouchard S, Le-Nguyen A, Laberge J-M, et al. Prenatal prediction of survival in congenital diaphragmatic hernia: An audit of postnatal outcomes. *J Pediatr Surg.* 2019;54: 925–931. doi:10.1016/j.jpedsurg.2019.01.021
58. Neff KW, Kilian AK, Schaible T, Schütz E-M, Büsing KA. Prediction of mortality and need for neonatal extracorporeal membrane oxygenation in fetuses with congenital diaphragmatic hernia: logistic regression analysis based on MRI fetal lung volume measurements. *AJR Am J Roentgenol.* 2007;189: 1307–11. doi:10.2214/AJR.07.2434
59. Lee TC, Lim FY, Keswani SG, Frischer JS, Haberman B, Kingma PS, et al. Late gestation fetal magnetic resonance imaging-derived total lung volume predicts postnatal survival and need for extracorporeal membrane oxygenation support in isolated congenital diaphragmatic hernia. *J Pediatr Surg.* 2011;46: 1165–71. doi:10.1016/j.jpedsurg.2011.03.046
60. Alfaraj MA, Shah PS, Bohn D, Pantazi S, O’Brien K, Chiu PP, et al. Congenital

diaphragmatic hernia: lung-to-head ratio and lung volume for prediction of outcome. Am J Obstet Gynecol. 2011;205: 43.e1–8. doi:10.1016/j.ajog.2011.02.050


61. Russo FM, Eastwood MP, Keijzer R, Al-Maary J, Toelen J, Van Mieghem T, et al. Lung size and liver herniation predict need for extracorporeal membrane oxygenation but not pulmonary hypertension in isolated congenital diaphragmatic hernia: systematic review and meta-analysis. Ultrasound Obstet Gynecol. 2017;49: 704–713. doi:10.1002/uog.16000

62. Walleyo A, Debus A, Kehl S, Weiss C, Schönberg SO, Schaible T, et al. Periodic MRI lung volume assessment in fetuses with congenital diaphragmatic hernia: prediction of survival, need for ECMO, and development of chronic lung disease. AJR Am J Roentgenol. 2013;201: 419–26. doi:10.2214/AJR.12.8655


63. Büsing KA, Kilian AK, Schaible T, Endler C, Schaffelder R, Neff KW. MR relative fetal lung volume in congenital diaphragmatic hernia: survival and need for extracorporeal membrane oxygenation. Radiology. 2008;248: 240–6. doi:10.1148/radiol.2481070952

64. Schaible T, Büsing KA, Felix JF, Hop WCJ, Zahn K, Wessel L, et al. Prediction of chronic lung disease, survival and need for ECMO therapy in infants with congenital diaphragmatic hernia: additional value of fetal MRI measurements? Eur J Radiol. 2012;81: 1076–82. doi:10.1016/j.ejrad.2011.02.060


Receiver Operating Characteristic (ROC) Curve



Precision-Recall Curve



Receiver Operating Characteristic (ROC) Curve



Precision-Recall Curve

