

Title

Health Utility Value of Overactive Bladder in Japanese Older Adults

Authors and affiliations

Takashi Yoshioka, MD, PhD^{1,2*}

Kenji Omae, MD, DrPH^{3,4}

Satoshi Funada, MD, PhD¹

Tetsuji Minami, MD, PhD⁵

Rei Goto, MD, PhD^{6,7}

¹ Health Technology Assessment Unit, Department of Preventive Medicine and Public Health, Keio University School of Medicine, Tokyo, Japan

² Institute of Clinical Epidemiology, Showa University, Tokyo, Japan

³ Department of Innovative Research and Education for Clinicians and Trainees (DiRECT), Fukushima Medical University Hospital, Fukushima, Japan

⁴ Department of Health Promotion and Human Behavior, Graduate School of Medicine/School of Public Health, Kyoto University, Kyoto, Japan

⁵ Division of Quality Assurance Programs, Institute for Cancer Control, National Cancer Center, Tokyo, Japan

⁶ Graduate School of Health Management, Keio University, Kanagawa, Japan

⁷ Graduate School of Business Administration, Keio University, Kanagawa, Japan

*** Corresponding author**

Takashi Yoshioka

Health Technology Assessment Unit, Department of Preventive Medicine and Public Health, Keio University School of Medicine, Tokyo, Japan

35 Shinano-machi, Shinjuku-ku, Tokyo, 160-8582, Japan

Tel: +81-3-5363-3205

E-mail: ty5733@keio.jp

30

31 **E-mail address:**

32 Takashi Yoshioka, ty5733@keio.jp

33 Kenji Omae, oranz416@gmail.com

34 Satoshi Funada, sfunada@keio.jp

35 Tetsuji Minami, t.minami.for.urology@gmail.com

36 Rei Goto, reigoto@kbs.keio.ac.jp

37

38 **Word counts:** 3,324

39 **Number of Tables:** 3

40 **Number of Figures:** 3

41 **Number of References:** 30

42

43 Abstract

44 **Objectives:** To determine the health utility values (HUVs) of overactive bladder (OAB)
45 among adults aged ≥ 65 years and to assess the HUV decrements (disutilities) of OAB
46 according to its severity.

47 **Methods:** This cross-sectional Internet-based study was conducted between 2 and 9
48 November 2023, with quota sampling with equal probability for each gender and age group
49 (age 65–74 years and ≥ 75 years). OAB was defined as an urgency score of ≥ 2 points and a
50 total score of ≥ 3 points based on the Overactive Bladder Symptom Score. OAB severity was
51 categorized as mild (total score, ≤ 5 points) or moderate-to-severe (total score, 6–15 points).
52 HUVs were measured using the EuroQol 5-Dimension 5-Level value set for the Japanese
53 population. Multivariable linear regression models were fitted to estimate the
54 covariate-adjusted disutilities of OAB, with eight covariates selected based on previous
55 studies.

56 **Results:** Among the 998 participants (51.9% male; mean age, 73.2 years), 158 (15.9%) had
57 OAB, of whom 87 (8.8%) had moderate-to-severe OAB. The mean HUVs for participants
58 with mild and moderate-to-severe OAB were 0.874 and 0.840, respectively, which were
59 lower compared with the HUV for those without OAB (0.913). After adjusting for relevant
60 covariates, disutilities (95% confidence intervals [CIs]) for mild and moderate-to-severe
61 OAB were -0.0334 (-0.0602 to -0.0066) and -0.0591 (-0.0844 to -0.0339), respectively.

62 **Conclusions:** This study examined HUVs in older Japanese adults with and without OAB.
63 The results demonstrate that increased OAB severity is associated with greater disutility.

64
65

66 **Keywords:**

67 Overactive bladder; lower urinary tract symptoms; health utility; health-related quality of life;

68 health economic evaluation

69

70

71 Introduction

72 Overactive bladder (OAB), characterized by urinary urgency with or without urinary
 73 incontinence, is associated with a substantial disease burden and significant economic impact
 74 [1]. The extensive disease burden of OAB includes a decrease in health-related quality of life
 75 (HRQoL), mood disorders, and various comorbid or consequential conditions (*e.g.*
 76 constipation, fecal incontinence, and irritable bowel syndrome) [2]. The prevalence of OAB
 77 increases with age [3]. Furthermore, evidence has demonstrated a longitudinal association
 78 between OAB or its treatment with anticholinergics and an increased risk of falls, which are
 79 the most critical outcomes in older adults [4]. Consequently, patients with OAB have higher
 80 healthcare resource utilization than those without the condition, which contributes to a
 81 substantial economic burden [5]. The high global prevalence of OAB, estimated at 15.5%
 82 between 2007 and 2020, and its high healthcare cost of \$82.6 billion by 2020 in the United
 83 States [1, 6] have drawn significant attention from both policymakers and clinicians
 84 worldwide.

85 OAB is widely managed in various healthcare settings, including primary care,
 86 urology, and urogynecology [7]. Treatment options for OAB include behavioral therapy and
 87 pharmacotherapy, such as antimuscarinic and β -adrenergic agents [7]. Surgical treatments are
 88 available for refractory cases, including intradetrusor injection of onabotulinum toxin A,
 89 sacral neuromodulation, and percutaneous tibial nerve stimulation [7]. Although these
 90 therapeutic options have the potential to relieve patients' symptoms and improve their
 91 HRQoL, they also raise concerns about the increasing economic burden on healthcare
 92 systems owing to higher treatment costs. Consequently, the importance of effectively
 93 allocating finite healthcare resources for the management of OAB is becoming increasingly
 94 recognized. In fact, the National Institute for Health and Care Excellence (NICE), an
 95 independent organization providing evidence-based guidance to inform healthcare decisions
 96 in the United Kingdom (UK), is currently conducting a health economic evaluation (HEE) of
 97 vibegron [8], a novel β 3-adrenergic agonist with fewer antimuscarinic side effects compared
 98 with traditional OAB medications [9]. Considering the example of the NICE in the UK, new

medical technologies for OAB, such as vibegrons, may also be subject to HEEs in other countries, such as Japan.

To make HEEs useful for policy decision-making, it is essential to assess their clinical effectiveness (*e.g.* quality-adjusted life years [QALYs]) using generic measures that can be collected for any disease. QALYs are calculated based on health utility values (HUVs) measured using generic preference-based measures (PBMs) rather than disease-specific ones [10]. Importantly, according to the NICE technology appraisal guidance, HUVs should be calculated using the EuroQoL 5-dimensions (EQ-5D), with data collected directly from patients in the country where the evaluation is being conducted (in the case of NICE, the UK) [11]. Previous studies have evaluated the association between OAB symptoms and HUVs using EQ-5D scores in the United States, United Kingdom, Spain, and South Korea [12–14]. However, these studies have utilized the older version of the EQ-5D, the three-level EQ-5D version (EQ-5D-3L), which raises concerns about the potential overestimation of positive effects and underestimation of negative effects [15]. The newer version of the EQ-5D, the five-level EQ-5D version (EQ-5D-5L), was developed to overcome the ceiling effects and low sensitivity inherent in EQ-5D-3L [16]. Ideally, the estimation of HUVs based on EQ-5D-5L is essential. However, few studies have used this instrument.

To address the unmet needs of global policies regarding HEEs of OAB, this study aimed to describe the HUVs of patients with OAB and assess the impacts of OAB on HUVs using EQ-5D-5L among older adults, the most vulnerable and policy-relevant subgroup, using an Internet-based survey.

Participants and methods

Study design and participants

This cross-sectional study was conducted as part of a research project aimed at developing an EQ-5D-5L value set for the Japanese general older adults [16]. To obtain responses from a sample of older adults, we used a sampling method that ensured a 1:1 ratio for both sex and age groups (age 65–74 and ≥ 75 years). The target sample size was 1,000 participants. The sampling strategy and sample size were determined based on a previous study [17]. All

participants were recruited from a survey panel provided by a Japanese Internet research agency (Intage, Inc., <https://www.intage.co.jp/english/>) with approximately 2.6 million panelists and their demographic information. The survey was conducted between 2 and 9 November 2023. We designed the survey to prevent participants from proceeding to the next question without providing a valid response, thereby eliminating missing data. The survey questionnaire and response options are presented in **Method S1**.

Inclusion and exclusion criteria

All respondents were included in the analysis. However, participants who responded ‘prefer not to answer’ to at least one of the questionnaires except for income items were excluded from this study.

Exposures (overactive bladder)

The variable of interest in this study was the presence of OAB. OAB was assessed using the Overactive Bladder Symptom Score (OABSS), a validated questionnaire developed in Japanese [18]. The OABSS is a self-reported questionnaire consisting of four domains (daytime frequency, 0–2 points; nighttime frequency, 0–3 points; urgency, 0–5 points; urgency incontinence, 0–5 points), with a total score of 15 points [18]. A higher score indicates more severe OAB symptoms. Based on previous studies and decision modelling on HEEs, OAB was defined as an urgency score of ≥ 2 points and a total score of ≥ 3 points, and the severity of OAB was categorized as mild (OAB with a total score of ≤ 5 points) or moderate-to-severe (OAB with a total score of 6–15 points) [14].

Outcomes (health utility values)

Our primary outcome of interest was HUVs measured using the EQ-5D-5L instrument. HUVs are an indicator of an individual’s current health state, where 0 represents death or the worst imaginable health state and 1 represents perfect health, with higher scores indicating better health status [10]. In HEEs, HUVs are calculated using PBMs developed with choice methods (e.g. standard gambling or time trade-off) as reference standards [10, 11]. The

EQ-5D-5L is a PBM developed by the EuroQoL Group that is used to measure HUVs based on time trade-off methods [10, 11]. Importantly, the EQ-5D-5L has been translated and validated in Japanese, demonstrating substantial reliability and validity for calculating HUVs in the general Japanese population [17].

In this study, we used the EuroQoL Visual Analogue Scale (EQ VAS) score as another measure of health utility. The EQ VAS is a scale developed by the EuroQoL Group that measures the current health state using a VAS on a vertically oriented ruler with markings from 0 to 100 [17]. On the EQ-VAS, 0 represents death or the worst imaginable health state, and 100 indicates perfect health. The EQ VAS has often been used in conjunction with the EQ-5D in similar studies [14], and consistent results have been reported. Therefore, we used the EQ-VAS score as an outcome measure for sensitivity analysis in this study.

This study used the online formats of the EQ-5D-5L and EQ VAS. The research protocol was pre-registered on the EuroQoL website [16], and permission was obtained to use these instruments. The online versions were administered in strict accordance with the manufacturers' instructions. The EuroQoL registration number for this study is 59331.

Covariates

The covariates of interest in this study were demographic, socioeconomic, and health-related variables. These variables were selected from previous studies on OAB that focused on the factors that could potentially influence HUVs. The demographic factors included age (continuous), gender (male or female), and body mass index BMI (continuous). Socioeconomic status included educational attainment (high school education or less and college education or more) and income [6]. As a proxy for income, this study used equivalent household income, derived by adjusting total household income by the square root of household size (then grouped by median, with ranges provided: lower half, JPY 0–2.8 million; upper half, JPY ≥ 2.9 million), along with a declined to answer category. Health-related factors included smoking status (never/past/current), alcohol consumption (never/past/current), and self-reported coexisting physical and psychiatric comorbidities.

Comorbidity data were used to record the presence or absence of the following diseases: hypertension, dyslipidemia, diabetes, low back pain, ischemic heart disease, stroke, chronic kidney disease, stress urinary incontinence, depression, and other psychiatric disorders [5, 19–22].

Statistical analyses

This study used three analytical approaches. First, a primary analysis was performed targeting all respondents. Second, subgroup analyses were performed, stratified based on sex (male/female) and age (65–74 years old/ ≥ 75 years), assuming that the impacts of OAB on HUVs may vary according to these factors. Third, an exploratory analysis was performed to evaluate the impact of each of the four domains constituting the OAB on HUVs. In all analyses, we initially described the baseline characteristics, including the prevalence of OAB and HUVs. Subsequently, we described the mean (standard deviation [SD]) and median (interquartile range [IQR]) of HUVs based on summary statistics for each group for all analyses. Finally, we estimated the adjusted HUV decrements (disutilities) using a multivariable linear regression analysis after adjusting for covariates. In the multivariable linear regression analysis with ordinal categories (non-OAB/mild OAB/moderate-to-severe OAB), a linear trend test was conducted to assess linearity. To confirm the consistency of the results, a sensitivity analysis was performed in the primary analysis to fit a similar linear regression model using the EQ VAS score as the outcome measure.

For all regression models, β coefficients, 95% confidence intervals (CIs), and p -values were calculated. Statistical significance was set at $p < 0.05$. All analyses were performed using STATA version 18.0 (Stata Corp, College Station, Texas, USA).

Ethical considerations

All respondents completed online questionnaires after providing digital informed consent indicating their willingness to participate in the survey. Participants received monetary points as an incentive for participation. All the procedures adhered to the ethical guidelines of the 1975 Declaration of Helsinki and its subsequent 2013 revision. This study was approved by

the Institutional Review Board of Keio University School of Medicine (approval number: 20221120). This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for cross-sectional studies.

Results

Baseline characteristics

Of the 1,099 respondents, 994 (90.7%) were included in the analysis (**Figure 1**). The median age of the participants was 73.2 (SD, 5.3) years, and 516 (51.9%) were male. Regarding socioeconomic indicators, 557 (56.0%) participants had college or higher education. Regarding health-related factors, 107 (10.8%) participants were current tobacco users, 519 (52.2%) were current alcohol users, 486 (48.9%) had hypertension, 166 (16.7%) had diabetes, 114 (11.5%) had stress urinary incontinence, and 50 (5.0%) had depression. The mean OABSS was 2.7 (SD, 2.2). The mean HUV was 0.90 (SD, 0.11), and the mean EQ VAS score was 75.2 (SD, 17.6) (**Table 1**).

Health utility values based on overactive bladder status

Table 2 presents a summary of the HUVs stratified according to the presence and severity of OAB. The mean HUV for participants without OAB was 0.913 (SD, 0.105), whereas that for those with OAB was 0.855 (SD, 0.145). When categorizing OAB severity into mild and moderate-to-severe OAB, the mean HUV for participants with mild OAB was 0.874 (SD, 0.131), and for those with moderate-to-severe OAB, it was 0.840 (SD, 0.155).

Disutilities of overall respondents

Table 3 presents the results of the multivariable linear regression analyses. After adjusting for covariates, the estimated disutility for the presence of OAB was −0.0472 (95% CI, −0.0664 to −0.0279; $p < 0.001$). The estimated disutilities for mild and moderate-to-severe OAB were −0.0334 (95% CI, −0.0602 to −0.0066; $p = 0.014$) and −0.0591 (95% CI, −0.0844 to −0.0339; $p < 0.001$), respectively, indicating a linear relationship (p for trend < 0.001). The sensitivity analysis using the EQ VAS as the outcome variable showed similar results

(presence of OAB, $\beta=-5.42$, 95% CI -8.47 to -2.37 , $p=0.001$; mild OAB, $\beta=-4.36$, 95% CI -8.60 to -0.12 , $p=0.044$; moderate-to-severe OAB, $\beta=-6.26$, 95% CI -10.26 to -2.27 , $p=0.002$; p for trend <0.001).

Results of the subgroup analysis

Tables S1 and **S2** present the baseline characteristics of the sex and age subgroups, respectively. **Table S3** summarizes the HUVs for sex and age subgroups of HUVs. In the sex subgroup, regardless of the presence of OAB, females ($n=478$) had lower mean HUVs than males ($n=516$) (*e.g.* presence of any OAB: male, 0.872 [SD, 0.128]; female, 0.821 [SD, 0.173]). In the age subgroup, regardless of the presence of OAB, the 75–94-year-old group ($n=502$) showed lower HUVs compared with the 65–74-year-old group ($n=492$) (*e.g.* presence of any OAB: 65–74 years old, 0.889 [SD, 0.130]; 75–94 years old, 0.829 [SD, 0.152]).

The results of the multivariable linear regression analysis are shown in **Figure 2**. In the gender subgroup, a substantial disutility was observed in females with moderate-to-severe OAB ($\beta=-0.1026$; 95% CI, -0.1510 to -0.0542 ; $p<0.001$). In the age subgroup, substantial disutilities were observed in the 75–94-year-old group with mild OAB ($\beta=-0.0688$; 95% CI, -0.1106 to -0.0271 ; $p=0.001$) and moderate-to-severe OAB ($\beta=-0.0664$; 95% CI, -0.1006 to -0.0322 ; $p<0.001$).

Results of the exploratory analysis

Table S4 presents a summary of the HUVs for each domain and the score of the OABSS. The higher OABSS scores indicated the lower mean HUVs in the three domains, excluding urgency urinary incontinence. The results of the multivariable linear regression analysis after rounding scores of $<2.5\%$ of the total are shown in **Figure 3**. After adjusting for covariates, disutilities were observed for each score in the three domains, excluding daytime frequency (*e.g.* nighttime frequency: 1 point, $\beta=-0.0089$, 95% CI -0.0276 to 0.0099 , $p=0.035$; 2 points, $\beta=-0.0193$, 95% CI -0.0411 to 0.0025 , $p=0.082$; 3 points, $\beta=-0.0409$, 95% CI -0.0682 to -0.0136 , $p=0.003$). A linear relationship between scores and disutilities was observed in all

domains, except for daytime frequency (p for trend: nighttime frequency, 0.003; urgency, <0.001; and urgency incontinence, <0.001).

Discussion

This study described the HUVs of Japanese adults aged ≥ 65 years with or without OAB and estimated the associated disutility of OAB using regression models. The results demonstrated that the mean HUVs of individuals with OAB were lower and the disutility of OAB was greater, depending on the severity of the condition. Subgroup analyses suggested that females and those aged 75 years might experience greater disutility owing to OAB than their counterparts. Furthermore, exploratory analyses indicated the possibility of score-proportional disutility in all domains of the OABSS, except daytime frequency.

Previous studies have evaluated the associations of OAB with HUVs in several countries. Kay *et al.* conducted a study estimating EQ-5D-3L-based HUVs from the Incontinence-Specific Quality of Life Questionnaire in 2,505 patients with idiopathic or neurogenic OAB in the United States and Europe [12]. Similarly, Ruiz *et al.* used the Overactive Bladder Symptom and Health-Related Quality of Life Questionnaire in 246 patients with OAB to estimate the effects of OAB symptoms on EQ-5D-3L-based HUVs in Spain [13]. Additionally, Kim *et al.* assessed the effects of OAB severity as defined by the OABSS on EQ-5D-3L-based HUVs in 226,867 participants (including 12,303 patients with OAB) from the 2012 Korean Community Health Survey [14]. These studies shared two common features. First, they estimated the associations of OAB with HUVs using OAB measures that are policy-relevant in their respective countries. Second, they employed the EQ-5D-3L for evaluation. These factors significantly influenced the design of this study in a Japanese setting. Second, regarding the choice of OAB measure, we used the OABSS, similar to Kim *et al.*, as it is explicitly used to define OAB in the Japanese OAB guidelines [23]. The OABSS was used to define OAB in a randomized controlled trial evaluating the effects of OAB interventions in Japan [24]. Therefore, decision modelling based on the OABSS is expected when conducting HEEs targeting patients with OAB in Japan. Second, we used the

EQ-5D-5L, for which a value set developed and validated by Ikeda *et al.* exists in Japan [17]. Evidence suggests that the EQ-5D-3L may potentially overestimate positive associations and underestimate negative associations compared with the EQ-5D-5L in HEEs [15]. Consequently, the EQ-5D-5L is recommended for cost-effectiveness evaluations in Japan [25], and our study adhered to this recommendation. This study, which fully considers Japanese clinical practices and healthcare systems, provides HUVs appropriate for HEEs in the Japanese setting. We believe that our approach offers valuable insights for the design of future studies in any country.

In this study, the disutility values for mild and moderate-to-severe OABs as defined by the OABSS were -0.034 and -0.0591 , respectively. When referring to the Japanese population norms using the EQ-5D-5L, these disutilities are comparable to those of other chronic conditions that affect HRQoL, such as allergic rhinitis (-0.027), atopic dermatitis (-0.031), and other skin diseases (-0.043) [26]. In a report from Korea, which used the same scale (OABSS) and criteria to define OAB as our study, the covariate-adjusted disutility values for mild OAB, moderate OAB, and severe OAB were -0.029 , -0.089 , and -0.179 , respectively [14]. The smaller disutility for moderate-to-severe OAB in our study compared to Kim *et al.*'s findings may be explained by differences in the mean age (75 years vs. 45.2 years), outcome measure (EQ-5D-5L vs. EQ-5D-3L), and ethnicity (Japanese vs. Korean). However, both studies consistently showed that disutility increased with severity, emphasizing the importance of the associations of OAB with HUVs. Furthermore, our study found high disutility (-0.1026) among female participants with moderate-to-severe OAB in the subgroup analyses. This finding may be supported by the results of EpiLUTS, an Internet survey conducted among individuals aged >40 years in the United States, which showed that a higher proportion of women (39.4%) than men (28.7%) reported moderate-to-severe problems in their patient perception of bladder condition for OAB with bother [27]. Additionally, in the subgroup of individuals aged ≥ 75 years, mild and moderate-to-severe OABs were associated with disutility values of similar magnitude (mild OAB -0.0688 vs. moderate-to-severe OAB -0.0664). This finding suggests that the presence of OAB, regardless of severity, may have a substantial impact on older adults who are more likely to

be frail than younger respondents. For example, evidence has shown that even in the absence of incontinence, the presence of OAB is a significant risk factor for falls [28].

Our study has several strengths. First, we comprehensively provided the descriptive and analytical information necessary for the HEE of OAB, specifically for older adults. The mean and median HUVs at baseline and the disutilities after covariate adjustment for each severity level of OAB are essential information when conducting HEEs. Second, this study estimated disutility by adjusting for a wide range of evidence-based covariates, from socioeconomic to health-related indicators. Third, a subgroup analysis was performed to identify vulnerable populations that are essential for HEEs from the perspective of equity considerations but are understudied [29]. Moreover, our subgroup analysis results provide HUVs that can be referenced in future HEEs, particularly when treatments specifically targeting nocturia or urgent urinary incontinence among the OAB symptoms become available.

However, this study has also some limitations. First, as this study utilized an Internet survey, there were concerns regarding the external validity of the results. However, the prevalence of OAB (15.9%) and mean HUV (0.89) in this study did not substantially differ from previous epidemiological data from Japan [26, 30]. Thus, these values are also expected to apply to HEEs. Second, the sample size was limited (approximately 1,000); consequently, there were few individuals with high scores on the OABSS (*e.g.* 2 points for daytime frequency and 4–5 points for urgency urinary incontinence). As a result, it was not possible to differentiate between moderate and severe OAB, and some scores were grouped in the exploratory analysis. Although the number of individuals with such high scores may not be large and may not be considered in actual HEEs, revalidation of our study with a larger sample size would be valuable. Third, there may have been residual confounding factors. For example, orthopedic factors, such as falls and fractures, can be considered in future models.

In conclusion, this study examined HUVs in older Japanese adults with and without OAB. The results also demonstrated that increasing the severity of OAB indicated greater disutility, highlighting the substantial effect of OAB on HRQoL. These findings serve as

359 valuable resources for future HEEs of OAB, possibly guiding decision-making in healthcare

360 resource allocation for policymakers.

361

Acknowledgments

This study was supported by the Japan Society for the Promotion of Science KAKENHI grant (grant number: 21K17228) for conducting the survey. This study was also supported by the National Institute of Public Health for the language editing fee and article publishing charge.

Conflicts of interest

TY has received the Japan Society for the Promotion of Science (JSPS) KAKENHI grant (grant number: 21K17228) for conducting the survey. RG has received grants for the evaluation of the cost-effectiveness of medicines and medical devices from the National Institute of Public Health, Japan since 2019, which support the consultation fees for the development of search strategies and will support the article publication fee. KO, SF, and TM have no conflicts of interest to declare. The funders did not participate in the study design, data collection, analysis, interpretation, manuscript preparation, review, approval, or decision to submit the manuscript for publication.

Data availability statement

The datasets generated and analyzed during the current study are available from the corresponding author upon reasonable request.

References

1. Milsom I, Coyne KS, Nicholson S, Kvasz M, Chen CI, Wein AJ. Global prevalence and economic burden of urgency urinary incontinence: a systematic review. *Eur Urol* 2014; 65: 79–95.
2. Peyronnet B, Mironska E, Chapple C, *et al.* A comprehensive review of overactive bladder pathophysiology: on the way to tailored treatment. *Eur Urol* 2019; 75: 988–1000.
3. Eapen RS, Radomski SB. Review of the epidemiology of overactive bladder. *Res Rep Urol* 2016; 8: 71–6.
4. Just KS, Schultze KA, Dormann H, *et al.* Use of overactive bladder anticholinergic medications associated with falls leading to emergency department visits: results from the ADRED study. *Eur J Clin Pharmacol* 2023; 79: 1185–93.
5. Durden E, Walker D, Gray S, Fowler R, Juneau P, Gooch K. The economic burden of overactive bladder (OAB) and its effects on the costs associated with other chronic, age-related comorbidities in the United States. *Neurourol Urodyn* 2018; 37: 1641–9.
6. Lin W, Li T, Xu Z, *et al.* Association of socioeconomic status and overactive bladder in US adults: a cross-sectional analysis of nationally representative data. *Front Public Health* 2024; 12: 1345866.
7. Raju R, Linder BJ. Evaluation and treatment of overactive bladder in women. *Mayo Clin Proc* 2020; 95: 370–7.
8. Project information | Vibegron for treating symptoms of overactive bladder ID6300 | Guidance | NICE. [cited 2024 Jun 5]. Available from: <https://www.nice.org.uk/guidance/indevelopment/gid-ta11348>.
9. Keam SJ. Vibegron: first global approval. *Drugs* 2018; 78: 1835–9.
10. Chang EM, Saigal CS, Raldow AC. Explaining health state utility assessment. *JAMA*

- 2020; 323: 1085–6.
11. Foreword | Guide to the methods of technology appraisal, 2013. *Guidance* | NICE. [cited 2024 Jun 5]. Available from: <https://www.nice.org.uk/process/pmg9/chapter/foreword>.
12. Kay S, Tolley K, Colayco D, Khalaf K, Anderson P, Globe D. Mapping EQ-5D utility scores from the Incontinence Quality of Life Questionnaire among patients with neurogenic and idiopathic overactive bladder. *Value Health* 2013; 16: 394–402.
13. Ruiz MA, Gutiérrez LL, Monroy M, Rejas J. Mapping of the OAB-SF questionnaire onto EQ-5D in Spanish patients with overactive bladder. *Clin Drug Investig* 2016; 36: 267–79.
14. Kim SK, Kim SH. The impact of overactive bladder on health-related quality of life in Korea: based on the results of a Korean Community Health Survey. *Qual Life Res* 2021; 30: 1017–24.
15. Wailoo A, Alava MH, Pudney S, *et al.* An international comparison of EQ-5D-5L and EQ-5D-3L for use in cost-effectiveness analysis. *Value Health* 2021; 24: 568–74.
16. EuroQol [Internet], 2022 [cited 2024 Jun 10]. Available from: <https://euroqol.org/information-and-support/euroqol-instruments/eq-5d-5l/>. Eq-5d-5l.
17. Ikeda S, Shiroywa T, Igarashi A, *et al.* Developing a Japanese version of the EQ-5D-5L value set. *J Natl Inst Public Health* 2015; 64: 47–55.
18. Homma Y, Yoshida M, Seki N, *et al.* Symptom assessment tool for overactive bladder syndrome--overactive bladder symptom score. *Urology* 2006; 68: 318–23.
19. Welk B, Baverstock R. Is there a link between back pain and urinary symptoms? *Neurourol Urodyn* 2020; 39: 523–32.
20. Ceyhun G, Erbay G. Relationship between the severity of coronary artery disease and overactive bladder. *Acta Cardiol Sin* 2021; 37: 254–60.

- 429 21. Lai S, Pastore S, Piloni L, *et al.* Chronic kidney disease and urological disorders:
430 systematic use of uroflowmetry in nephropathic patients. *Clin Kidney J* 2019; 12: 414–9.
- 431 22. Leron E, Weintraub AY, Mastrolia SA, Schwarzman P. Overactive bladder syndrome:
432 evaluation and management. *Curr Urol* 2018; 11: 117–25.
- 433 23. The Japanese Continence Society, The Japanese Urological Association, eds. *Clinical*
434 *Guidelines for Overactive Bladder Syndrome*, 3rd edn, 2022.
- 435 24. Funada S, Luo Y, Uozumi R, *et al.* Multicomponent intervention for overactive bladder in
436 women: A randomized clinical trial. *JAMA Netw Open* 2024; 7: e241784.
- 437 25. Center for Outcomes Research and economic evaluation for health, National Institute of
438 Public Health, 2024. *Core to Evidence-Based Health*, Policy [cited 2024 Jun 7].
439 Guideline for Preparing Cost-Effectiveness Evaluation to the Central Social Insurance
440 Medical Council. Available from:
441 https://c2h.niph.go.jp/tools/guideline/guideline_en_2024.pdf.
- 442 26. Shiroiwa T, Noto S, Fukuda T. Japanese population norms of EQ-5D-5L and health
443 utilities index Mark 3: disutility catalog by disease and symptom in community settings.
444 *Value Health* 2021; 24: 1193–202.
- 445 27. Milsom I, Kaplan SA, Coyne KS, Sexton CC, Kopp ZS. Effect of bothersome overactive
446 bladder symptoms on health-related quality of life, anxiety, depression, and treatment
447 seeking in the United States: results from EpiLUTS. *Urology* 2012; 80: 90–6.
- 448 28. Omae K, Kurita N, Takeshima T, *et al.* Significance of overactive bladder as a predictor
449 of falls in community dwelling older adults: 1-year followup of the Sukagawa study. *J*
450 *Urol* 2021; 205: 219–25.
- 451 29. Husereau D, Drummond M, Augustovski F, *et al.* Consolidated health economic
452 evaluation reporting standards 2022 (CHEERS 2022) statement: updated reporting
453 guidance for health economic evaluations. *Pharmacoeconomics* 2022; 40: 601–9.

- 454 30. Funada S, Kawaguchi T, Terada N, *et al.* Cross-sectional epidemiological analysis of the
455 Nagahama study for correlates of overactive bladder: genetic and environmental
456 considerations. *J Urol* 2018; 199: 774–8.

457

458 Tables

459 **Table 1** Baseline characteristics of respondents (n=994)

Variables	Category	n	%
Age	Mean (SD)	73.2	(5.3)
	Median [IQR]	75	[68–77]
Gender	Male	516	51.9
BMI	Mean (SD)	22.6	(3.4)
	Median [IQR]	22.3	[20.5–24.3]
Educational attainment	College graduate or more	557	56.0
Equivalent household income	Lower half	409	41.2
	Upper half	360	36.2
	Decline to answer	225	22.6
Smoking status	Never	477	48.0
	Past	410	41.3
	Current	107	10.8
Alcohol consumption	Never	246	24.8
	Past	229	23.0
	Current	519	52.2
Presence of comorbidities	Hypertension	486	48.9
	Dyslipidemia	407	41.0
	Diabetes	166	16.7
	Low back pain	566	56.9
	Ischemic heart disease	91	9.2
	Stroke	63	6.3
	Chronic kidney disease	36	3.6
	Stress urinary incontinence	114	11.5
	Depression	50	5.0
	Other psychiatric disorders	44	4.4
OABSS	Mean (SD)	2.7	(2.2)
	Median [IQR]	2	[1–4]
Health utility value	Mean (SD)	0.90	(0.11)
	Median [IQR]	0.89	[0.83–1]
EQ-VAS score	Mean (SD)	75.2	(17.6)
	Median [IQR]	80	[70–88]

460 **Note.** BMI, body mass index; EQ VAS, EuroQol Visual Analogue Scale; IQR, interquartile range; LUTS,
461 lower urinary tract symptom; OABSS, Overactive Bladder Symptom Scale; SD, standard deviation.

462

Table 2 Prevalence of OAB and utility values among overall respondents

OAB severity*	n (%)	Mean utility value (SD)	Median utility value [IQR]
Non-OAB	836 (84.1)	0.913 (0.105)	0.894 [0.867–1]
Any OAB	158 (15.9)	0.855 (0.145)	0.885 [0.784–1]
Mild OAB	71 (7.1)	0.874 (0.131)	0.895 [0.823–1]
Moderate-to-severe OAB	87 (8.8)	0.840 (0.155)	0.844 [0.759–1]

Note. IQR, interquartile range; OAB, overactive bladder; SD, standard deviation.

*Mild OAB is defined as domain 3 (urgency) of overactive bladder symptom score (OABSS) ≥ 2 and total score of $3 \leq \text{OABSS} \leq 5$, and moderate-to-severe OAB is defined as domain 3 of OABSS ≥ 2 and total score of $6 \leq \text{OABSS} \leq 15$.

Table 3 Results of multivariable linear regression models in the main and sensitivity analyses

Variables	Category	β^*	SE	95% CI	<i>p</i> -value	<i>p</i> for trend
Main analysis (outcome measure: EQ-5D-5L)						
Presence or absence of OAB	Non-OAB			Reference		Not applicable
	Any OAB	−0.0472	0.0098	−0.0664 to −0.0279	<0.001	
Based on OAB severity	Non-OAB			Reference		<0.001
	Mild OAB	−0.0334	0.0136	−0.0602 to −0.0066	0.014	
	Moderate-to-severe OAB	−0.0591	0.0129	−0.0844 to −0.0339	<0.001	
Sensitivity analysis (outcome measure: EQ-VAS)						
Presence or absence of OAB	Non-OAB			Reference		Not applicable
	Any OAB	−5.42	1.55	−8.47 to −2.37	0.001	
Based on OAB severity	Non-OAB			Reference		<0.001
	Mild OAB	−4.36	2.16	−8.60 to −0.12	0.044	
	Moderate-to-severe OAB	−6.26	2.04	−10.26 to −2.27	0.002	

Note. EQ-5D-5L, the 5-level EuroQoL 5 dimensions version; EQ VAS, EuroQol Visual Analogue Scale; CI, confidence interval; OAB, overactive bladder; SE, standard error.

* Adjusted for age, sex, body mass index, educational attainment, equivalent household income, smoking status, alcohol consumption, and presence of comorbidities (hypertension, dyslipidemia, diabetes, low back pain, ischemic heart disease, stroke, chronic kidney disease, stress urinary incontinence, depression, and other psychiatric disorders).

Figure legends

Figure 1 Flow diagram of this study

Figure 2 Results of multivariable linear regression models in the subgroup analyses

* Adjusted for age, gender, body mass index, educational attainment, equivalent household income, smoking status, alcohol consumption, and presence of comorbidities (hypertension, dyslipidemia, diabetes, low back pain, ischemic heart disease, stroke, chronic kidney disease, stress urinary incontinence, depression, and other psychiatric disorders).

Abbreviations: CI, confidence interval; OAB, overactive bladder.

Figure 3 Results of multivariable linear regression models in the exploratory analyses

* Adjusted for age, gender, body mass index, educational attainment, equivalent household income, smoking status, alcohol consumption, and presence of comorbidities (hypertension, dyslipidemia, diabetes, low back pain, ischemic heart disease, stroke, chronic kidney disease, stress urinary incontinence, depression, and other psychiatric disorders).

Abbreviations: CI, confidence interval; OABSS, overactive bladder symptom score.





