Treatment Efficacy of Theophylline in ADYC5 Dyskinesia: A Retrospective Case Series Study

Dirk Taenzler^{1,2}, Frank Hause^{1,2,3} (ORCID: 0000-0002-6879-6944), Andreas Merkenschlager, Prof. Dr.⁴ (ORCID: 0000-0001-9959-2306), Andrea Sinz, Prof. Dr.^{1,2} (ORCID: 0000-0003-1521-4899)

¹Department of Pharmaceutical Chemistry and Bioanalytics, Martin-Luther-University Halle-Wittenberg

²Center for Structural Mass Spectrometry, Martin-Luther-University Halle-Wittenberg

³Institute of Molecular Medicine, Section for Molecular Cell Biology, Faculty of Medicine, Martin Luther University Halle-Wittenberg

⁴Department of Pediatrics, University of Leipzig Medical Center

Corresponding author:

Prof. Dr. Andrea Sinz Center for Structural Mass Spectrometry Martin Luther University Halle-Wittenberg 06120 Halle (Saale), Germany. Phone: +49-345-5525170 Email: andrea.sinz@pharmazie.uni-halle.de

Word Count: 2,346

Running Title:

Theophylline in ADYC5 Case Series

Financial Disclosure/Conflict of Interest:

The authors declare that there are no conflicts of interest regarding the publication of this paper. AS acknowledges financial support by the DFG (RTG 2467, project number 391498659 "Intrinsically Disordered Proteins-Molecular Principles, Cellular Functions, and Diseases", INST 271/404-1 FUGG, INST 271/405-1 FUGG), the Federal Ministry for Economic Affairs and Energy (BMWi, ZIM project KK5096401SK0), the region of Saxony-Anhalt, and the Martin Luther University Halle-Wittenberg (Center for Structural Mass Spectrometry).

Abstract

Background

ADCY5-related dyskinesia is a rare disorder caused by mutations in the ADCY5 gene resulting in abnormal involuntary movements. Currently, there are no standardized guidelines to treat this condition.

Objectives

The aim of this study is to evaluate the efficacy of theophylline administration in improving symptoms and quality of life in patients with ADCY5-related dyskinesia.

Methods

A retrospective study was conducted involving 12 patients (aged 2-34 years) with ADCY5-related dyskinesia. Participants completed a questionnaire about theophylline administration, including dosage, improvement of symptoms, adverse effects, and changes in quality of life. Data were analyzed for reported efficacy and side effects.

Results

Theophylline administration demonstrated substantial efficacy, with 92% (11 out of 12) of patients reporting significant improvements in their movement disorders. The average improvement score was 7.0 (\pm 1.9) on a 10-point scale. Notable improvements included reductions in severity and frequency of episodes, improved gait, more independent mobility, psycho-social well-being, and quality of sleep. Adverse effects were reported by 6 patients, including dystonia, speech worsening, headaches, nausea, impaired sleep, and agitation.

Conclusions

Theophylline shows substantial promise as a treatment option for ADCY5-related dyskinesia, improving various aspects of patients' quality of life and movement disorder symptoms. Further research is needed to optimize dosing, to understand long-term effects, and to explore combinational drug therapies. Despite the small cohort size and the retrospective nature of this study, the results support theophylline

administration to decrease dyskinetic movements and enhance overall quality of life in patients.

Introduction

ADCY5-related dyskinesia is a rare neurological condition caused by mutations in the ADCY5 gene that codes for the enzyme adenylyl cyclase 5. This enzyme is essential for intracellular signaling, as it converts ATP into cyclic AMP (cAMP), a secondary messenger involved in numerous cellular functions. Mutations in the ADCY5 gene cause increased enzymatic activity,¹ leading to an overproduction of cAMP and clinically resulting in abnormal involuntary movements referred to as dyskinesia.²

ADCY5-related dyskinesia manifests through a variety of impaired voluntary movements, including but not limited to choreoathetosis, dystonia, myoclonus and/or axial hypotonia,³ ultimately resulting in poor motor control. These symptoms typically emerge in infancy or early childhood, affecting the limbs, face, and neck.⁴ The movements are often continuous during waking hours and can disrupt sleep. They tend to worsen with stress, fatigue, or illness but may also occur without any identifiable trigger.⁴ The majority of individuals diagnosed with ADCY5-related dyskinesia are simplex cases, meaning they are the only affected member in their family, and their condition arises from a de novo pathogenic variant. Diagnosis is primarily based on clinical presentation and confirmed through genetic testing to identify mutations in the ADCY5 gene.⁵⁻⁷

Based on case reports, patients with ADCY5-related dyskinesia may benefit from deep brain stimulation (DBS).⁸⁻¹⁰ Bilateral pallidal DBS can significantly reduce hyperkinetic movements and paroxysomal episodes. While there is generally partial improvement patients often remain impaired and daily living activities may not significantly improve.¹¹

Currently, there are no randomized controlled trials or consensus guidelines for treating ADCY5-related dyskinesia. Case reports suggest that benzodiazepines, such as clonazepam, clobazam, diazepam, and lorazepam, can help managing dyskinetic episodes, also those associated with sleep.⁸ In a retrospective study involving 30 individuals, a beneficial effect of caffeine on hyperkinetic movements was observed.¹² This effect is thought to be attributed to the attenuating action of caffeine as an adenosine 2A antagonist, which is believed to lead to a reduction of cAMP levels in the striato-pallidal projection neurons.¹²

4

We were able to demonstrate that treatment with the purine derivatives caffeine, theophylline, and istradefylline reduced ADCY5-catalyzed cAMP production in cell lines overexpressing wild-type and mutant ADCY5. The most pronounced effects on cAMP reduction were observed in ADCY5 R418W mutant cells¹³, a mutation found in the majority of individuals with ADCY5-related dyskinesia¹⁴. Following these findings, a slow-release theophylline formulation was administered to a preschool-aged patient with ADCY5-related dyskinesia, resulting in a striking improvement of symptoms, surpassing the effects of caffeine that had previously been administered to the same patient.¹³

Based on these encouraging results, we systematically present the first comprehensive account of the effects of theophylline administration in a series of retrospective reports from patients with ADCY5-related dyskinesia.

Patients and Methods

Patients and/or their parents, if the patients were too young or unable to complete the questionnaire themselves, were asked to fill out a form that had been specifically designed for this study. The questionnaire targeted individuals who had administered a slow-release formulation of theophylline to treat ADCY5-related dyskinesia. The questionnaire was made available to interested participants via the patient organization portal adcy5.org. It was provided in German, English, French, and Spanish. Interested individuals were given the opportunity to answer the questionnaire anonymously. The questionnaire was uploaded by DT, and the collection of responses including their anonymization was also conducted by DT. Anonymized forms were forwarded to AM for analysis.

The aim of the ADCY5 questionnaire for parents was to systematically gather comprehensive information about symptoms, treatment responses, and overall patients' quality of life with ADCY5-related movement disorders and potential changes following theophylline administration. In addition to demographic questions, the questionnaire includes items related to the patients' genotype, cardinal dyskinetic symptoms and their frequency, current theophylline dosage, positive and adverse effects of theophylline administration, as well as former and ongoing (co-)medications for the treatment of ADCY5-related dyskinesia.

5

The study protocol was approved by the Ethics Committee of the Medical Faculty, University of Leipzig, on April 30, 2024 and listed in the German Clinical Trials Register (DRKS00034720), on Juli 25, 2024. All patients, or their authorized representatives, provided written informed consent prior to participation in this study. Data produced in the present study are available upon reasonable request to the authors. The questionnaire provided to the patients is available as online supplemental data.

Results

Demographics

The study included 12 patients, comprising 7 females and 5 males. The patients' age ranged from 2 to 34 years, with a median age of 8 years. All patients were characterized by ADYC5-related dyskinesia with a diverse range of baseline hyperkinetic and paroxysmal movement disorders. Genotyping data of the ADCY5 gene was provided by the 12 participants in the questionnaire and confirmed mutations in the ADCY5 gene, with R418W being the most prominent mutation, present in 9 out of 12 individuals.

Theophylline Efficacy

Theophylline administration demonstrated substantial efficacy across the study cohort. Out of the 12 patients, 11 reported improvements in their movement disorder symptoms, constituting 92% of the cohort. The average improvement score on a 10-point scale was approximately 7.0, with a standard deviation of 1.9, indicating a high level of efficacy with minor variability.

These improvements were comprehensive, affecting various aspects of the disorder symptoms. Most patients noted a reduction in the severity and frequency of their episodes (8 of 12), with a substantial number reporting reduced duration per episode (8 of 12) (see Table 1).

Additional patient-reported outcomes include improvements in gait, more independent mobility, and increased psycho-social well-being in terms of confidence and happiness, as well as improved quality of sleep (on a 10-point scale: M = 5.0, SD = 3.8), significantly contributing to the overall improved functionality and quality of life reported by the patients.

Theophylline Dosing and Tolerance

Theophylline was administered as a slow-release formulation at a daily dose ranging between 200 mg and 1,050 mg, adjusted to the patient's body weight (9.5 to 28.6 mg/kg/day). The medication was generally well-tolerated. Adverse effects were reported by 6 patients. These included pronounced dystonia (1 of 6), worsening of speech (1 of 6), headaches of idiopathic origin (1 of 6), nausea (1 of 6), impaired sleep (1 of 6), and episodes of agitation (2 of 6).

Other Treatments

Prior to theophylline administration, 10 patients had been treated with different regimens, including caffeine, levetiracetam, and benzodiazepines, with mixed results. After starting theophylline administration, 8 patients continued to use adjunct therapies, such as zopiclone and duloxetine, which were generally well-tolerated and enhanced overall treatment efficacy.

).	Age Range (y)	Gender (F/M)	Body Weight (kg)	ADCY5 Mutant	Theophylline Dosage Daily (mg)	Theophylline Dosage Daily (mg/kg/d)	Blood Level (mg/l)	Improvement Movement Disorder (0: No Improvement - 10: Drastic Improvement)	Improvement of Episode Severeness (less severe) (Yes/No)	improvement of Episode Duration (shorter) (Yes/No)	Improvement of Episode Frequency (less frequent) (Yes/No)	Change in QoL (0: significantly deteriorated - 6: significantly improved)	Change in QoS (0: No Improvement - 10: Drastic Improvement)	Adverse Effects
1	0-5	М	10	R418W	200	20.00	10	6	Yes	Yes	Yes	6	0	١
2	31-35	F	51	M696K	700	13.73	6.1	8.5	Yes	Yes	Yes	6	10	١
3	11-16	M	32	R418W	400	12.50	10.3	9	Yes	Yes	Yes	6	9	N
4	6-10	г г	21	R410W	600	28.57	17.5	0	Voc	Voc	Voc	6	9	
6	6-10	F	21	R418W	500	20.57	NA	4	No	No	No	5	9	More Dyston
7	6-10	F	22	R62L, R412L, R45L	550	25.00	14.3	NA	NA	NA	NA	4	2	Episodes Agitatio
8	6-10	М	21	R418W	200	9.52	2	7.5	Yes	Yes	Yes	5	6.5	Worsening
9	11-16	F	29	R418W	600	20.69	85	8	Yes	Yes	Yes	5	8	N
0	6-10	М	45	R418Q	750	16.67	11.4	8	Yes	Yes	Yes	6	4	Headach (Idiopathic Origi
1	6-10	F	18	R418W	200	11.11	NA	4	NA	NA	NA	6	3	Nause
2	41-45	М	66	R418W	1,050	15.91	8	4.5	NA	NA	NA	5	0	Episodes Agitatio Impaired Slee
JIE	viation	S. QUL.	Quanty	Of Life	, QUO. QUA	ity of Sieep								

Discussion

The present study indicates that theophylline administration substantially improves the symptoms and quality of life in patients with ADCY5-related dyskinesia, as evidenced by the high level of reported efficacy and alleviated symptoms.

Theophylline administration demonstrated substantial efficacy, with 92% of patients reporting improvements in their movement disorder symptoms. The average improvement score was 7.0 on a 10-point scale, with a standard deviation of 1.9, indicating a high overall efficacy. Improvements were noted in various symptom domains, including the severity and frequency of episodes, with 8 out of 12 patients reporting shorter episodes. Additionally, patients experienced improvements in gait, more independent mobility, enhanced psycho-social well-being, and higher quality of sleep, significantly contributing to their overall functionality and quality of life. Only minor adverse effects have been reported.

We are aware that there are several aspects in this study that should be considered. Firstly, the retrospective nature of the study introduces a potential bias and limits the ability to draw causal inferences. It is uncertain whether the participants were aware of the results from the initial case report involving the preschool-aged patient.¹³ Patients #2, #6, and #12 (see Table 1) had just started theophylline administration when completing the questionnaire. The relatively small cohort size is attributed to the rarity of ADCY5-related dyskinesia, which poses challenges to the feasibility of conducting randomized trials.

It has to be considered that the administration of theophylline might not improve sleep quality in ADCY5-related dyskinesia: Three patients (#1, #6, and #12, see Table 1) reported no improvement in sleep quality after theophylline administration. For patient #1, theophylline administration began at an age before the typical symptoms of the disease manifested. As evident from the overall improvement in quality of life for patient #1, theophylline has a positive effect, but this patient never experienced sleep disruption due to the disease. Patient #6 underwent surgery for DBS before administering theophylline, therefore it is difficult to attribute the effects to the theophylline treatment alone. Finally, patient #12 indicated that the reduced sleep quality might also be credited to the concurrent use of diazepam. In conclusion, while theophylline demonstrates substantial promise as a treatment for ADCY5-related dyskinesia, further research is warranted to optimize dosing, to better understand long-term effects, and to explore the benefits of combinational drug therapies, especially regarding above mentioned limitations. Based on our results, we recommend theophylline administration in patients with ADCY5-related dyskinesia. Blood drug levels should optimally range between 15 to 20 mg/l to decrease dyskinetic movements and to improve overall quality of life.

Authors' Roles

DT, AM, and AS designed the study. DT informed willing participants about, and collected and anonymized the answers to the questionnaire. DT, AM, and FH analyzed the data. DT, FH, and AS wrote the final manuscript.

Financial Disclosures of all authors

AS acknowledges financial support by the DFG (RTG 2467, project number 391498659 "Intrinsically Disordered Proteins-Molecular Principles, Cellular Functions, and Diseases", INST 271/404-1 FUGG, INST 271/405-1 FUGG), the Federal Ministry for Economic Affairs and Energy (BMWi, ZIM project KK5096401SK0), the region of Saxony-Anhalt, and the Martin Luther University Halle-Wittenberg (Center for Structural Mass Spectrometry).

References

¹ Chen YZ, Friedman JR, Chen DH, et al. Gain-of-function ADCY5 mutations in familial dyskinesia with facial myokymia. Ann Neurol 2014; 75(4): 542-549.

² Doyle TB, Hayes MP, Chen DH, Raskind WH, Watts VJ. Functional characterization of AC5 gain-of-function variants: Impact on the molecular basis of ADCY5-related dyskinesia. Biochem Pharmacol. 2019; 163: 169–77.

³ Hisama FM, Friedman J, Raskind WH, Bird TD. ADCY5 Dyskinesia. In: Adam MP, Mirzaa GM, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2024.

⁴ Vijiaratnam N, Bhatia KP, Lang AE, Raskind WH, Espay AJ. (2019). ADCY5-related dyskinesia: Improving clinical detection of an evolving disorder. Mov Disord Clin Pract 2019; 6(7): 512-520.

⁵ Bohlega SA, Abou-Al-Shaar H, AlDakheel A, et al. Autosomal recessive ADCY5related dystonia and myoclonus: Expanding the genetic spectrum of ADCY5-related movement disorders. Parkinsonism Relat Disord 2019: 64: 145-149.

⁶ Carapito R, Paul N, Untrau M, et al. A de novo ADCY5 mutation causes early-onset autosomal dominant chorea and dystonia. Mov Disord 2015; 30(3): 423-427.

⁷ Carecchio M, Mencacci NE, Iodice A, et al. ADCY5-related movement disorders: Frequency, disease course and phenotypic variability in a cohort of paediatric patients. Parkinsonism Relat Disord 2017: 41:37-43.

⁸ de Almeida Marcelino AL, Mainka T, Krause P, Poewe W, Ganos C, Kuhn AA. Deep brain stimulation reduces (nocturnal) dyskinetic exacerbations in patients with ADCY5 mutation: a case series. J Neurol 2020; 267(12): 3624-3631. ⁹ Eisenberg HJ, Malinova V, Mielke D, Bahr M, Goricke MB, van Riesen C. ADCY5induced dyskinetic storm rescued with pallidal deep brain stimulation in a 46-year-old man. Mov Disord Clin Pract 2021; 8(1): 142-144.

¹⁰ Meijer IA, Miravite J, Kopell BH, Lubarr N. Deep brain stimulation in an additional patient with ADCY5-related movement disorder. J Child Neurol 2017; 32(4): 438-439.

¹¹ Menon PJ, Nilles C, Silveira-Moriyama L, et al. Scoping Review on ADCY5-Related Movement Disorders. Mov Disord Clin Pract 2023;10(7): 1048-1059.

¹² Méneret A, Mohammad SS, Cif, L, et al. Efficacy of caffeine in ADCY5-related dyskinesia: A retrospective study. Mov Disord 2022; 37(6): 1294-1298.

¹³ Tänzler D, Kipping M, Lederer M, et al. Effects of theophylline on ADCY5 activation—From cellular studies to improved therapeutic options for ADCY5-related dyskinesia patients. PLoS One 2023;18(3): e0282593.

¹⁴ Ferrini A, Steel D, Barwick K, Kurian MA. An Update on the Phenotype, Genotype and Neurobiology of ADCY5-Related Disease. Mov Disord 2021; 36(5): 1104-1114.