- 1 Load and muscle dependent changes in triceps surae motor unit firing properties and
- 2 motor unit firing-torque relationships in individuals with non-insertional Achilles
- 3 tendinopathy
- 4
- 5 Authors: Ignacio Contreras-Hernandez¹, Deborah Falla¹, Michail Arvanitidis¹,
- 6 Francesco Negro², David Jimenez-Grande¹, Eduardo Martinez-Valdes¹
- 7
- 8 Corresponding author: Eduardo Martinez-Valdes¹
- 9 Lecturer (Assistant Professor) in Physiotherapy
- 10 Centre of Precision Rehabilitation for Spinal Pain (CPR Spine)
- 11 School of Sport, Exercise and Rehabilitation Sciences
- 12 College of Life and Environmental Sciences
- 13 University of Birmingham
- 14 Edgbaston B15 2TT, United Kingdom
- 15 T: +44 (0) 121 4158187
- 16 M: +44 (0) 7961 548851

Email: <u>e.a.martinezvaldes@bham.ac.uk</u>

Author Affiliations

- ¹⁷¹Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), School of Sport,
- 18 Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences,
- 19 University of Birmingham, Birmingham, UK.
- ²Department of Clinical and Experimental Sciences, Università degli Studi di Brescia,
- 21 Brescia, Italy.
- 22
- 23 Email: Ignacio Contreras-Hernandez¹ <u>iac921@student.bham.ac.uk</u> Deborah Falla¹
- 24 d.falla@bham.ac.uk Michail Arvanitidis MXA1016@student.bham.ac.uk -
- 25 Francesco Negro² <u>francesco.negro@unibs.it</u> David Jimenez-Grande
- 26 dbj893@student.bham.ac.uk Eduardo Martinez-Valdes¹
- 27 <u>e.a.martinezvaldes@bham.ac.uk</u>
- 28 WORD COUNT MAIN TEXT: 8963 (excluding title page, abstract, references,
- 29 figures, tables, and acknowledgments)

30 ABSTRACT

Non-insertional Achilles tendinopathy (NIAT) induces morpho-mechanical changes 31 32 to the Achilles tendon (AT). However, evidence on how triceps surae motor unit firing properties are influenced by altered tendon mechanics in NIAT is limited. This study 33 34 investigated motor unit firing properties (mean discharge rate (DR), recruitment and de-recruitment thresholds, and discharge rate variability (COVisi)), motor unit firing-35 torque relationships (cross-correlation coefficient between cumulative spike train 36 37 (CST) and torque, and neuromechanical delay), and neural drive distribution 38 (connectivity strength and functional networks) of the medial gastrocnemius (MG), 39 lateral gastrocnemius (LG), and soleus (SO) muscles during isometric plantarflexion 40 contractions at 10%, 40%, and 70% maximal voluntary contraction (MVC) using high-density surface electromyography on 26 individuals with NIAT and 25 healthy 41 controls. Furthermore, AT's morpho-mechanical properties (thickness, cross-42 43 sectional area, length and stiffness) were assessed via ultrasound imaging. NIAT individuals showed reduced tendon stiffness and increased thickness (p<0.01). 44 Motor unit properties changed in a load and muscle-dependent manner. LG DR 45 increased (p=0.002) and de-recruitment threshold decreased (p=0.039) at 70%MVC 46 47 in the NIAT group compared to controls. The CST-torque cross-correlation 48 coefficient of the LG decreased at 10%MVC (p<0.0001) and increased at 70%MVC (p=0.013) in the NIAT group. Connectivity strength for the 0-5 Hz and 5-15 Hz 49 frequency bands decreased (p<0.01) in the NIAT group at 10%MVC. This study 50 shows that individuals with NIAT exhibit load-dependent changes in motor unit firing 51 52 properties, motor unit-torque relationships, and neural drive distribution to the triceps 53 surae. These alterations may be due to muscle-specific compensations for the 54 modified mechanical properties of the AT.

55

Keywords: High-density surface electromyography; motoneuron, stiffness, tendon
 degeneration, tendinosis, neuromechanics.

58

59 ABSTRACT WORD COUNT: 249

- 60
- 61
- 62

63 KEY POINTS

- 64
- Individuals with non-insertional Achilles tendinopathy (NIAT) have changes of
 the neural drive to the lateral gastrocnemius (LG) muscle and altered
 contribution of the LG to the net plantarflexion torque.
- Individuals with NIAT show a more uneven distribution of neural drive to the
 triceps surae muscle at low force levels, characterized by reduced
 intermuscular coherence between the medial and lateral gastrocnemius in the
 0-5 Hz and 5-15 Hz bands compared to the control group.
- Our findings support the idea that the LG may have a central role in the
 pathophysiology of this condition, possibly affecting the load transmission to
 the Achilles tendon (AT).
- 75
- 76

77 INTRODUCTION

78

79 Non-insertional Achilles tendinopathy (NIAT) is a common debilitating tendon disorder in the lower extremity, characterized by an insidious onset of pain 80 81 accompanied by swelling, stiffness and thickening in the midportion of the Achilles 82 tendon (AT) (Maffulli et al., 1998; van Dijk et al., 2011; Bahari et al., 2023; Sara et al., 2023). NIAT can affect athletes and the general population, causing disability, 83 functional impairment and decreasing performance (Kvist, 1994; de Jonge et al., 84 2011). It's estimated that more than half (55-65%) of the AT problems occur about 2-85 7 cm proximal to superior border of the calcaneus (Järvinen et al., 2005). The 86 87 etiology of the NIAT remains debated (Maffulli et al., 2020); however, excessive loading of the tendon is considered the major causative factor (Leadbetter, 1992). 88 Recent evidence suggests that uneven activation of the triceps surae muscle may 89 influence the magnitude and distribution of AT load, stress and strain (O'Neill et al., 90 91 2015; Debenham et al., 2017); therefore, deficits in the performance of this muscle 92 group has been identified as a key factor for the development of NIAT (O'Neill et al., 93 2019). In humans, plantarflexion is primarily accomplished by the triceps surae 94 muscle group, which comprises the medial gastrocnemius (MG), lateral 95 gastrocnemius (LG) and soleus (SO) muscles (Héroux et al., 2014). The resultant 96 force generated by these muscles is then transmitted to the AT, which links the 97 triceps surae muscle to the calcaneus (Crouzier et al., 2020). The AT is composed of 98 three subtendons, arising from each of the triceps surae muscles (Crouzier et al., 99 2020); together, these subtendons exhibit a twisted structure, with the deep layer of the tendon formed by fibers from the LG and SO, and the superficial layer formed by 100 101 fibers from the MG and SO, with a wide range of rotation of the subtendons between 102 individuals (Szaro et al., 2009). Ultrasound imaging during isometric contractions has 103 revealed nonuniform displacements within the Achilles tendon, consistent with an 104 uneven distribution of load across the tendon (Arndt et al., 1998; Clark & Franz, 105 2018). Additionally, 3D models suggest that the nonuniform distribution of load within 106 the AT is largely due to the distribution of force among the MG, LG, and SO muscles 107 (Handsfield et al., 2017). Consequently, it has been proposed that an imbalance of 108 the force produced by these three muscles may cause an heterogenous load within 109 the tendon, contributing to the development of NIAT(Bojsen-Møller & Magnusson, 2015; Hug & Tucker, 2017). 110

111 The morpho-mechanical properties of the Achilles tendon in individuals with NIAT have been studied extensively. The morphological changes include an 112 113 increase in tendon thickness, cross-sectional area (CSA), and volume (Shalabi et al., 2004; Arya & Kulig, 2010; Child et al., 2010; Grigg et al., 2012; Docking & Cook, 114 115 2016) and the mechanical changes comprise an increase in longitudinal and 116 transverse strain, hysteresis, and lower stiffness and Young's modulus compared to healthy tendons under the same tensile load (Arya & Kulig, 2010; Child et al., 2010; 117 118 Wang et al., 2012; Chang & Kulig, 2015). Alterations in the AT morpho-mechanical 119 properties are relevant because of their potential to modify the stress-strain patterns 120 experienced by the AT during muscle contraction (Hansen et al., 2017), and 121 muscle's capacity to function within the force-length curve's optimal region (Arya & 122 Kulig, 2010), possibly influencing the triceps surae motor unit firing properties.

123 For the generation of movement, the central nervous system (CNS) must 124 effectively coordinate the many degrees of freedom of the musculoskeletal system, 125 considering the nonlinear features of muscles and the dynamic interactions between 126 the segments of the body, as well as the interactions between the body and the 127 environment (d'Avella & Bizzi, 2005). This complex task might be simplified by 128 organizing a modular and hierarchical control (Full & Koditschek, 1999; Loeb et al., 129 1999; Bizzi et al., 2002). It has been hypothesized that the CNS uses a set of muscle 130 synergies, the coherent activation in space or time of a group of muscles, as output 131 modules. Thus, supraspinal and afferent signals combine with a determined number 132 of muscle synergies to generate a variety of muscle activation patterns (d'Avella & 133 Bizzi, 2005). Based on this, it has been theorized that alterations in how the CNS 134 control the coordination of the triceps surae muscles could create uneven loading of 135 the AT and contribute to tendinopathy (Cook & Purdam, 2009). Although these 136 muscles act as agonists that share the same common distal tendon, they exhibit 137 anatomical, neurophysiological, and functional differences, indicating diverse 138 functional roles (Héroux et al., 2014). These functional roles have been linked to unique properties in motor unit firing properties between muscles when performing 139 140 different tasks (Héroux et al., 2014).

The force exerted by a muscle during a voluntary contraction depends, partly, on the recruitment and discharge rate of motor units (Enoka & Duchateau, 2017). Hence, alterations in motor performance may be the consequence of the inability of the CNS to increase the recruitment and/or the discharge rate of the pool of motor

145 units (Enoka & Duchateau, 2017). Within this framework, the most reliable approach 146 to investigate how the CNS modulates the recruitment and discharge firing rates of 147 the triceps surae motor units during a voluntary contraction is through the individual analysis of the MG, LG, and SO muscles (Hug et al., 2021; Fernandes et al., 2023). 148 149 To the best of our knowledge, only one study has investigated the motor unit firing 150 rates properties of the triceps surae muscles in individuals with NIAT. Fernandes et al. 2021 reported no differences in the discharge rate of the LG between a group of 151 152 runners with NIAT and a control group at low forces; however, they observed that in the NIAT group the DR of the LG remained unchanged as the target torque 153 154 increased from 10% to 20% MVC, suggesting that the neural drive to the LG is affected in individuals with NIAT (Fernandes et al., 2023). However, the authors only 155 156 assessed motor unit firing parameters during low-force isometric plantarflexion 157 contractions in a limited group of participants, without examining variations in AT's 158 morpho-mechanical properties and the distribution of the neural drive received by the 159 triceps surae. Considering the stress-strain properties of tendons, it is known that the 160 rate of tendon deformation changes according to the load placed on the tendon 161 (Kongsgaard et al., 2011). Furthermore, the pain experienced by individuals with 162 NIAT increases with load (Magnusson et al., 2010). Therefore, studying these 163 properties across a wider range of torgues is essential to understand how different 164 loading conditions affect tendon behavior and motor unit firing patterns across the 165 triceps surae muscle. Moreover, since common modulations in firing rate reflect the 166 effective neural drive received by muscles (i.e., common synaptic input (Farina et al., 167 2014), it is crucial to examine these modulations between the different heads of the triceps surae and their relationship with the resultant torque. This information could 168 provide insights into whether NIAT-induced changes in tendon structure might alter 169 the contribution of each triceps surae muscle head to the produced torque. 170

We aimed to investigate triceps surae motor unit firing properties (mean 171 172 discharge rate (DR), recruitment/de-recruitment thresholds and discharge rate variability (coefficient of variation of the interspike interval (COVisi)), motor unit firing-173 174 torque relationships (cross-correlation coefficient between cumulative spike train 175 (CST) and torque, and neuromechanical delay (NMD)), and common motor unit firing 176 modulations between triceps surae muscle heads (connectivity strength and 177 muscular functional networks via intermuscular coherence) in individuals with NIAT 178 at low, intermediate and high force levels during isometric plantarflexion

contractions. In addition, morpho-mechanical properties of the AT (length, thickness, cross-sectional area (CSA) and stiffness) were determined using B-mode ultrasonography and shear-wave elastography (SWE). Our primary hypothesis is that NIAT-induced alterations of the AT's morpho-mechanical properties would induce load and muscle-dependent changes in motor unit firing properties, motor unit firing-torque relationships and unbalanced distribution of the neural drive received by the triceps surae.

186

187 METHODS

188

189 Ethical approval

This study was approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee, University of Birmingham, UK (ERN_20-0604A).

193

194 **Participants**

Twenty-five healthy (17 males, 8 females, 28.60 ± 3.92 years, 74.00 ± 11.57 kg, 171.10 ± 9.22 cm) and twenty-six participants with NIAT (14 males, 12 females, 29.04 ± 8.46 years, 75.09 ± 17.20 kg, 173 ± 9.16 cm) were recruited from the University of Birmingham staff/student population and the local community via leaflets, e-mail, and social media.

200 Inclusion criteria included men or women aged 18 to 55 years old. This range 201 was chosen to prevent the influence of ageing-related alterations in tendon 202 properties, since previous studies have found that the AT has lower stiffness and 203 Young's modulus in older individuals (Lindemann et al., 2020). For the NIAT group inclusion criteria were as follows: presence of NIAT determined by an experienced 204 205 physiotherapist based on defined clinical findings (VISA-A (Robinson et al., 2001), 206 NRS (Numeral Rating Scale) (Alghadir et al., 2018) and pain for more than 3 months 207 (Beyer et al., 2015)), physical examination and ultrasound imaging. Ultrasound 208 imaging included assessing echoic pattern (focal hypoechoic and hyperechoic areas 209 within the tendon) and tendon thickness (focal or diffuse thickening) (Bleakney & 210 White, 2005). VISA-A scores less than 90 were considered as a reference to identify 211 individuals with NIAT (Iversen et al., 2012). Due to the high variability in the NRS 212 scores in individuals with NIAT(Rompe et al., 2009; Beyer et al., 2015), we considered individuals with an NRS score \geq 2. Inclusion criteria for the healthy group included confirmation of a healthy tendon determined by the same physiotherapist through physical examination and ultrasound imaging.

216 The exclusion criteria for both groups were as follows: (1) current or history of 217 chronic neurological, respiratory, or cardiovascular diseases, (2) systemic or 218 inflammatory conditions including rheumatic, neuromuscular disorders, and 219 malignancy, and (3) history of lower limb surgery. Specific exclusion criteria for the 220 participants with NIAT were if they had undergone a corticosteroid injection in the Achilles tendon within the last 12 months or if they had been diagnosed with 221 222 insertional Achilles tendinopathy. Specific exclusion criteria for the healthy group 223 were history of NIAT, lower limb surgery or pain/injury in the lower limbs within the 224 previous 6 months.

225

226 Study design

This cross-sectional study was conducted from October 2021 to April 2023 at a laboratory within the Centre of Precision Rehabilitation for Spinal Pain (CPR Spine), University of Birmingham, UK. The study was conducted according to the declaration of Helsinki and participants signed written informed consent prior to participation. The guideline for Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) was used to facilitate reporting (Vandenbroucke *et al.*, 2007).

Participants made a single visit to the laboratory for the experimental session (2.5 hours). They were instructed to avoid any vigorous physical activity within 24 hours prior to testing. The assessed leg was randomized in the control group, and the leg with tendinopathy symptoms was evaluated in the NIAT group. In the case of individuals with bilateral symptoms, the most painful leg was assessed.

239

240 Experimental setup and tasks

Experimental sessions included completing a battery of questionnaires, physical examination, ultrasonography of the AT, high-density surface electromyography (HD-sEMG) of the triceps surae muscles and torque recordings during plantarflexion isometric contractions. A representation of the experimental setup is shown in **Figure 1**.



Figure 1. Representation of the experimental setup.

247

246

Anthropometric data (age, gender, weight, height, and foot dominance) was 248 249 obtained across the participants. Foot preference in specific daily activities (foot 250 dominance) was determined using a behavioral foot-preference inventory (Chapman 251 et al., 1987). For the NIAT group, we obtained information regarding their symptoms 252 (intensity of pain (NRS), duration of symptoms, and presence of bilateral symptoms). 253 Then, all participants were asked to complete a battery of questionnaires, which 254 included the Victorian Institute of Sports Assessment – Achilles Questionnaire 255 (VISA-A) (Robinson et al., 2001), the International Physical Activity Questionnaire 256 short form (IPAQ-SF) (Craig et al., 2003), Foot and Ankle Ability Measure (FAAM) 257 (Martin et al., 2005), Pain Catastrophizing Scale (PCS) (Sullivan et al., 1995) and 258 Tampa Scale for Kinesiophobia (TSK) (Kori, 1990). Then, participants lay prone on 259 the chair of a Biodex System 3 dynamometer (Biodex Medical System), with their 260 knees extended and their assessed foot tightly strapped on the footplate. The ankle 261 was positioned in 0° of plantarflexion with the axis of the dynamometer aligned with 262 the inferior tip of the lateral malleolus. The pelvis was stabilized with a strap to 263 minimize compensatory movements. Ultrasonography was used to determine the 264 length, thickness, and cross-sectional area of the AT (see procedure below). Then,

the skin was shaved, cleaned, and prepared, and the electrodes grids were placed on the MG, LG, and SO muscles (see details below). Following the placement of the electrode grids, we assessed AT stiffness through SWE during rest conditions. HDsEMG was used to confirm that the muscles were not active during these measurements as this could influence the estimation of stiffness.

270 Next, participants performed a warm-up protocol consisting of 3 submaximal isometric plantarflexion contractions at their perceived 30% of maximal voluntary 271 272 force for 5 seconds with 30 seconds rest between contractions. The maximal voluntary contraction (MVC) was then determined during 3 isometric plantarflexion 273 274 contractions of 5 seconds with 2 minutes rest between contractions (Martinez-Valdes et al., 2018) at 0° of plantarflexion. The highest MVC value was used as the 275 276 reference maximal torgue. Then, we measured the electromyographic activity of the 277 MG, LG, and SO muscles during two isometric plantarflexion contractions at 10, 40 and 70% MVC (10% MVC/s ramp-up, 10 s hold, 10% MVC/s ramp-down and 30 s 278 rest) with HD-sEMG. Visual feedback of the torque produced by the participant was 279 280 provided through a computer monitor positioned 1 m in front of the participant. The 281 order of the contractions at different target torque levels (10, 40, and 70% MVC) was 282 randomized using а randomization app (Randomizer, https://play.google.com/store/apps/details?id=com.giannis.randomizer&hl=en_US). A 283 284 schematic representation of the experimental session is shown in Figure 2. 285

- ____
- 286
- 287



288

Figure 2. Schematic of the experimental procedure. The order of the contractions performed at each target torque (10, 40, and 70% MVC) was randomized. VISA-A, Victorian Institute of Sports Assessment- Achilles Questionnaire; IPAQ-SF, International Physical Activity Questionnaire short form; FAAM, Foot and Ankle Ability Measure; PCS, Pain Catastrophizing Scale; TSK, Tampa Scale for Kinesiophobia; MVC, maximum voluntary contraction.

294

295 Ultrasonography

All ultrasound images were obtained using an ultrasound imaging device equipped with SWE (LOGIQ S8 GE Healthcare, Milwaukee, USA). Tendon thickness, length, and cross-sectional area were recorded in B-mode with a 16-linear array probe (50 mm, 4-15 MHz). SWE was recorded in elastography mode with a 9linear array probe (44 mm, 2-8 MHz).

301 The morphological properties of the AT were determined using an adaptation of the protocol developed by Arya and Kulig (Arya & Kulig, 2010). The ultrasound 302 303 probe was placed longitudinally over the posterior part of the heel with the ankle at 304 0° of plantarflexion, and the calcaneal notch was identified. Then, a fine wire (3.2 x 305 40 mm) was used under the probe to create an artifact in the ultrasound image. The 306 artifact was aligned with the calcaneal notch and the corresponding point was 307 marked on the skin with a marker (this mark represented the AT's insertion). The 308 ultrasound probe was moved proximally to identify the musculotendinous junction of 309 the MG, and again, the wire was aligned with the musculotendinous junction of the 310 MG and the corresponding point was marked on the skin. The distance between 311 these two marks (calcaneal notch and musculotendinous junction of the MG

represented the length of the AT). Subsequently, marks were made at 2, 4 and 6 cm above the AT's insertion and these marks were used as reference to place the middle part of the probe in the longitudinal plane to obtain the thickness of the AT at 2, 4 and 6 cm of its insertion. Similarly, these reference marks were used to place the probe in the transversal plane and determine the cross-sectional area at 2, 4 and 6 cm of the AT's insertion. Three ultrasound images were obtained in the longitudinal and transverse planes for each mark.

For the HD-sEMG electrode grid placement, a tape and marker were used to 319 draw a line following the direction of the AT. This line and the distal 320 321 musculotendinous junction of the MG were used as references for the positioning of 322 the electrode grids. Briefly, for the MG electrode grid, a mark was made 10 cm 323 above the distal musculotendinous junction of the MG and 4 cm medial to the line 324 that followed the direction of the AT. Similarly, for the LG electrode grid, the leg was 325 marked 10 cm above the distal musculotendinous junction of the MG and 4 cm 326 lateral to the line that followed the direction of the AT. Likewise, for the SO electrode 327 grid, the leg was marked 5 cm below the distal musculotendinous junction of the MG 328 and 4 cm lateral to the line that followed the direction of the AT. The central 329 electrodes of the HD-sEMG grids (electrode in row 7 and column 3) were placed on 330 top of all these marks.

331 For the mechanical parameters (i.e., estimated stiffness), the ankle was 332 placed at 0° of plantarflexion and the ultrasound probe was placed longitudinally, 333 with the middle part of the probe located at 4 cm above the AT's insertion. A probe 334 holder was used to avoid applying pressure over the tendon. A trial SWE 335 measurement was done to check for possible voids in the stiffness estimation and if 336 voids were detected, the probe was removed, ultrasound gel was added, and the probe was placed again. Elastography images were acquired for 12 seconds (twice). 337 338 Due to the equipment properties, as a SWE image was obtained every 2.4 seconds; 339 thus, to obtain 4 SWE images, the elastography measurements lasted 12 seconds.

340

341 HD-sEMG and torque recordings

Participants' skin was shaved, gently abraded (Nuprep, Skin Prep Gel, Weaver and Company, Aurora, Colorado), cleaned with water and dried with tissue paper. Three two-dimensional (2D) adhesive grids (OT Bioelettronica, Italy) of 13 x 5 equally spaced electrodes (each of 1 mm diameter, with an inter-electrode distance

of 8 mm) were used to record the HD-sEMG signals from the MG, LG, and SO muscles (position of each grid was described above). The HD-sEMG grid was prepared by attaching a double-sided adhesive foam to the grid surface (SPES Medica, Genova, Italy) and by filling the grid cavities with conductive paste which provided adequate electrode-skin contact (AC-CREAM, SPES Medica, Genova, Italy).

352 HD-sEMG signals were converted by a 16-bit analogue-digital converter (Quattrocento- OT Bioelecttronica, Torino, Italy). Signals were amplified by a factor 353 of 150, sampled at 2048 Hz, and filtered with a band-pass filter (bandwidth: 10-500 354 355 Hz, first order, -3 dB) (Arvanitidis et al., 2019). HD-sEMG signals were acquired in 356 monopolar mode with ground electrodes (WhiteSensor WS, Ambu A/S, Ballerup, 357 Denmark) located in the head of the fibula and with a wet strap in the thigh of the 358 assessed leg. Electrode grids and ground electrodes were connected to an bioelectrical amplifier (Quattrocento-OT-Bioelecttronica, Torino, Italy) and the torque 359 360 exerted by the participants was obtained with a Biodex System 3 dynamometer 361 (Biodex Medical System), which was synchronized with the HD-sEMG signals 362 through the auxiliary input of the EMG amplifier (Arvanitidis et al., 2019).

363

364 Image analysis

365 Analysis of Ultrasound images. After obtaining the ultrasound images, a 366 reference line of 1 cm was drawn using the ultrasound device tools. Then, using the 367 software ImageJ (http://imagej.nih.gov/ij), the reference line was converted into pixels, and set as a scale. Next, the width of the ultrasound image was determined, 368 369 and the middle point marked on the image. To obtain the AT thickness, the distance 370 between the superficial and deep part of the paratenon was measured at 2, 4, and 6 371 cm from the AT's insertion. The measurements of three ultrasound images were 372 averaged for each position. Conversely, ultrasound device tools were used to 373 determine the CSA. Briefly, a discontinuous line was drawn following the internal part 374 of the paratenon as reference and the CSA was measured at 2, 4, and 6 cm of its 375 insertion. Similarly, the measurements of three ultrasound images were averaged for 376 each position.

For the stiffness measurements, we obtained approximately 4 SWE color maps (height x width, 2.5 cm x 1 cm) selected using the elastography tools. Then, using the ultrasound device tools, a line was drawn in the middle of the ultrasound

image (this line indicates the 4 cm distance from the AT). A region of interest (ROI)
of 3 mm diameter (Siu *et al.*, 2016) was aligned with the reference line and located in
the middle of the tendon, and the stiffness (kPa) was determined. Lastly, mean
stiffness was calculated over the ROIs of the 4 consecutive images (Coombes *et al.*,
2018).

385

386 HD-sEMG signal analysis

387 Motor unit analysis. The HD-sEMG signals recorded during the two 388 isometric plantarflexion contractions at each target torque (10, 40, and 70% MVC) 389 were decomposed into motor unit spike trains with an algorithm based on blind 390 source separation, which provides automatic identification of multiple single motor 391 units (Martinez-Valdes et al., 2022). Single motor units were assessed for 392 decomposition accuracy with a validated metric (Silhouette, SIL), which was set to \geq 393 0.90 (Negro et al., 2016). SIL is a normalized measure of the relative height of the 394 peaks of the decomposed spike trains with respect to the baseline noise (Martinez-395 Valdes et al., 2022).

396 The HD-sEMG signals were decomposed throughout the duration of the 397 submaximal contractions, and the discharge times of the identified motor units were 398 converted into binary spike trains (Martinez-Valdes et al., 2018). Recruitment 399 threshold was defined as the torque (%MVC) at the time when motor units began 400 firing action potentials and de-recruitment threshold was defined as the torque 401 (%MVC) at the time when motor units ceased firing action potentials (Martinez-402 Valdes et al., 2020). DR and COVisi were calculated during the steady phase of the 403 torque signal (10 s duration). Missing pulses producing non-physiological firing rates 404 i.e., inter-spike intervals > 250 ms, were manually and iteratively deleted and the 405 pulse train was re-estimated. Furthermore, if the algorithm assigned two or more 406 pulses for a single firing, we removed the additional firing, and the final pulse trains 407 were re-calculated (Martinez-Valdes et al., 2022). Motor unit data was recorded, 408 analyzed, and reported according to the consensus for experimental design in 409 electromyography: single motor unit matrix (Martinez-Valdes et al., 2023).

410

411 *Neuromechanical delay and cross-correlation coefficient.* Motor unit rate 412 coding and force generation interactions were determined using cross-correlation to

413 assess delays and similarities between fluctuations in cumulative motor unit firing of each individual muscle (MG, LG, and SO) and torque. Motor unit discharge times 414 415 obtained were summed to generate a cumulative spike train (CST) that represents 416 the cumulative activity of multiple motor units (Martinez-Valdes et al., 2022). CST and torque parameters were low pass filtered (4th order zero-phase Butterworth, 2 417 Hz) and then high pass filtered (4th order zero-phase Butterworth, 0.75 Hz) as 418 419 presented previously (Martinez-Valdes et al., 2021). Then, CST and torque were 420 cross-correlated to determine the delay in their fluctuations (calculated from the lags 421 obtained from the cross-correlation function) and the similarities in their fluctuations 422 (cross-correlation coefficient) (Martinez-Valdes et al., 2022). Delays between the 423 motor unit firing activity and torque were used as a measure of the NMD. Finally, 424 cross-correlation coefficient between signals was computed in 5-s segments with 425 50% overlap (Martinez-Valdes et al., 2022). The average NMD and cross-correlation 426 coefficient calculated from these segments was reported. Furthermore, the NMD and 427 cross-correlation coefficient of the triceps surae muscle (MG, LG, and SO combined) 428 was also calculated.

429

Torque signal analysis. Torque signals were low pass filtered at 15 Hz and then used to quantify torque steadiness (coefficient of variation of torque (COV Torque), SD torque/mean torque * 100) from the steady phase of the contractions (Martinez-Valdes *et al.*, 2020). The torque exerted by each participant was plotted using a custom-made MATLAB script allowing us to identify and select the steady phase of the contraction needed for the analysis (Arvanitidis *et al.*, 2022).

436

437 Intermuscular coherence and functional muscle network analysis

438 Intermuscular coherence analysis was performed to identify the presence of a 439 common descending drive shared by motor unit pools of the MG, LG, and SO 440 muscles. Intermuscular coherence is a well-established measure of association of motor unit firing activity between muscles (Dobie & Wilson, 1989; Simpson et al., 441 2000; Miranda de Sá & Felix, 2002). Its calculation generates a connectivity strength 442 443 value for each pair of EMG signals (in our case motor unit CSTs) within a specified 444 frequency band, resulting in a weighted adjacency matrix for each subject. 445 Significant coherence at low frequencies (<5 Hz) is associated with force production;

446 in the alpha band (5-15 Hz), with involuntary force oscillations (physiological tremor) 447 and afferent feedback loops; in the beta band (15-30 Hz), with corticospinal 448 projections; and in the gamma band (30-60 Hz), with signals relevant to efferent 449 motor commands in the motor cortex (Brown et al., 1998). Hence, these four 450 frequency bands were considered in the calculation of the connectivity matrices. 451 Furthermore, to isolate significant connections and exclude less relevant ones, a threshold was applied as has been done previously (Halliday et al., 1995). To ensure 452 453 consistency across subjects, a proportional threshold of 25% was used to maintain 454 the same connection density (Fornito et al., 2016).

After obtaining the average coherence matrix for each group, the muscular connectivity was visually represented using graph theory. In this representation, each muscle was depicted as a node within the network, while the functional connections, revealed by intermuscular coherence, were illustrated as edges between these nodes. Graph theory, widely utilized in EEG studies, allows for the analysis of networks with numerous connections and nodes, enabling the application of various metrics for comparison (Fornito *et al.*, 2016).

462 Due to the limited number of muscles accessible for data collection in this 463 study (N=3), the resulting networks have fewer connections and nodes. 464 Consequently, the analysis was centered on the assessment of the strength metric. 465 The strength metric represents the cumulative weights of links connected to a node, 466 indicating the overall strength of connections between muscles. This metric reflects 467 the muscle's influence or contribution to collective muscle activity, with higher 468 strength values indicating a more prominent role in the neural control of movement (Mheich et al., 2020). All network measures were computed using the Brain 469 470 Connectivity Toolbox, a specialized tool for analyzing functional connectivity and 471 graph-based interactions between signals (Rubinov & Sporns, 2010).

472

473 Statistical analysis

Descriptive statistics were used to report the data which are presented as mean \pm SD, unless otherwise specified. The Shapiro-Wilk test was used to determine the normality of the data, and the Levene test was used to evaluate the assumption of homogeneity of variance. As these assumptions were met, parametric statistical tests were deemed appropriate. The level of significance for all statistical procedures was set at P<0.05 and 95% confidence intervals (CI) of differences were

reported. Participant´ characteristics, questionnaires scores, morpho-mechanical
properties of the AT and connectivity strength were compared between groups using
independent T-test for continuous variables. IBM SPSS Statistics software, V. 29.0
(Armonk, New York, USA) was used for these comparisons.

484 Statistical analyses for the HD-sEMG motor unit data were done using R 485 (version 4.2.1, R Development Core Team, 2023). A generalized linear mixed-effects 486 model (GLMM) analysis was performed with the R package lme4 (version 1.1.31). 487 The following model was used "variable of interest ~ group * muscle * torque (1) 488 subject)". This model is read as a "variable of interest predicted by each of the 489 factors". The variable of interest (on the left) is the dependent variable and the ones 490 on the right (except the subject) are the independent (explanatory) variables, or 491 "fixed effects", group (Control, NIAT), muscle (MG, LG, SO), and torque (10, 40 and 492 70% MVC). The 1/subject in the model is the "random effect", which characterizes 493 the variation that is due to individual differences. The normality of residuals was 494 assessed using the Shapiro-Wilk test after running the model. Residual outliers were 495 removed using Cook's distance method (considering as threshold, 4 times the SD) 496 when the normality assumption was not met. Post-hoc pairwise comparisons were 497 conducted using Sidak correction and least-squares contrasts, as implemented in 498 the R package emmeans (version 1.8.8). The post-hoc tests were used based on the 499 interactions identified with the GLMM. Post-hoc results were reported as p-value and 500 95% confidence intervals (CI).

Torque steadiness was compared between groups at each torque level with a linear mixed model analysis with factors group (Control, NIAT) and torque (10, 40, 70% MVC) as fixed effects, and participants as random effect using GraphPad Prism software V.8.0.2 (San Diego, California, USA).

505 Relationships between pain level and questionnaire scores and 506 neuromuscular and morpho-mechanical parameters were analyzed through 507 correlation analysis using IBM SPSS Statistics software, V. 29.0. Pearson correlation 508 coefficients and p-values for each significant correlation were reported.

- 509
- 510
- 511
- 512
- 513

514 **RESULTS**

515

516 *Participants characteristics*

517 All 51 participants (26 NIAT, 25 asymptomatic controls) completed the study. 518 Participants anthropometrics, peak plantar flexion torque and information regarding 519 their symptoms are reported in Table 1. No statistical differences were observed 520 between the groups.

521

Table 1.	Participan	ts charac	teristics.
----------	------------	-----------	------------

		NIAT group	Control group	p-value	95% Cl
		(N=26)	(N=25)		
Age (years)		29.0 ± 8.5	28.6 ± 3.9	0.815	[-4.17 to 3.30]
Sex (males/female	es)	14/12	17/8		
Height (cm)		173.0 ± 9.2	171.1 ± 9.2	0.463	[-7.08 to 3.27]
Weight (kg)		75.1 ± 17.2	74.0 ± 11.6	0.793	[-9.37 to 7.20]
Peak torque (Nm)		85.7 ± 24.7	91.6 ± 30.1	0.448	[-9.60 to 21.38]
Laterality (Right/Left)		24/2	19/6		
Assessed leg (Right/Left)		16/10	17/8		
Bilateral symptom	S	13/26	N/A		
Symptoms	duration	42.10 ± 38.45	N/A		
(months)					
NRS		3.12 ± 1.24	N/A		

Values are presented as mean ± SD (Standard deviation). 95% Confidence intervals are reported. NIAT, non-insertional Achilles tendinopathy; NRS, Numerical Rating Scale; N/A, non-applicable. * Indicates significant statistical difference, p<0.05.

522

523 Questionnaires

Questionnaire scores between groups are reported in Table 2. Individuals in the NIAT group presented lower scores in the VISA-A (p<0.001), IPAQ-SF (p=0.019), FAAM (Subscale Activities of Daily Living) (p<0.001), and FAAM (Subscale Sports) (p<0.001) compared to the Control group. Additionally, individuals in the NIAT group showed higher TSK scores compared to the Control group (p=0.021). No differences were observed in the PCS scores between groups.

- 530
- 531
- 532

	NIAT group	Control group	p-value	95% CI
	(N=26)	(N=25)		
VISA-A	65.73 ± 16.16	100 ± 0	<0.001*	[23.99 to 37.35]
IPAQ-SF	128.46 ± 58.36	171.60 ± 68.19	0.019*	[7.47 to 78.81]
FAAM (Subscale Activities	85.42 ± 11.15	99.44 ± 1.83	<0.001*	[9.48 to 18.56]
of Daily Living)				
FAAM (Subscale Sports)	68.15 ± 17.04	98.68 ± 3.68	<0.001*	[23.52 to 37.53]
PCS	12.15 ± 10.65	8.72 ± 8.25	0.205	[-8.81 to 1.94]
TSK	36.08 ± 7.64	30.80 ± 8.21	0.021*	[-9.74 to -0.82]

Table 2. Questionnaires score.

Values are presented as mean ± SD (Standard deviation). 95% Confidence intervals are reported. NIAT, non-insertional Achilles tendinopathy; VISA-A, Victorian Institute of Sport Assessment – Achilles Questionnaire; IPAQ-SF, International Physical Activity Questionnaire short form; FAAM, Foot and Ankle Ability Measure, PCS, Pain Catastrophizing Scale; TSK, Tampa Scale for Kinesiophobia. * Indicates significant statistical difference, p<0.05.

533

534

535 Morpho-mechanical properties of the AT

Morpho-mechanical properties of the AT between groups are reported in Table 3. Individuals in the NIAT group presented higher thickness at 6 cm (p=0.034), and CSA at 4 cm (p=0.039) and 6 cm (p<0.001) of the AT compared to Control group. Additionally, individuals in the NIAT group showed lower stiffness of the AT compared to the Control group (p=0.003). No differences were observed in length, thickness at 2 and 4 cm and CSA at 2 cm of the AT between groups.

542

Table 3. Morpho-mechanical properties of the Achilles tendon.

	NIAT group	Control group	p-value	95% CI
	(N=26)	(N=25)		
Length (cm)	21.23 ± 3.07	19.89 ± 2.57	0.096	[-2.94 to 0.25]
Thickness 2 cm (cm)	0.35 ± 0.06	0.37 ± 0.04	0.400	[-0.02 to 0.04]
Thickness 4 cm (cm)	0.45 ± 0.10	0.41 ± 0.05	0.074	[-0.08 to 0.004]
Thickness 6 cm (cm)	0.47 ± 0.12	0.40 ± 0.05	0.008*	[-0.12 to -0.02]
CSA 2 cm (cm ²)	0.46 ± 0.08	0.44 ± 0.09	0.315	[-0.07 to -0.02]
CSA 4 cm (cm ²)	0.44 ± 0.09	0.40 ± 0.06	0.039*	[-0.09 to -0.002]
CSA 6 cm (cm ²)	0.48 ± 0.08	0.38 ± 0.08	<0.001*	[-0.14 to -0.05]
Stiffness (kPa)	68.16 ± 7.80	75.95 ± 9.98	0.003*	[2.76 to 12.82]

Values are presented as mean ± SD (Standard deviation). 95% Confidence intervals are reported. NIAT, non-insertional Achilles tendinopathy; CSA, cross-sectional area * Indicates significant

statistical difference, p<0.05.

543

544 *Motor unit decomposition*

545 A total of 5,681 motor units were identified in the triceps surae muscle during submaximal contractions in both the NIAT and control groups. At 10% MVC, the 546 average number of motor units identified in the NIAT group were 14.3 ± 9.2 for MG. 547 548 6.5 ± 5.1 for LG, and 15.6 ± 8.4 for SO; in the control group, the averages were 18.9 ± 18.6 for MG, 8 ± 10.9 for LG, and 12.6 ± 8.8 for SO. At 40% MVC, the NIAT group 549 550 had averages of 25.3 ± 19.5 for MG, 11.4 ± 9.6 for LG, and 16.5 ± 8.0 for SO, while 551 the control group had 25.5 ± 23.4 for MG, 11.1 ± 11.3 for LG, and 12.3 ± 7.5 for SO. At 70% MVC, the NIAT group had averages of 18.5 ± 15.7 for MG, 9.3 ± 11.1 for LG, 552 553 and 12.1 \pm 8.1 for SO, compared to the control group's 10.4 \pm 12.6 for MG, 3.9 \pm 2.5 554 for LG, and 11.9 ± 8.0 for SO.

555

556 *Recruitment and de-recruitment threshold*

Recruitment thresholds obtained from motor units from MG, LG, and SO 557 muscles at 10, 40, and 70% MVC are presented in Figure 3A. Overall, no 558 differences in recruitment threshold were observed between groups across torque 559 560 levels (Group effect: F=0.56, P=0.46). Within-group muscle differences at each 561 torque level can be seen in Figure 3A. De-recruitment thresholds calculated from motor units from MG, LG, and SO muscles at 10, 40, and 70% MVC are presented 562 in Figure 3B. In general, de-recruitment threshold increased differently across 563 torque levels between muscles and groups (interaction effect: F=7.24, P<0.0001), 564 with between groups differences showing decreased de-recruitment threshold (i.e., 565 566 delayed de-recruitment) in the LG in the NIAT group compared to control group at 567 70% MVC (P=0.039, 95% CI= 0.11 to 10.41). Within-group muscle differences at 568 each torque level can be seen in Figure 3B.

Α



Recruitment Threshold







571

Figure 3. A) Recruitment threshold from medial gastrocnemius (MG: red bar), lateral gastrocnemius (LG: green bar) and soleus (SO: blue bar) muscles at 10, 40, and 70% maximum voluntary contraction (MVC) between control and non-insertional tendinopathy (NIAT) groups. B) Derecruitment threshold from MG, LG, and SO muscles at 10, 40, and 70% MVC between control and non-insertional tendinopathy (NIAT) groups. A linear mixed model was used for statistical comparisons. * Statistical differences between muscles and groups, p<0.05. # Statistical differences between muscles within groups, p<0.05.

579

580 Discharge Rate

DR obtained from motor units from MG, LG, and SO muscles at 10, 40, and 70% MVC are presented in **Figure 4.** Overall, DR increased differently across torque levels between muscles and groups (interaction effect: F=7.52, P<0.0001), with between-group differences showing significantly increased DR in the LG in the NIAT group compared to control group at 70% MVC (P=0.002, 95% CI= -3.27 to -0.34). Within-group muscle differences at each torque level can be seen in **Figure 4.**

Discharge Rate

588

Figure 4. Motor unit discharge rate (DR) from medial gastrocnemius (MG: red dot), lateral gastrocnemius (LG: green square) and soleus (SO: blue triangle) muscles at 10, 40, and 70% maximum voluntary contraction (MVC) between control and non-insertional tendinopathy (NIAT) groups. A linear mixed model was used for statistical comparisons. * Statistical differences between muscles and groups, p<0.05. # Statistical differences between muscles within groups, p<0.05.</p>

595

596 Discharge Rate Variability

597 COVisi calculated from motor units from MG, LG, and SO muscles at 10, 40, 598 and 70% MVC are presented in **Figure 5.** Overall, COVisi increased differently 599 between groups and muscles (interaction effect: F=4.41, P=0.012); however, no 600 differences between NIAT and control groups were observed. Within-group muscle 601 differences at each torque level can be seen in **Figure 5.**

Coefficient of variation of the interspike interval (COVisi)

Figure 5. Motor unit coefficient of variation of the interspike interval (COVisi) from medial gastrocnemius (MG: red dot), lateral gastrocnemius (LG: green square) and soleus (SO: blue triangle) muscles at 10, 40, and 70% maximum voluntary contraction (MVC) between control and non-insertional tendinopathy (NIAT) groups. A linear mixed model was used for statistical comparisons. # Statistical differences between muscles within groups, p<0.05.</p>

608 Neuromechanical delay and cross-correlation coefficient

609 NMD calculated from motor units from MG, LG, and SO muscles at 10, 40, 610 and 70% MVC are presented in Figure 6A. Overall, NMD decreased distinctly 611 across torque levels between muscles and groups (interaction effect: F=11.74, 612 P<0.0001); however, no differences between NIAT and control groups were 613 observed. Within-group muscle differences at each torque level can be seen in 614 Figure 6A. CST and torque cross-correlation coefficient calculated from motor units 615 from MG, LG, and SO muscles at 10, 40, and 70% MVC are presented in Figure 6B. In general, cross-correlation coefficients varied across torque levels between 616 muscles and groups (interaction effect: F=84.27, P<0.0001), with between groups 617 618 differences showing significantly decreased cross-correlation coefficient in the LG in 619 the NIAT group compared to control group at 10% MVC (P<0.0001, 95% CI= 0.07 to 0.27) and increased cross-correlation coefficient in the LG in the NIAT group 620 621 compared to control group at 70% MVC (P=0.013, 95% CI= -0.21 to -0.01). Withingroup muscle differences at each torque level can be seen in Figure 6B. 622

A Neuromechanical Delay (NMD)

B Cross-correlation coefficient (CST vs Torque)

623

Figure 6. A) Neuromechanical delay (MND) from medial gastrocnemius (MG: red bar), lateral gastrocnemius (LG: green bar) and soleus (SO: blue bar) muscles at 10, 40, and 70% maximum voluntary contraction (MVC) between control and non-insertional Achilles tendinopathy (NIAT) groups.
B) Cross-correlation coefficients between cumulative spike train (CST) vs torque from MG, LG, and SO muscles at 10, 40, and 70% MVC between control and NIAT groups. A linear mixed model was used for statistical comparisons. * Statistical differences between muscles and groups, p<0.05. #

631 632

633 Torque steadiness

Overall, torque steadiness varied across torque levels (Torque effect: F=29.94 P<0.0001, MD=0.28, 95% CI= -0.04 to 0.60); however, no differences between groups were observed (Group effect: P=0.09).

- 637
- 638

640 Connectivity strength and functional muscle networks

The connectivity strength and functional muscle networks derived from 641 642 intermuscular coherence analysis for the 0-5 Hz frequency band, calculated from the 643 CSTs of the MG, LG, and SO muscles at 10% MVC, are shown in Figure 7A. 644 Individuals in the control group presented higher connectivity strength compared to 645 the NIAT group (p=0.03). Functional muscle networks (MG:M1, LG:M2, and SO:M3) 646 showed decreased connectivity between MG and LG muscles in the NIAT group 647 compared to the control group. Connectivity strength for the frequency band 5-15 Hz obtained from the CSTs of the MG, LG, and SO muscles at 10% MVC can be seen 648 649 in **Figure 7B.** Individuals in the control group showed higher connectivity strength in 650 this bandwidth compared to the NIAT group (p<0.001). Functional muscle networks 651 showed decreased connectivity between MG and LG muscles in the NIAT group 652 compared to the control group. No differences in connectivity strength for the 653 frequency bands 15-30 Hz and 30-60 Hz were observed between control and NIAT groups at 10% MVC. Additionally, no differences in connectivity strength for the 654 655 frequency bands 0-5 Hz, 5-15 Hz, 15-30 Hz, and 30-60 Hz were observed between 656 control and NIAT groups at 40 and 70% MVC (p>0.09 in all cases).

657

A Coherence 0-5 Hz at 10% MVC

659

в

Figure 7. A) Left, intermuscular coherence for the frequency band 0-5 Hz at 10% maximum voluntary contraction (MVC) between control and non-insertional Achilles tendinopathy (NIAT) groups. Right, functional muscle networks for the medial gastrocnemius (MG:M1), lateral gastrocnemius (LG:M2), and soleus (SO:M3) muscles between control and NIAT groups. B) Left, intermuscular coherence for the frequency band 5-15 Hz at 10% MVC between control and NIAT groups. Right, functional muscle networks for the MG, LG, and SO muscles between control and NIAT groups. An unpaired t-test was used for statistical comparisons. * Statistical differences between groups, p<0.05.</p>

668

669 Relationships between pain level and questionnaires and neuromuscular and 670 morpho-mechanical parameters in individuals with NIAT

671 A correlation analysis was conducted to assess the relationship between pain level and questionnaires and neuromuscular and morpho-mechanical parameters. 672 The pain level was significantly negatively correlated with LG's de-recruitment 673 threshold at 70% MVC (r = -0.52, P<0.05). VISA-A was significantly negatively 674 correlated with AT thickness 6 cm (r= -0.46, P<0.01). Moreover, FAAM (Subscale 675 676 Activities of Daily Living) was significantly negatively correlated with AT thickness 6 cm (r= -0.64, P<0.001) and CSA 6 cm (r= -0.42, P<0.05). Finally, PCS was 677 significantly positively correlated with AT thickness 6 cm (r=0.49, P<0.05). 678 679 Additionally, there were no significant correlations between the IPAQ-SF, FAAM

(Sports Subscale), and TSK questionnaires with motor unit properties or the tendon'smorpho-mechanical parameters.

682

683 **DISCUSSION**

684

685 This study demonstrates that individuals with NIAT exhibit load- and muscle-686 dependent alterations in motor unit discharge rate, motor unit firing-torque 687 relationships, and intermuscular coherence of the triceps surae during isometric 688 plantarflexion contractions. Specifically, our findings show that, at higher force levels, 689 individuals with NIAT display an increased discharge rate and a reduced de-690 recruitment threshold in the LG compared to controls. Additionally, the results reveal 691 that the cross-correlation coefficient between motor unit activity and torque 692 production (CST vs. Torque) in the LG is lower at low forces and higher at high 693 forces in individuals with NIAT, suggesting an altered contribution of the LG to the 694 overall plantarflexion torque. Moreover, our analysis indicates a reduced connectivity 695 strength within the triceps surae for low-frequency bands (0-5 and 5-15 Hz) at low 696 forces in individuals with NIAT. Notably, the functional muscle network analysis 697 highlights a decreased connectivity between the MG and LG muscles, possibly 698 reflecting an imbalance in the synergistic activity of the triceps surae in individuals 699 with NIAT.

700

Differential changes in motor unit firing properties for MG, LG, and SO muscles in
 individuals with NIAT

703 There is evidence that some muscles are predominantly controlled by a 704 common neural drive (Bremner et al., 1991; Gibbs et al., 1995; Laine et al., 2015; 705 Kerkman et al., 2018), with the strength of this shared drive being greater among 706 muscles closely related anatomically and functionally (Gibbs et al., 1995; Kerkman et 707 al., 2018). However, recent evidence indicates that the common neural drive 708 between the MG, LG, and SO muscles is reduced during isometric plantarflexion 709 contractions and heel-raise tasks (Hug et al., 2021). Our findings are in accordance 710 with those results since we observed muscle differences in recruitment and de-711 recruitment thresholds, DR, COVisi, NMD and cross-correlation coefficients at 712 different torque levels, providing further evidence of the limited shared neural drive to 713 the triceps surae muscles in both healthy individuals and those with NIAT. However,

we also observed specific motor unit firing changes for the NIAT group in the LG muscle. The NIAT group showed increased DR and decreased de-recruitment thresholds (as illustrated in Fig. 4 and Fig. 3 B, respectively) at 70% MVC, revealing a load-dependent adaptation in motor unit firing properties for this muscle, specifically.

719

720 To our knowledge, only one study has investigated the motor unit firing 721 properties of the triceps surae in individuals with NIAT. Consistent with our findings, 722 Fernandez et al. (2022) reported no significant differences in DR of the MG, LG, and 723 SO muscles between NIAT and control groups during isometric plantarflexion 724 contractions at 10% and 20% of the MVC. However, they observed that in the NIAT 725 group, the DR of the LG remained unchanged as the target torque increased from 726 10% to 20% MVC. Contrary to this, our study found no such blunted firing rate response in the LG; instead, we observed a more pronounced increase in LG's DR 727 728 at higher force levels. Additionally, we noted a reduction in the de-recruitment 729 threshold of the LG at 70% MVC. Several factors may explain these observations. 730 First, NIAT is known to alter the mechanical properties of the Achilles tendon, 731 potentially requiring greater motor unit firing to maintain the necessary torque as a 732 compensatory mechanism for the reduced tendon stiffness. This hypothesis is 733 supported by De la Fuente et al. (De la Fuente et al., 2023), who observed increased 734 activation of the MG muscle at high force levels (90% MVC) in individuals post-735 Achilles tendon rupture repair—an injury that increases tendon compliance. Indeed, changes in tendon mechanical properties can alter force transmission and 736 movement generation. However, it is crucial to consider why only the LG exhibits this 737 738 increased DR at higher forces. NIAT is also associated with increased tendon pain, 739 which may alter the plantarflexion force vector, leading to greater recruitment of the 740 LG muscle at high forces. This idea is supported by the findings of Hug et al. (Hug et 741 al., 2021), who demonstrated that the mean discharge rate of the LG can be 742 modulated by changes in foot position during plantarflexion. Therefore, it is possible 743 that NIAT-induced pain-related motor adaptations result in increased neural drive to 744 the LG at higher forces. This pain-related adaptation is further evidenced by the 745 observed reduction in de-recruitment threshold at high forces. Martinez-Valdes et al. 746 (Martinez-Valdes et al., 2020) previously showed that experimental muscle pain 747 reduces the recruitment and de-recruitment thresholds of higher-threshold motor

748 units, increasing the duration of their activity during contraction. These higher-749 threshold motor units are associated with greater force production but have reduced 750 resistance to muscle fatigue. Consequently, the reduced de-recruitment threshold observed in our study might suggest that the increased activation of the LG during 751 752 the task could lead to inefficient recruitment within the triceps surae group. Typically, 753 the LG exhibits a higher recruitment threshold compared to the MG and SO during plantarflexion (Hug et al., 2021). Thus, prolonged overactivation of the LG could 754 755 potentially increase fatigue in the triceps surae due to altered motor strategies in 756 response to pain, which may exacerbate symptoms over time. Future studies should 757 investigate potential associations between pain and fatigue in individuals with NIAT 758 to further understand these adaptations.

759

760 NMD and CST-torque cross-correlation coefficients

761 The conversion of neural signals to force output has a latency caused by the 762 dynamic responsiveness of motor neurons and the time required to stretch the series 763 elastic components of the muscle-tendon unit after the activation of the muscle fibers 764 (Del Vecchio et al., 2018). In this context, the NMD represents the time difference 765 between the neural drive and the generated torgue during a voluntary contraction 766 and can be estimated from the time lag of the peak of the cross-correlation between the CST and torque (Del Vecchio et al., 2018). Our results showed no differences in 767 768 NMD of the MG, LG, and SO between the NIAT group and control group at 10%, 769 40%, and 70% MVC; however, NMD decreased as the target torque increased in both groups. The decrease in the NMD as the target torque increase may be 770 explained by the Henneman's size principle (Henneman, 1957), since as the target 771 772 torque increases larger motor units, which innervate fast-twitch muscle fibers, are 773 recruited. Consequently, the recruitment of fast-twitch muscle fibers may contribute 774 to the observed decrease in NMD. Theoretically, the stiffness of the AT affects the 775 time to stretch the elastic components of the muscle-tendon unit during a voluntary contraction. Given our observation of reduced tendon stiffness in individuals with 776 777 NIAT, we anticipated that this difference might influence NMD results. However, no 778 significant differences between the NIAT and control groups were observed. In 779 contrast, a previous study by Chang and Kulig (Chang & Kulig, 2015) reported an 780 increase in electromechanical delay (quantified as the time difference between the 781 onset of force/torque and muscle activation during electrically induced contraction) in

782 individuals with NIAT. These conflicting findings may be partially explained by 783 differences in how NMD and electromechanical delay are calculated. 784 Electromechanical delay is typically estimated at a single time point when the muscle activates from a passive state (Chang & Kulig, 2015). In contrast, NMD is influenced 785 786 by active motor unit responses, such as twitch duration and afferent inputs, during 787 contraction and is measured when the tendon is under sustained tension, reflecting a state of greater tendon stiffness. It is possible that delays between motor unit/EMG 788 789 activity and torque are more pronounced in NIAT during the initial phase of the force-790 tension curve, particularly at the contraction intensities generated by electrically 791 induced contractions. However, testing this hypothesis would require a simultaneous 792 assessment of motor unit firing properties and tendon stiffness changes across the 793 full force-tension curve of the tendon.

794

795 Recent research has established that CST can be used to estimate the 796 effective neural drive received by a group of synergistic muscles in humans (Mazzo 797 et al., 2021). Mazzo and colleagues demonstrated that the neural drive to the triceps 798 surae muscle group (via CST analysis) is correlated with fluctuations in isometric 799 plantarflexion torque (Mazzo et al., 2022). Therefore, assessing CST-torque 800 relationships for individual heads of the triceps surae can provide insight into their 801 independent contributions to the resultant plantarflexion torque. Our results indicate 802 that the cross-correlation coefficient between CST and torque decreased in the 803 lateral gastrocnemius (LG) in the NIAT group compared to the control group at 10% 804 MVC and increased in the NIAT group at 70% MVC. These findings suggest a 805 reduced contribution of the LG to net torque at low forces in individuals with NIAT, 806 consistent with previous research that indirectly showed a significant reduction in 807 LG's contribution to plantarflexion torque at low forces in this population. However, 808 the same study also reported a decreased contribution of the LG at intermediate 809 forces, a difference we did not observe. This discrepancy may be partly due to 810 methodological differences, as Crouzier et al. (2020) used LG activation and relative 811 physiological cross-sectional area to indirectly estimate the muscle's contribution to 812 net plantarflexion torque. Since surface EMG can be influenced by muscle 813 properties, any change in LG morphology induced by NIAT can alter sEMG 814 calculations and therefore affect individual muscle force estimations, particularly at 815 higher forces due to amplitude cancellation (Dideriksen & Farina, 2019).

816 Furthermore, Crouzier et al. did not assess force levels as high as the ones used in 817 the present study. The observed decrease in LG's CST-torque cross-correlation at 818 low forces and the increase in cross-correlation at higher forces is noteworthy. As 819 previously discussed, these different force levels exert varying degrees of tension on 820 the tendon. The tension produced at 10% MVC likely demands greater muscular 821 control due to reduced tendon stiffness, which may necessitate a more balanced 822 contribution from all triceps surae muscle heads to the resultant torque. Indeed, at 823 this lower force level, despite significant muscle differences, the control group 824 exhibits more consistent CST-torque cross-correlation coefficients compared to the 825 NIAT group. Interestingly, this pattern reverses at higher forces, where the NIAT 826 group demonstrates a greater contribution of LG to the resultant torque compared to 827 the control group. Given that both MG and SO have larger CSAs and contribute 828 more significantly to the resultant plantarflexion torque (Maganaris, 2003) compared 829 to LG, it would be reasonable to expect that individuals in the control group 830 predominantly rely on these muscles to increase plantarflexion torque. However, this 831 pattern was not observed in the NIAT group. As mentioned in the discussion of the 832 DR results, it is plausible that the NIAT group employed different plantarflexion 833 strategies at various force levels in response to pain. Since tendinopathic pain 834 typically intensifies with increased load (Magnusson et al., 2010), the NIAT group 835 may have adjusted the plantarflexion torque vector to reduce the engagement of the 836 MG and SO muscles, thereby attempting to minimize the load on a potentially painful 837 tendon region. However, this hypothesis should be validated in future studies that 838 assess both overall plantarflexion torque and plantar pressure distribution.

- 839
- 840
- 841

842 Strength of synergistic activity in individuals with NIAT

When the CNS initiates a voluntary movement, it triggers the activation of several muscles through the simultaneous activation and coordination of thousands of motor units (Bizzi & Cheung, 2013). It has been proposed that the CNS simplifies this task by controlling groups of muscles, known as muscle synergies, instead of commanding each muscle independently (Singh *et al.*, 2018). Thus, the CNS encodes several muscle synergies, and it combines them according to a specific task, generating the muscle contractions necessary for the desired movement

850 (Alessandro et al., 2013). Intermuscular coherence analysis is a well-established 851 method for identifying shared neural drive, as it quantifies the frequency spectrum of 852 correlated activity between motor units from different muscle pairs (Laine et al., 853 2021). For this reason, we employed intermuscular coherence together with network 854 analysis, to assess the strength of the neural connectivity (shared neural drive) 855 between the three triceps surae muscle heads. Our network analysis demonstrated a 856 decrease in the connectivity strength within the 0-5 Hz band (delta band) for the triceps surae in individuals with NIAT at 10% MVC. This decrease in connectivity 857 858 strength was primarily given by a reduction in intermuscular coherence between MG 859 and LG muscles. Motor unit coherence at frequencies below 5 Hz reflects concurrent 860 fluctuations in motor unit firing rates (common neural drive) between muscles (De 861 Luca et al., 1982; Myers et al., 2004) and plays a crucial role in controlling muscle 862 force (Farina & Negro, 2015). These findings, coupled with the reductions in LG's 863 CST-torque cross-correlation estimates, further support the evidence of reduced LG 864 involvement in generating plantarflexion torgue at low forces in individuals with NIAT. 865

866 Additionally, our findings showed a decrease in the connectivity strength 867 within the frequency band 5-15 Hz (alpha band) for the triceps surae at 10% MVC in 868 individuals with NIAT. Again, observing a decrease of the intermuscular coherence 869 between MG and LG muscles. Since motor unit coherence at frequencies between 870 5-15 Hz has been associated with involuntary force oscillations (physiological tremor) (Laine et al., 2015), it can be hypothesized that these oscillations might be 871 872 transmitted to the Achilles tendon and subsequently influence torque steadiness. 873 However, our findings did not reveal any differences in torque steadiness between 874 NIAT and control groups across torque levels. Considering that increased alpha-875 band intermuscular coherence is typically linked to tasks demanding a high level of 876 muscle coordination (Nazarpour et al., 2012; de Vries et al., 2016; Laine & Valero-877 Cuevas, 2017; Reyes et al., 2017), and given that the task in our study required 878 precise matching of plantarflexion torque to visual feedback, it is plausible to 879 speculate that individuals with NIAT may experience altered coordination between 880 the MG and LG muscles during low-force contractions, potentially due to reduced 881 afferent input. Indeed, literature suggests that coordinated patterns of this frequency 882 band may reflect afferent feedback loops (arising from 1a or 1b fibers), which is 883 supported by observations that cycles of excitation around the monosynaptic stretch

reflex can induce oscillations in the 5-15 Hz frequency band within individual muscles (Lippold, 1970; Christakos *et al.*, 2006; Erimaki & Christakos, 2008). Additionally, it has been proposed that the oscillations in the 5-15 Hz frequency band may be linked to the activity of the cerebello-thalamo-cortical-circuit, which resonates at these frequencies and synchronizes with motor output via its projections to the cortex and/or pontomedullary reticular formation (Gross *et al.*, 2002; Soteropoulos & Baker, 2006; Williams *et al.*, 2010).

891

Relationships between pain level and questionnaires scores and neuromuscular and
morpho-mechanical parameters

894 Our results showed a negative correlation between pain level and LG de-895 recruitment threshold at 70% MVC, suggesting that as the pain level increases, 896 individuals with NIAT decrease the de-recruitment threshold of the LG at high forces, 897 allowing them to maintain the LG active for a greater period to accomplish the task. 898 Additionally, we found a negative correlation between VISA-A and AT thickness 6 899 cm, indicating that as the tendon thickness increases, the severity of the symptoms 900 increases. Nevertheless, previous studies have shown inconsistent relationships 901 between the extent and/or severity of tendon abnormalities and severity of symptoms 902 (Martin *et al.*, 2018). Similarly, we observed a negative correlation between FAAM 903 (Subscale Activities of Daily Living) and AT thickness 6 cm and CSA 6 cm, showing 904 that as the tendon thickness and cross-sectional area increase, the physical function 905 of individuals with NIAT decreases, providing additional evidence of the relationship 906 between tendon morphology and physical function. Finally, we found a positive 907 correlation between PCS and AT thickness at 6 cm, revealing that as the tendon 908 pain catastrophizing (rumination, magnification, and thickness increases, 909 helplessness about pain) increases. Since morphological changes to the AT in 910 individuals with NIAT are commonly linked to the duration of symptoms, our results 911 support the idea that chronic NIAT is linked to changes in psychological attitudes 912 about pain. Understanding the interplay between symptoms, functionality, 913 psychological attitudes towards pain and the neuromuscular and morpho-mechanical 914 properties is essential for an enhanced comprehension of the different factors 915 involved in NIAT. Designing intervention strategies that consider the dynamic 916 interplay between these components is imperative to achieve optimal outcomes in 917 the management of this condition.

918 Limitations and future developments

919 There are some methodological aspects of this study which should be 920 considered.

921 First, tendon stiffness measurements were conducted under resting conditions. 922 Unfortunately, assessing tendon stiffness with SWE during plantarflexion 923 contractions presents significant challenges, mainly due to foot movement within the 924 isokinetic ankle attachment. This movement introduces artifacts in the elastography 925 color map, which compromises the accuracy of stiffness estimations. Additionally, 926 current SWE devices have very low sampling resolution (e.g., up to 2 SWE images 927 per second) and do not reliably estimate tendon stiffness during contractions (Khair 928 et al., 2024). Future studies should utilize more advanced SWE devices to 929 simultaneously assess AT stiffness and triceps surae motor unit firing rates during 930 plantarflexion contractions, offering a more comprehensive understanding of the 931 interplay between neural coding and muscle-tendon behavior in this condition. 932 Furthermore, given that recent studies have successfully identified motor units during 933 both shortening and lengthening contractions (Glaser & Holobar, 2019; Oliveira & 934 Negro, 2021), future research should explore triceps surae motor unit firing 935 parameters during dynamic plantarflexion contractions in individuals with NIAT.

936

937 CONCLUSION

938

939 This study demonstrates that individuals with NIAT exhibit load- and muscle-940 dependent changes in triceps surae motor unit discharge rate, motor unit firing-941 torque relationships, and intermuscular coherence during isometric plantarflexion 942 contractions. Our research found increased DR and decreased de-recruitment 943 thresholds in the LG at high forces, indicating an altered neural drive to the triceps 944 surae in individuals with NIAT. Additionally, the results revealed a decreased cross-945 correlation coefficient (CST vs. Torque) in the LG at low forces, with an increase at 946 high forces, suggesting a shift in the LG's contribution to net plantarflexion torque. 947 Furthermore, our findings indicated reduced strength connectivity of the triceps surae 948 in low-frequency bands at low forces, pointing to a more unbalanced synergistic 949 activity within the triceps surae in individuals with NIAT. Collectively, these results 950 support the hypothesis that the LG may play a central role in the pathophysiology of 951 this condition, potentially affecting load transmission to the Achilles tendon.

952 953

954 ADDITIONAL INFORMATION

955

956 **ACKNOWLEDGEMENTS**

957 We thank Joeri van Helden for his assistance during data collection.

958 All figures were created using BioRender.

959

960 DATA AVAILABILITY STATEMENT

The data and analysis codes are available from the corresponding author, EM-V, upon reasonable request.

963

964 **GRANTS**

This work was supported by ANID PhD Scholarship awarded by the Government of Chile. Recipient: Ignacio Contreras-Hernandez, Scholarship ID number: 72200295

Francesco Negro was funded by the European Research Council Consolidator Grant INCEPTION (contract no. 101045605)

Eduardo Martinez-Valdes is supported by an Orthopaedic Research UK Early
 Career Research Fellowship (ORUK ref-574)

972

973

974 DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

976

977 AUTHOR CONTRIBUTIONS

IC-H, EM-V, and DF conceived and designed research; IC-H and MA
performed experiments; IC-H, DJ-G, and EM-V analysed data; IC-H and EM-V
interpreted results of experiments; IC-H prepared figures; IC-H and EM-V drafted
manuscript; IC-H, EM-V, DF, FN, DJ-G and MA edited and revised manuscript; IC-H,
EM-V, DF, FN, DJ-G and MA approved final version of manuscript.

983

984

986	
987	
988	References
989	
990 991	Alessandro C, Delis I, Nori F, Panzeri S & Berret B. (2013). Muscle synergies in neuroscience and robotics: from input-space to task-space perspectives. <i>Front Comput Neurosci</i> 7, 43.
992 993 994 995	Alghadir AH, Anwer S, Iqbal A & Iqbal ZA. (2018). Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. <i>J Pain Res</i> 11, 851-856.
996 997 998	Arndt AN, Komi PV, Brüggemann GP & Lukkariniemi J. (1998). Individual muscle contributions to the in vivo achilles tendon force. <i>Clin Biomech (Bristol, Avon)</i> 13, 532-541.
999 1000 1001 1002	Arvanitidis M, Falla D & Martinez-Valdes E. (2019). Can visual feedback on upper trapezius high- density surface electromyography increase time to task failure of an endurance task? <i>J</i> <i>Electromyogr Kinesiol</i> 49, 102361.
1003 1004 1005 1006	Arvanitidis M, Jiménez-Grande D, Haouidji-Javaux N, Falla D & Martinez-Valdes E. (2022). People with chronic low back pain display spatial alterations in high-density surface EMG-torque oscillations. <i>Sci Rep</i> 12, 15178.
1007 1008 1009	Arya S & Kulig K. (2010). Tendinopathy alters mechanical and material properties of the Achilles tendon. <i>J Appl Physiol (1985)</i> 108, 670-675.
1010 1011 1012	Bahari M, Taslimipour S & Moazenzadeh A. (2023). Different clinical presentations of Achilles tendinopathy: a cross-sectional study. <i>Rheumatol Int</i> .
1013 1014 1015 1016	Beyer R, Kongsgaard M, Hougs Kjær B, Øhlenschlæger T, Kjær M & Magnusson SP. (2015). Heavy Slow Resistance Versus Eccentric Training as Treatment for Achilles Tendinopathy: A Randomized Controlled Trial. <i>Am J Sports Med</i> 43, 1704-1711.
1017 1018	Bizzi E & Cheung VC. (2013). The neural origin of muscle synergies. <i>Front Comput Neurosci</i> 7, 51.
1019 1020 1021	Bizzi E, D'Avella A, Saltiel P & Tresch M. (2002). Modular organization of spinal motor systems. <i>Neuroscientist</i> 8, 437-442.
1022 1023	Bleakney RR & White LM. (2005). Imaging of the Achilles tendon. <i>Foot Ankle Clin</i> 10, 239-254.
1024 1025 1026	Bojsen-Møller J & Magnusson SP. (2015). Heterogeneous Loading of the Human Achilles Tendon In Vivo. <i>Exerc Sport Sci Rev</i> 43, 190-197.
1027	

1028 1029	Bremner FD, Baker JR & Stephens JA. (1991). Correlation between the discharges of motor units recorded from the same and from different finger muscles in man. <i>J Physiol</i> 432, 355-380.
1030 1031 1032	Brown P, Salenius S, Rothwell JC & Hari R. (1998). Cortical correlate of the Piper rhythm in humans. <i>J</i> Neurophysiol 80, 2911-2917.
1033 1034 1035	Chang YJ & Kulig K. (2015). The neuromechanical adaptations to Achilles tendinosis. <i>J Physiol</i> 593, 3373-3387.
1036 1037 1038	Chapman JP, Chapman LJ & Allen JJ. (1987). The measurement of foot preference. <i>Neuropsychologia</i> 25, 579-584.
1039 1040 1041 1042	Child S, Bryant AL, Clark RA & Crossley KM. (2010). Mechanical properties of the achilles tendon aponeurosis are altered in athletes with achilles tendinopathy. <i>Am J Sports Med</i> 38, 1885- 1893.
1043 1044 1045 1046	Christakos CN, Papadimitriou NA & Erimaki S. (2006). Parallel neuronal mechanisms underlying physiological force tremor in steady muscle contractions of humans. <i>J Neurophysiol</i> 95, 53- 66.
1047 1048 1049	Clark WH & Franz JR. (2018). Do triceps surae muscle dynamics govern non-uniform Achilles tendon deformations? <i>PeerJ 6,</i> e5182.
1050 1051 1052 1053	Cook JL & Purdam CR. (2009). Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. <i>British Journal of Sports Medicine</i> 43, 409-416.
1054 1055 1056 1057	Coombes BK, Tucker K, Vicenzino B, Vuvan V, Mellor R, Heales L, Nordez A & Hug F. (2018). Achilles and patellar tendinopathy display opposite changes in elastic properties: A shear wave elastography study. <i>Scand J Med Sci Sports 28,</i> 1201-1208.
1058 1059 1060 1061	Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, Pratt M, Ekelund U, Yngve A, Sallis JF & Oja P. (2003). International physical activity questionnaire: 12-country reliability and validity. <i>Med Sci Sports Exerc</i> 35, 1381-1395.
1062 1063 1064 1065	Crouzier M, Tucker K, Lacourpaille L, Doguet V, Fayet G, Dauty M & Hug F. (2020). Force-sharing within the Triceps Surae: An Achilles Heel in Achilles Tendinopathy. <i>Med Sci Sports Exerc</i> 52, 1076-1087.
1066 1067 1068	d'Avella A & Bizzi E. (2005). Shared and specific muscle synergies in natural motor behaviors. <i>Proc</i> Natl Acad Sci U S A 102, 3076-3081.

1070 1071 1072	de Jonge S, van den Berg C, de Vos RJ, van der Heide HJ, Weir A, Verhaar JA, Bierma-Zeinstra SM & Tol JL. (2011). Incidence of midportion Achilles tendinopathy in the general population. <i>Br J</i> <i>Sports Med</i> 45, 1026-1028.
1073 1074 1075 1076	De la Fuente C, Martinez-Valdes E, da Rocha ES, Geremia JM, Vaz MA & Carpes FP. (2023). Distal overactivation of gastrocnemius medialis in persistent plantarflexion weakness following Achilles tendon repair. <i>J Biomech</i> 148, 111459.
1077 1078 1079	De Luca CJ, LeFever RS, McCue MP & Xenakis AP. (1982). Control scheme governing concurrently active human motor units during voluntary contractions. <i>J Physiol</i> 329, 129-142.
1080 1081 1082	de Vries IE, Daffertshofer A, Stegeman DF & Boonstra TW. (2016). Functional connectivity in the neuromuscular system underlying bimanual coordination. <i>J Neurophysiol</i> 116, 2576-2585.
1083 1084 1085 1086	Debenham JR, Gibson WI, Travers MJ, Campbell AC & Allison GT. (2017). Modulation of Stretch- Shortening-Cycle Behavior With Eccentric Loading of Triceps Surae: A Possible Therapeutic Mechanism. <i>Journal of Sport Rehabilitation</i> 26, 151-158.
1087 1088 1089 1090	Del Vecchio A, Úbeda A, Sartori M, Azorín JM, Felici F & Farina D. (2018). Central nervous system modulates the neuromechanical delay in a broad range for the control of muscle force. <i>J</i> <i>Appl Physiol (1985)</i> 125, 1404-1410.
1091 1092 1093 1094	Dideriksen JL & Farina D. (2019). Amplitude cancellation influences the association between frequency components in the neural drive to muscle and the rectified EMG signal. <i>PLoS</i> <i>Comput Biol</i> 15, e1006985.
1095 1096 1097	Dobie RA & Wilson MJ. (1989). Analysis of auditory evoked potentials by magnitude-squared coherence. <i>Ear Hear</i> 10, 2-13.
1098 1099 1100 1101	Docking SI & Cook J. (2016). Pathological tendons maintain sufficient aligned fibrillar structure on ultrasound tissue characterization (UTC). <i>Scandinavian Journal of Medicine & Science in</i> <i>Sports 26, 675-683.</i>
1102 1103 1104	Enoka RM & Duchateau J. (2017). Rate Coding and the Control of Muscle Force. <i>Cold Spring Harb</i> Perspect Med 7 .
1105 1106 1107 1108	Erimaki S & Christakos CN. (2008). Coherent motor unit rhythms in the 6-10 Hz range during time- varying voluntary muscle contractions: neural mechanism and relation to rhythmical motor control. <i>J Neurophysiol</i> 99, 473-483.
1109 1110 1111	Farina D & Negro F. (2015). Common synaptic input to motor neurons, motor unit synchronization, and force control. <i>Exerc Sport Sci Rev</i> 43, 23-33.
1112	

1113 1114	Farina D, Negro F & Dideriksen JL. (2014). The effective neural drive to muscles is the common synaptic input to motor neurons. <i>J Physiol</i> 592, 3427-3441.
1115 1116 1117 1118	Fernandes GL, Orssatto LBR, Sakugawa RL & Trajano GS. (2023). Lower motor unit discharge rates in gastrocnemius lateralis, but not in gastrocnemius medialis or soleus, in runners with Achilles tendinopathy: a pilot study. <i>Eur J Appl Physiol</i> 123, 633-643.
1119 1120	Fornito A, Zalesky A & Bullmore E. (2016). <i>Fundamentals of brain network analysis</i> . Academic Press.
1121 1122 1123	Full RJ & Koditschek DE. (1999). Templates and anchors: neuromechanical hypotheses of legged locomotion on land. <i>J Exp Biol</i> 202, 3325-3332.
1124 1125 1126	Gibbs J, Harrison LM & Stephens JA. (1995). Organization of inputs to motoneurone pools in man. <i>J</i> <i>Physiol</i> 485 (Pt 1), 245-256.
1127 1128 1129	Glaser V & Holobar A. (2019). Motor Unit Identification From High-Density Surface Electromyograms in Repeated Dynamic Muscle Contractions. <i>IEEE Trans Neural Syst Rehabil Eng</i> 27, 66-75.
1130 1131 1132	Grigg NL, Wearing SC & Smeathers JE. (2012). Achilles tendinopathy has an aberrant strain response to eccentric exercise. <i>Med Sci Sports Exerc</i> 44, 12-17.
1133 1134 1135	Gross J, Timmermann L, Kujala J, Dirks M, Schmitz F, Salmelin R & Schnitzler A. (2002). The neural basis of intermittent motor control in humans. <i>Proc Natl Acad Sci U S A</i> 99, 2299-2302.
1136 1137 1138 1139 1140	Halliday DM, Rosenberg JR, Amjad AM, Breeze P, Conway BA & Farmer SF. (1995). A framework for the analysis of mixed time series/point process datatheory and application to the study of physiological tremor, single motor unit discharges and electromyograms. <i>Prog Biophys Mol</i> <i>Biol</i> 64, 237-278.
1141 1142 1143 1144	Handsfield GG, Inouye JM, Slane LC, Thelen DG, Miller GW & Blemker SS. (2017). A 3D model of the Achilles tendon to determine the mechanisms underlying nonuniform tendon displacements. <i>J Biomech</i> 51, 17-25.
1145 1146 1147 1148	Hansen W, Shim VB, Obst S, Lloyd DG, Newsham-West R & Barrett RS. (2017). Achilles tendon stress is more sensitive to subject-specific geometry than subject-specific material properties: A finite element analysis. <i>Journal of Biomechanics</i> 56, 26-31.
1149 1150 1151	Henneman E. (1957). Relation between size of neurons and their susceptibility to discharge. <i>Science</i> 126, 1345-1347.
1152 1153 1154 1155	Héroux ME, Dakin CJ, Luu BL, Inglis JT & Blouin JS. (2014). Absence of lateral gastrocnemius activity and differential motor unit behavior in soleus and medial gastrocnemius during standing balance. <i>J Appl Physiol (1985)</i> 116, 140-148.
1156	

1157 1158 1159	Hug F, Del Vecchio A, Avrillon S, Farina D & Tucker K. (2021). Muscles from the same muscle group do not necessarily share common drive: evidence from the human triceps surae. <i>J Appl Physiol (1985)</i> 130, 342-354.
1160 1161 1162	Hug F & Tucker K. (2017). Muscle Coordination and the Development of Musculoskeletal Disorders. <i>Exerc Sport Sci Rev</i> 45, 201-208.
1163 1164 1165 1166	Iversen JV, Bartels EM & Langberg H. (2012). The victorian institute of sports assessment - achilles questionnaire (visa-a) - a reliable tool for measuring achilles tendinopathy. <i>Int J Sports Phys</i> <i>Ther</i> 7, 76-84.
1167 1168 1169	Järvinen TA, Kannus P, Maffulli N & Khan KM. (2005). Achilles tendon disorders: etiology and epidemiology. <i>Foot Ankle Clin</i> 10, 255-266.
1170 1171 1172 1173	Kerkman JN, Daffertshofer A, Gollo LL, Breakspear M & Boonstra TW. (2018). Network structure of the human musculoskeletal system shapes neural interactions on multiple time scales. <i>Sci</i> <i>Adv</i> 4, eaat0497.
1174 1175 1176	Khair RaM, Sukanen M & Finni T. (2024). Achilles Tendon Stiffness: Influence of Measurement Methodology. <i>Ultrasound in Medicine & Biolog</i> y.
1177 1178 1179	Kongsgaard M, Nielsen CH, Hegnsvad S, Aagaard P & Magnusson SP. (2011). Mechanical properties of the human Achilles tendon, in vivo. <i>Clin Biomech (Bristol, Avon</i>) 26, 772-777.
1180 1181	Kori SH. (1990). Kinesiophobia : a new view of chronic pain behavior. <i>Pain Manage</i> 3, 35-43.
1182 1183	Kvist M. (1994). Achilles tendon injuries in athletes. Sports Med 18, 173-201.
1184 1185 1186	Laine CM, Cohn BA & Valero-Cuevas FJ. (2021). Temporal control of muscle synergies is linked with alpha-band neural drive. <i>J Physiol</i> 599, 3385-3402.
1187 1188 1189	Laine CM, Martinez-Valdes E, Falla D, Mayer F & Farina D. (2015). Motor Neuron Pools of Synergistic Thigh Muscles Share Most of Their Synaptic Input. <i>J Neurosci</i> 35, 12207-12216.
1190 1191 1192	Laine CM & Valero-Cuevas FJ. (2017). Intermuscular coherence reflects functional coordination. <i>J</i> Neurophysiol 118, 1775-1783.
1193 1194	Leadbetter WB. (1992). Cell-matrix response in tendon injury. <i>Clin Sports Med</i> 11, 533-578.
1195 1196 1197 1198	Lindemann I, Coombes BK, Tucker K, Hug F & Dick TJM. (2020). Age-related differences in gastrocnemii muscles and Achilles tendon mechanical properties in vivo. <i>J Biomech</i> 112, 110067.
1199	

1200 1201	Lippold OC. (1970). Oscillation in the stretch reflex arc and the origin of the rhythmical, 8-12 C-S component of physiological tremor. <i>J Physiol</i> 206, 359-382.
1202 1203 1204	Loeb GE, Brown IE & Cheng EJ. (1999). A hierarchical foundation for models of sensorimotor control. <i>Exp Brain Res</i> 126, 1-18.
1205 1206 1207	Maffulli N, Khan KM & Puddu G. (1998). Overuse tendon conditions: time to change a confusing terminology. <i>Arthroscopy</i> 14, 840-843.
1208 1209 1210	Maffulli N, Longo UG, Kadakia A & Spiezia F. (2020). Achilles tendinopathy. <i>Foot Ankle Surg</i> 26, 240- 249.
1211 1212 1213	Maganaris CN. (2003). Force-length characteristics of the in vivo human gastrocnemius muscle. <i>Clin</i> Anat 16, 215-223.
1214 1215 1216	Magnusson SP, Langberg H & Kjaer M. (2010). The pathogenesis of tendinopathy: balancing the response to loading. <i>Nat Rev Rheumatol</i> 6, 262-268.
1217 1218 1219 1220	Martin RL, Chimenti R, Cuddeford T, Houck J, Matheson JW, McDonough CM, Paulseth S, Wukich DK & Carcia CR. (2018). Achilles Pain, Stiffness, and Muscle Power Deficits: Midportion Achilles Tendinopathy Revision 2018. <i>Journal of Orthopaedic & Sports Physical Therapy</i> 48, A1-A38.
1221 1222 1223	Martin RL, Irrgang JJ, Burdett RG, Conti SF & Van Swearingen JM. (2005). Evidence of validity for the Foot and Ankle Ability Measure (FAAM). <i>Foot Ankle Int 26,</i> 968-983.
1224 1225 1226 1227 1228 1229	Martinez-Valdes E, Enoka RM, Holobar A, McGill K, Farina D, Besomi M, Hug F, Falla D, Carson RG, Clancy EA, Disselhorst-Klug C, van Dieën JH, Tucker K, Gandevia S, Lowery M, Søgaard K, Besier T, Merletti R, Kiernan MC, Rothwell JC, Perreault E & Hodges PW. (2023). Consensus for experimental design in electromyography (CEDE) project: Single motor unit matrix. <i>J</i> <i>Electromyogr Kinesiol</i> 68, 102726.
1230 1231 1232 1233	Martinez-Valdes E, Negro F, Arvanitidis M, Farina D & Falla D. (2021). Pain-induced changes in motor unit discharge depend on recruitment threshold and contraction speed. <i>J Appl Physiol (1985)</i> 131, 1260-1271.
1234 1235 1236 1237	Martinez-Valdes E, Negro F, Botter A, Pincheira PA, Cerone GL, Falla D, Lichtwark GA & Cresswell AG. (2022). Modulations in motor unit discharge are related to changes in fascicle length during isometric contractions. <i>J Appl Physiol (1985)</i> 133, 1136-1148.
1238 1239 1240 1241	Martinez-Valdes E, Negro F, Falla D, De Nunzio AM & Farina D. (2018). Surface electromyographic amplitude does not identify differences in neural drive to synergistic muscles. <i>J Appl Physiol</i> <i>(1985)</i> 124, 1071-1079.
1242	

1243 1244	Martinez-Valdes E, Negro F, Farina D & Falla D. (2020). Divergent response of low- versus high- threshold motor units to experimental muscle pain. <i>J Physiol</i> 598, 2093-2108.
1245 1246 1247 1248	Mazzo MR, Holobar A & Enoka RM. (2022). Association between effective neural drive to the triceps surae and fluctuations in plantar-flexion torque during submaximal isometric contractions. <i>Exp Physiol</i> 107, 489-507.
1249 1250 1251 1252	Mazzo MR, Weinman LE, Giustino V, McLagan B, Maldonado J & Enoka RM. (2021). Changes in neural drive to calf muscles during steady submaximal contractions after repeated static stretches. <i>J Physiol</i> 599, 4321-4336.
1253 1254 1255	Mheich A, Wendling F & Hassan M. (2020). Brain network similarity: methods and applications. <i>Netw</i> <i>Neurosci</i> 4, 507-527.
1256 1257 1258 1259	Miranda de Sá AM & Felix LB. (2002). Improving the detection of evoked responses to periodic stimulation by using multiple coherenceapplication to EEG during photic stimulation. <i>Med</i> <i>Eng Phys</i> 24, 245-252.
1260 1261 1262	Myers 니, Erim Z & Lowery MM. (2004). Time and frequency domain methods for quantifying common modulation of motor unit firing patterns. <i>J Neuroeng Rehabil</i> 1, 2.
1263 1264 1265	Nazarpour K, Barnard A & Jackson A. (2012). Flexible cortical control of task-specific muscle synergies. <i>J Neurosci</i> 32, 12349-12360.
1266 1267 1268 1269	Negro F, Muceli S, Castronovo AM, Holobar A & Farina D. (2016). Multi-channel intramuscular and surface EMG decomposition by convolutive blind source separation. <i>J Neural Eng</i> 13, 026027.
1270 1271 1272	O'Neill S, Barry S & Watson P. (2019). Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. <i>Physical Therapy in Sport</i> 37, 69-76.
1273 1274 1275	O'Neill S, Watson PJ & Barry S. (2015). WHY ARE ECCENTRIC EXERCISES EFFECTIVE FOR ACHILLES TENDINOPATHY? <i>Int J Sports Phys Ther</i> 10, 552-562.
1276 1277 1278	Oliveira AS & Negro F. (2021). Neural control of matched motor units during muscle shortening and lengthening at increasing velocities. <i>J Appl Physiol (1985)</i> 130, 1798-1813.
1279 1280 1281	Reyes A, Laine CM, Kutch JJ & Valero-Cuevas FJ. (2017). Beta Band Corticomuscular Drive Reflects Muscle Coordination Strategies. <i>Front Comput Neurosci</i> 11, 17.
1282 1283 1284 1285	Robinson JM, Cook JL, Purdam C, Visentini PJ, Ross J, Maffulli N, Taunton JE & Khan KM. (2001). The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. <i>British Journal of Sports Medicine</i> 35, 335-341.
1286	

1287 1288 1289	Rompe JD, Furia J & Maffulli N. (2009). Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. <i>Am J Sports Med</i> 37, 463-470.	
1290 1291 1292	Rubinov M & Sporns O. (2010). Complex network measures of brain connectivity: uses and interpretations. <i>Neuroimage</i> 52, 1059-1069.	
1293 1294 1295 1296	Sara LK, Gutsch SB, Hoeger Bement M & Hunter SK. (2023). Plantar Flexor Weakness and Pain Sensitivity Cannot Be Assumed in Midportion Achilles Tendinopathy. <i>Exercise, Sport, and</i> <i>Movement</i> 1, 1-7.	
1297 1298 1299 1300	Shalabi A, Kristoffersen-Wilberg M, Svensson L, Aspelin P & Movin T. (2004). Eccentric training of the gastrocnemius-soleus complex in chronic Achilles tendinopathy results in decreased tendon volume and intratendinous signal as evaluated by MRI. <i>Am J Sports Med</i> 32, 1286-1296.	ž
1301 1302 1303 1304	Simpson DM, Tierra-Criollo CJ, Leite RT, Zayen EJ & Infantosi AF. (2000). Objective response detection in an electroencephalogram during somatosensory stimulation. <i>Ann Biomed Eng</i> 28, 691-698.	
1305 1306 1307 1308	Singh RE, Iqbal K, White G & Hutchinson TE. (2018). A Systematic Review on Muscle Synergies: From Building Blocks of Motor Behavior to a Neurorehabilitation Tool. <i>Appl Bionics Biomech</i> 2018 , 3615368.	,
1309 1310 1311 1312	Siu WL, Chan CH, Lam CH, Lee CM & Ying M. (2016). Sonographic evaluation of the effect of long- term exercise on Achilles tendon stiffness using shear wave elastography. <i>J Sci Med Sport</i> 19 883-887.	,
1313 1314 1315	Soteropoulos DS & Baker SN. (2006). Cortico-cerebellar coherence during a precision grip task in the monkey. <i>J Neurophysiol</i> 95, 1194-1206.	
1316 1317 1318	Sullivan MJL, Bishop SR & Pivik J. (1995). The Pain Catastrophizing Scale: Development and validation. <i>Psychological Assessment</i> 7, 524-532.	
1319 1320 1321	Szaro P, Witkowski G, Smigielski R, Krajewski P & Ciszek B. (2009). Fascicles of the adult human Achilles tendon - an anatomical study. <i>Ann Anat</i> 191, 586-593.	
1322 1323 1324	van Dijk CN, van Sterkenburg MN, Wiegerinck JI, Karlsson J & Maffulli N. (2011). Terminology for Achilles tendon related disorders. <i>Knee Surg Sports Traumatol Arthrosc</i> 19, 835-841.	
1325 1326 1327 1328	Vandenbroucke JP, Elm Ev, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, Poole C, Schlesselman J. & Egger M. (2007). Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. <i>Annals of Internal Medicine</i> 147, W-163-W-194.	I
1329		

- Wang HK, Lin KH, Su SC, Shih TT & Huang YC. (2012). Effects of tendon viscoelasticity in Achilles
 tendinosis on explosive performance and clinical severity in athletes. *Scand J Med Sci Sports*
- 1332 **22**, e147-155.

1333

- Williams ER, Soteropoulos DS & Baker SN. (2010). Spinal interneuron circuits reduce approximately
 10-Hz movement discontinuities by phase cancellation. *Proc Natl Acad Sci U S A* 107, 11098 11103.
- 1337