

Estimated morbimortality and costs attributable to child and adolescent obesity in Brazil from 2024 to 2060: a multistate life table study

Eduardo Augusto Fernandes Nilson^{1,2,3}, Ana Carolina Rocha de Oliveira⁴, Bruna Gusmão⁴, Raphael Barreto da Conceição Barbosa⁴

¹ – Oswaldo Cruz Foundation- Fiocruz, Brasilia, Brazil

² - Center for Epidemiological Research in Nutrition and Public Health, University of São Paulo, São Paulo, Brazil

³ – Universidad Autónoma de Chile, Chile

⁴ – Desiderata Institute, Rio de Janeiro, Brazil

Corresponding author: Eduardo Nilson (edunilson@gmail.com)

Abstract

Introduction

Childhood obesity is a major global public health issue globally and in Brazil. The impacts of childhood obesity include higher risk of disease during childhood and of obesity and non-communicable diseases in adulthood and represent an important epidemiological and economic burden to countries.

Methods

This study is based on the modeling of total costs of hospitalizations and the additional healthcare costs attributable to childhood obesity in the Brazilian National Health System.

Results

The hospitalizations of children and adolescents with obesity as a primary cause totaled

Int\$2.6 million to the Brazilian National Health System from 2013 to 2022, demonstrating

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

that obesity is rarely considered as a cause of hospitalization especially among children and adolescents. The additional costs of hospitalizations attributable to childhood obesity totaled Int\$101.5 million during the same period. The additional non-hospital, outpatient and medication cost attributable to childhood obesity in Brazil were estimated at Int\$6.0 million, so the total estimated healthcare costs were of approximately Int\$107.5 million in the last decade.

Conclusion

This study highlight that the costs of childhood obesity are not limited to the impacts on adult health and represent a relevant economic burden to the Brazilian National Health System and to families because of additional costs during childhood. Therefore, the prevention and control of childhood obesity is a public health priority that demands immediate and robust policies.

Introduction

Obesity among children and adolescents represents one of the greatest public health challenges today. Worldwide, childhood obesity has increased eightfold in the last four decades (1). According to the World Health Organization (WHO), in 2016, 340 million children and adolescents were overweight, and in 2020, 39 million children under the age of 5 were obese (2).

Considering consolidated analyses of cohort studies from 200 countries, different trends in height gain and variations in Body Mass Index (BMI) for age in children and adolescents were observed across different regions of the world. According to this study, in Brazil, cohorts of children and adolescents between 1985 and 2019 showed age-specific BMIs higher than WHO growth curves, consistent with the increasing trend in

the prevalence of overweight and obesity among children and adolescents in the country (3).

In Brazil, data from national surveys show that the prevalence of obesity in children aged 5 to 9 years increased from 2.4% to 14.2% between 1974 and 2009, while the prevalence among adolescents increased from 0.6% to 5.0% in the same period (4). However, according to the Food and Nutrition Surveillance System (Sisvan) in 2022, the prevalence of obesity was 16.3% for children aged 5 to 9 years and 12.8% among adolescents monitored by Primary Health Care. The situation of childhood obesity is even more severe, as more recently, the National Survey of Child Nutrition (Enani), conducted in 2019, revealed that the prevalence of obesity in children under 5 years old was 10.1% (5). Growing evidence shows that, compared to eutrophic children, children who are overweight or obese are more likely to face issues such as poorer health even in childhood, including type 2 diabetes, asthma, hypertension, sleep apnea, musculoskeletal problems, and metabolic disorders (6), as well as lower self-esteem, a higher likelihood of being bullied, and lower school attendance and performance. Additionally, children and adolescents with obesity are more likely to have health problems in the future, as childhood obesity is a strong predictor of obesity in adulthood, increasing the risk of non-communicable chronic diseases such as type 2 diabetes, cardiovascular diseases, and certain types of cancer. The future impacts of childhood obesity also include a higher likelihood of negative socioeconomic and labor consequences, such as lower employability, reduced productivity, and lower wages (7).

Given this association between childhood and adolescent obesity and negative health outcomes in adulthood, many studies have begun to project the impacts of childhood obesity in terms of morbidity and mortality due to associated comorbidities, as well as direct costs for its treatment and indirect costs to society resulting from higher morbidity

and mortality, early retirements, and productivity losses due to absenteeism and presenteeism (8)(9)(10)(11)(12).

Modeling studies of the impact of diseases and risk factors have been used to evaluate the impact of childhood obesity in other countries (13)(14), but they have not yet been developed for the Brazilian context. On the other hand, considering the adult population in Brazil, multi-state life table models have already been developed, from which it was possible to estimate the epidemiological burden of overweight and the associated direct and indirect costs. According to these studies, if the current trend of increasing overweight among adults continues, there will be approximately 5.26 million cases of diseases and 808.6 thousand deaths associated with overweight from 2021 to 2030, and the direct and indirect costs of these diseases will total around R\$49.2 billion during this period (15)(16).

However, the costs of childhood obesity do not only occur in adulthood, and in a groundbreaking study in the country, it was estimated that the direct costs attributable to childhood obesity in the Unified Health System (SUS) reached approximately R\$438.8 million in the last decade, including hospital, outpatient, and medication expenses (17).

In Brazil, there are still no projections of the epidemiological and economic burden of obesity in children and adolescents, so this study aims to develop a multi-state life table model to estimate the growth trends of childhood obesity, the cases and deaths from diseases associated with obesity, and the direct and indirect costs of childhood obesity from 2024 to 2044.

Methods

This study was based on the development and application of a multi-state life table model to estimate the future impacts of childhood and adolescent obesity in Brazil. The study was divided into the following stages: (1) projection of BMI evolution by age in children

and adolescents and BMI in Brazilian adults from 2024 to 2060, assuming the continuation of current trends of average BMI increase for each age group; (2) projection of changes in BMI by age in children and adolescents and BMI in adults from 2024 to 2060 under different scenarios of childhood obesity reduction; (3) estimation of cases and deaths attributable to obesity, assuming the continuation of current trends of average BMI increase for each age group and under different scenarios of childhood obesity reduction from 2024 to 2060; (4) estimation of direct costs to the Unified Health System (SUS) attributable to childhood and adolescent obesity from 2024 to 2060; (5) estimation of direct and indirect costs of diseases and deaths attributable to obesity in Brazil that could be avoided under different scenarios of childhood obesity reduction from 2024 to 2060. The age groups of 5 to 9 years, 10 to 14 years, and 15 to 19 years were considered to represent children and adolescents based on the aggregation of age groups available in health information systems and national surveys.

Estimation of Obesity Prevalence Trends

First, for the projection of BMI evolution by age in children and adolescents and BMI in Brazilian adults from 2024 to 2044, estimates obtained from the study by Rodriguez-Martinez et al. (2020) (18) were used to obtain average BMIs by sex and age and the trends in BMI growth by age in children and adolescents according to cohorts in Brazil from 1985 to 2019 using Prais Winsten regressions. The study by Estivaleti et al. (2022) (19) was used to obtain the trends in BMI growth in Brazilian adults according to projections from the Risk Factor Surveillance System for Chronic Disease Telephone Survey (Vigitel). The BMI information was then compiled and expanded in the multi-state life table model developed by Nilson et al. (2022) (15).

These trends in population obesity growth were then modeled for different scenarios of childhood obesity reduction in 2024 by 10% (most optimistic) and two scenarios with

smaller reductions, 5% and 2%, and their impact on adult obesity until 2060. For the modeling, it was assumed that after the reduction in obesity prevalence in 2024, the same prevalence would be maintained among children and adolescents until 2060, and that the same rates of obesity prevalence growth in the population estimated by Estivaleti et al. (19) and extrapolated to 2060 would be maintained. Furthermore, the obesity growth trend was smoothed in the model considering the midpoint of each 5-year age range by sex through simple linear regression for each age group according to the usual scenario (current trends maintained in the future, or BAU – business as usual), extrapolated for each scenario of reducing childhood and adolescent obesity prevalence.

Multi-State Life Table

The multi-state life table model assesses the association between overweight/obesity and eleven non-communicable chronic diseases (NCDs): ischemic heart disease, cerebrovascular diseases, hypertensive heart disease, diabetes, chronic kidney disease, liver cirrhosis, and cancers of the breast, colon, pancreas, kidney, and liver. The RR values for each NCD by age group were extracted to estimate the increased disease risk for each 5 kg/m² increase in BMI. Data on incidence, prevalence, and mortality from NCDs were extracted from the Global Burden of Disease (GBD) study (20), stratified by sex and age group (considering 5-year age ranges), identified by the International Classification of Diseases 10 (ICD-10) codes, and data on case fatality and remission were estimated using the Dismod II tool (21).

The cohort simulation model by multi-state life table (MSLT) of cases and deaths from 11 NCDs was adapted from another validated study for estimating NCDs attributable to overweight in Brazil (Nilson et al., 2022). The model considers a hypothetical cohort of individuals who, in each cycle (year), are classified as healthy, ill, or deceased based on the incidence, mortality, remission, and case fatality rates of the diseases considered in

the study. For this, within each sex and age stratum and for each disease evaluated, the RR values and the prevalence of overweight and non-communicable chronic diseases were considered, through systematic reviews and meta-analyses. The BMI distribution in the population was modeled considering a log-linear function of the average BMI and its standard deviation and the national population for each sex and age stratum.

Finally, within each sex and age stratum, for the scenario of maintaining BMI increase trends and for each reduction scenario, we calculated the potential impact fraction (PIF) for outcomes (o) in each age group (a) and sex (s). The PIF results represent the percentage difference in incidence and mortality rates of NCDs attributable to scenarios of population BMI increase. The model assumes that the impact of obesity reduction on NCD incidence and mortality rates would not occur immediately, so latency periods were incorporated into the model between BMI changes and outcomes, considering an average of 5 years for cardiovascular diseases, type 2 diabetes, and chronic kidney disease, and an average of 10 years for cancers. The uncertainty of cases and deaths attributable to different scenarios was quantified through Monte Carlo simulations ($n = 5,000$), where for each simulation, a draw was made from the distributions (uncertainties) of BMI and RR, which were reported for the 50th, 2.5th, and 97.5th percentiles of estimates across all simulations as the central estimate and 95% uncertainty intervals, respectively.

Estimates of Direct Costs attributable to Obesity

For estimating the direct costs to the Unified Health System (SUS) attributable to childhood and adolescent obesity until 2044, the same methodology by Nilson et al. (2023) (22) was used, which estimates excess costs for hospitalizations of children and adolescents with obesity compared to those with an adequate BMI, using data from the SUS Hospital Information System (SIH/SUS), using the base year of 2022. Next, to estimate non-hospital, outpatient, and medication expenses, it was assumed that these

would be proportional to additional hospitalization costs according to the meta-analysis by Ling et al., 2023 (23).

For estimating direct costs in adulthood, the cost data for NCDs in the SUS were extracted using the Tabnet tool for hospitalizations (Hospital Information System – SIH/SUS) and outpatient care (Outpatient Information System – SIA/SUS), publicly available from the SUS Department of Informatics (DATASUS). Costs were extracted by sex and age group (considering 5-year age ranges) and for the 11 diseases included in the multi-state life table model. To calculate the per capita cost of NCDs in the SUS, the costs from SIH and SIA were used, considering the average cost per patient for the SUS. This average cost per patient (numerator) was multiplied by the number of incident cases attributable to obesity until 2044.

Further details of the parameters for the estimations and the modeling methodology are described in the Supplementary Materials.

Results

Future Trends in Childhood and Adolescent Obesity prevalences and direct costs

Following the trends observed in Brazilian cohorts from 1985 to 2019, estimates show that obesity prevalence will increase across all age groups for both sexes until 2060. From 2023 to 2060, among boys aged 5 to 9 years, obesity prevalence will rise from 22.2% to 34.7%, while it will increase from 14.3% to 24.9% among girls of the same age. In the same period, obesity prevalence will increase from 12.6% to 21.9% among boys aged 10 to 14 years and from 8.3% to 15.5% among girls in this age group. Finally, in the next two decades, for the 15 to 19 age group, obesity prevalence among boys will rise from 8.6% to 16.0% and from 7.9% to 14.9% among girls.

Following the same methodology adopted in the previous study on direct costs of childhood and adolescent obesity, we used the annual per capita costs in 2022 as a reference to estimate hospital, non-hospital, outpatient, and medication costs attributable to obesity in individuals aged 5 to 19 years in Brazil from 2013 to 2022.

If current trends in prevalence increases continue, the annual direct attributable costs would rise from R\$83.2 million in 2023 to R\$115.5 million in 2060, totaling R\$3.84 billion over the period. Most of these costs (48.1%) would be in the 15 to 19 age group, while 18.8% would occur in the 10 to 14 age group, and 33.2% of these costs would be among individuals aged 5 to 9 years. In terms of cost distribution by type, 94.4% would be spent on hospital expenses, 2.7% on non-hospital expenses, 2.2% on medications, and 0.7% on outpatient care and procedures (Table 1).

Scenarios for Reducing Childhood and Adolescent Obesity

If current trends in obesity prevalence growth continue across all age groups in Brazil, the number of people aged 0 to 79 with obesity will rise from 26.7 million in 2023 to 38.6

million in 2060, representing a 44% increase. In terms of sex and age groups, this corresponds to 4.7 million boys and 2.8 million girls with obesity in 2023, and 5.9 million boys and 3.8 million girls with obesity in 2060 (an increase of 22% to 36%). Among adults, it is estimated that there would be 10.1 million men and 9.2 million women with obesity in 2023, and if the current trends in prevalence growth continue, these numbers would rise to 15.4 million men and 13.8 million women in 2060 (increases of 50% to 52%).

Considering the scenarios for reducing childhood and adolescent obesity, a 10% reduction in obesity prevalence among individuals aged 5 to 19 would slow the growth in adult obesity prevalence among those up to 59 years old. While in the baseline scenario (BAU – business-as-usual), which reflects the continuation of current trends, there would be an average 99.5% increase in obesity prevalence among adults (ranging from 86.5% to 148%, depending on sex and age group), in the scenario with a 10% reduction in obesity prevalence among children and adolescents, the average increase in adult obesity prevalence would be 17.8% (ranging from 1.2% to 72.6%). In the intermediate scenarios, the average increase in obesity prevalence among adults up to 59 years old would decrease to 23.4% for a 5% reduction in childhood and adolescent obesity and to 26.8% for a 2% reduction, with ranges from 6.8% to 75.0% and from 10.2% to 76.4%, respectively (Tables 2 and 3).

These impacts illustrate the combined effect of the initial reduction in childhood and adolescent obesity prevalence, coupled with its maintenance over time, ensuring that these impacts gradually extend to adults in the future, following the cohort in the life table. For example, reductions among individuals aged 15 to 19 will first impact younger adults (aged 20 to 24) in the first five years of the simulation. Over time, more adult age groups will be affected, eventually reaching those aged 55 to 59 by 2060.

Impact of Reduction Scenarios on the Direct Costs Attributable to Childhood and Adolescent Obesity

In terms of the potential impact of reducing and subsequently maintaining the prevalence of obesity among children and adolescents, significant cost savings could be achieved in all scenarios when considering the costs attributable to childhood and adolescent obesity within the SUS. In the most optimistic scenario, with a 10% reduction in prevalence, approximately R\$1.274 billion could be saved by 2060, while a 5% reduction would lead to savings of R\$1.136 billion, and a 2% reduction would result in savings of R\$1.053 billion over the same period (Table 4).

Table 1. Estimates of the future direct costs attributable to child and adolescent obesity from 2024 to 2060 in Brazil (in Brazilian Reals – R\$).

Hospital costs	2023	2025	2030	2035	2040	2045	2050	2055	2060	TOTAL
5 to 9 years	27,496,137	28,482,189	30,208,523	30,968,568	31,509,636	32,116,279	32,900,682	33,870,021	34,787,960	1,203,582,572
10 to 14 years	14,328,395	14,716,665	16,292,204	17,449,886	18,063,685	18,557,366	19,096,266	19,749,071	20,522,946	680,735,594
15 to 19 years	36,724,375	37,193,437	39,070,282	43,439,423	46,720,458	48,561,146	50,086,912	51,741,716	53,713,170	1,743,963,404
	78,548,907	80,392,291	85,571,008	91,857,877	96,293,779	99,234,792	102,083,859	105,360,808	109,024,076	3,628,281,569
Non hospital costs										
5 to 9 years	786,853	815,071	864,473	886,223	901,707	919,067	941,514	969,253	995,522	34,442,744
10 to 14 years	410,034	421,145	466,232	499,361	516,926	531,053	546,475	565,156	587,302	19,480,510
15 to 19 years	1,050,936	1,064,359	1,118,068	1,243,100	1,336,992	1,389,667	1,433,330	1,480,685	1,537,102	49,906,743
	2,247,822	2,300,574	2,448,773	2,628,683	2,755,625	2,839,787	2,921,319	3,015,095	3,119,926	103,829,997
Outpatient costs										
5 to 9 years	198,662	205,787	218,260	223,751	227,660	232,043	237,711	244,714	251,346	8,696,001
10 to 14 years	103,524	106,329	117,713	126,077	130,512	134,079	137,972	142,689	148,280	4,918,381
15 to 19 years	265,337	268,726	282,287	313,854	337,560	350,859	361,883	373,839	388,083	12,600,305
	567,523	580,842	618,259	663,682	695,732	716,981	737,566	761,242	787,710	26,214,686
Medications										
5 to 9 years	645,687	668,842	709,382	727,230	739,935	754,181	772,601	795,364	816,920	28,263,526
10 to 14 years	336,471	345,589	382,587	409,773	424,186	435,779	448,434	463,764	481,937	15,985,599
15 to 19 years	862,392	873,407	917,481	1,020,081	1,097,129	1,140,353	1,176,182	1,215,042	1,261,337	40,953,198
	1,844,551	1,887,839	2,009,450	2,157,083	2,261,251	2,330,314	2,397,218	2,474,170	2,560,194	85,202,323
TOTAL										
5 to 9 years	29,127,339	30,171,889	32,000,637	32,805,771	33,378,938	34,021,570	34,852,507	35,879,353	36,851,748	1,274,984,843
10 to 14 years	15,178,424	15,589,728	17,258,735	18,485,097	19,135,310	19,658,278	20,229,148	20,920,680	21,740,466	721,120,083
15 to 19 years	38,903,041	39,399,929	41,388,117	46,016,457	49,492,139	51,442,026	53,058,307	54,811,281	56,899,692	1,847,423,649
	83,208,804	85,161,546	90,647,489	97,307,325	102,006,387	105,121,874	108,139,962	111,611,314	115,491,906	3,843,528,575

Table 2. Estimated prevalences according to the maintenance of current trends and in different scenarios of reduction of the prevalence of obesity among children and adolescents in Brazil from 2024 to 2060.

	Current trends		10% reduction		5% reduction		2% reduction	
	2023	2060	2023	2060	2023	2060	2023	2060
Men								
20 to 24 years	15.34%	30.85%	15.34%	15.54%	15.34%	16.41%	15.34%	16.93%
25 to 29 years	29.49%	59.25%	29.49%	31.51%	29.49%	33.26%	29.49%	34.31%
30 to 34 years	36.82%	69.18%	36.82%	42.09%	36.82%	44.43%	36.82%	45.83%
35 to 39 years	37.37%	69.88%	37.37%	43.00%	37.37%	45.38%	37.37%	46.82%
40 to 44 years	37.70%	70.83%	37.70%	42.87%	37.70%	45.25%	37.70%	46.68%
45 to 49 years	36.12%	69.84%	36.12%	40.17%	36.12%	42.40%	36.12%	43.74%
50 to 54 years	36.27%	71.08%	36.27%	40.21%	36.27%	42.44%	36.27%	43.78%
54 to 59 years	34.08%	68.78%	34.08%	56.29%	34.08%	57.12%	34.08%	57.61%
Women								
20 to 24 years	24.37%	40.09%	24.37%	24.67%	24.37%	26.04%	24.37%	26.86%
25 to 29 years	31.99%	61.08%	31.99%	34.11%	31.99%	36.01%	31.99%	37.14%
30 to 34 years	34.38%	65.00%	34.38%	39.30%	34.38%	41.49%	34.38%	42.80%
35 to 39 years	35.58%	66.93%	35.58%	40.94%	35.58%	43.21%	35.58%	44.57%
40 to 44 years	36.21%	67.48%	36.21%	41.16%	36.21%	43.44%	36.21%	44.81%
45 to 49 years	36.82%	69.90%	36.82%	40.96%	36.82%	43.23%	36.82%	44.60%
50 to 54 years	32.22%	67.46%	32.22%	35.71%	32.22%	37.70%	32.22%	38.89%
54 to 59 years	30.83%	65.41%	30.83%	53.23%	30.83%	53.96%	30.83%	54.40%

Table 3. Estimates of the predicted number of individuals with obesity in Brazil from 2024 to 2060 by sex and age group.

	2023	2025	2030	2035	2040	2045	2050	2055	2060
Men									
5 to 9 years	1,677,255	1,717,557	1,785,681	1,812,848	1,835,969	1,866,307	1,907,812	1,955,494	1,992,756
10 to 14 years	1,554,238	1,606,763	1,682,035	1,713,656	1,737,900	1,767,555	1,807,421	1,855,549	1,897,024
15 to 19 years	1,444,340	1,493,108	1,578,791	1,614,732	1,639,954	1,668,819	1,706,930	1,754,372	1,798,582
20 to 24 years	1,274,297	1,316,212	1,410,170	1,449,871	1,476,604	1,505,084	1,541,536	1,587,368	1,632,998
25 to 29 years	1,088,077	1,136,473	1,224,829	1,271,932	1,300,484	1,328,935	1,363,950	1,407,695	1,453,807
30 to 34 years	939,068	982,464	1,067,391	1,120,514	1,150,693	1,179,091	1,212,649	1,254,008	1,299,662
35 to 39 years	801,436	829,233	917,650	972,956	1,004,926	1,033,302	1,065,430	1,104,290	1,148,689
40 to 44 years	694,657	715,541	796,111	855,486	888,967	917,005	947,503	983,810	1,026,209
45 to 49 years	671,262	686,832	758,640	826,514	862,401	890,514	919,834	954,440	995,485
50 to 54 years	664,353	673,581	740,699	809,518	852,158	880,787	909,249	942,433	982,084
55 to 59 years	662,109	670,896	727,927	799,725	848,921	878,535	906,458	938,376	976,580
60 to 64 years	693,458	706,958	756,833	850,002	913,459	953,983	989,968	1,028,776	1,073,558
65 to 69 years	763,991	782,429	841,448	956,642	1,051,392	1,113,445	1,166,320	1,220,212	1,279,658
70 to 74 years	844,149	866,579	943,875	1,081,590	1,223,831	1,318,163	1,396,909	1,473,621	1,554,973
75 to 79 years	1,015,269	1,040,802	1,138,631	1,306,845	1,492,222	1,632,757	1,737,071	1,834,041	1,933,642

(Table 3 – continuation)

	2023	2025	2030	2035	2040	2045	2050	2055	2060
Women									
5 to 9 years	1,029,984	1,062,418	1,120,224	1,154,680	1,187,167	1,224,751	1,270,119	1,319,932	1,362,934
10 to 14 years	928,074	966,314	1,025,906	1,061,037	1,092,302	1,127,446	1,169,592	1,217,484	1,261,319
15 to 19 years	813,293	846,719	909,120	944,677	974,623	1,007,213	1,045,885	1,090,843	1,134,303
20 to 24 years	723,421	752,072	819,265	855,574	884,492	914,749	950,282	992,209	1,034,797
25 to 29 years	639,156	670,947	733,857	772,911	800,690	828,660	861,110	899,601	940,374
30 to 34 years	589,685	618,543	679,961	721,878	749,021	775,221	805,139	840,645	879,677
35 to 39 years	542,192	561,917	627,732	671,414	699,278	724,754	753,087	786,502	824,400
40 to 44 years	529,436	545,798	610,056	658,955	688,341	713,555	740,835	772,863	809,988
45 to 49 years	519,423	531,613	589,387	645,313	676,631	701,860	728,159	758,828	794,903
50 to 54 years	551,309	558,908	615,121	674,316	712,436	738,527	764,552	794,673	830,431
55 to 59 years	586,472	594,139	643,692	708,052	753,106	780,498	806,429	835,996	871,247
60 to 64 years	709,509	723,594	769,367	862,698	923,219	959,316	989,351	1,021,255	1,058,453
65 to 69 years	929,183	949,884	1,008,210	1,132,458	1,230,156	1,283,636	1,322,169	1,359,187	1,400,589
70 to 74 years	1,256,785	1,277,344	1,358,874	1,516,134	1,675,318	1,755,987	1,807,987	1,852,651	1,900,099
75 to 79 years	1,602,398	1,621,787	1,723,759	1,914,564	2,120,283	2,244,384	2,309,785	2,360,394	2,411,380

Table 4. Estimated savings in the direct health costs attributable to child and adolescent obesity from 2024 to 2060 in different scenarios of prevalence reduction in Brazil (in Brazilian Reals – R\$).

TOTAL	10%	5%	2%
5 to 9 years	382,196,079	334,214,889	305,426,175
10 to 14 years	240,655,798	214,806,583	199,297,055
15 to 19 years	651,362,302	587,075,730	548,503,786
	1,274,214,180	1,136,097,203	1,053,227,016

Table 5. Prevented or averted incident cases of obesity-related diseases from 2024 to 2060 in different scenarios of obesity reduction among children and adolescents in Brazil (Mean and Uncertainty Intervals 95%).

Scenario	Incident cases	Deaths
10%		
Men	74,227 (44,818-101,972)	26,298 (13,391-37,486)
Women	170,034 (80,730-244,107)	44,542 (17,366-64,952)
Total	244,261 (125,547-346,079)	70,840 (30,757-102,438)
5%		
Men	65,847 (39,744-90,476)	23,371 (11,899-33,319)
Women	151,732 (72,052-217,867)	39,817 (15,523-58,072)
Total	217,579 (111,796-308,343)	63,189 (27,422-91,391)
2%		
Men	60,937 (36,772-83,738)	21,651 (11,022-30,869)
Women	140,996 (66,963-202,469)	37,038 (14,439-54,024)
Total	201,932 (103,734-286,207)	58,689 (25,461-84,893)

Table 6. Incident cases of obesity related diseases prevented from 2024 to 2060 in different scenarios of obesity reduction among children and adolescents by disease group in Brazil (Mean and Uncertainty Intervals 95%).

	Prevented cases		
	2%	5%	10%
Ischaemic heart disease	7,743 (3,623-11,298)	8,320 (3,894-12,139)	9,294 (4,351-13,560)
Cerebrovascular diseases	23,959 (11,792-34,629)	25,779 (12,687-37,259)	28,873 (14,210-41,730)
Hypertensive heart disease	17,24 (832-2,391)	1,857 (896-2,576)	2,085 (1,006-2,893)
Diabetes	116,268 (64,109-157,003)	125,451 (69,174-169,397)	141,167 (77,843-190,611)
Cancers	136 (73-207)	147 (79-224)	165 (89-251)
Cirrhosis	11,715 (6,464-16,608)	12,627 (6,968-17,901)	14,178 (7,826-20,099)
Chronic kidney disease	40,388 (16,842-64,070)	43,399 (18,097-68,846)	48,499 (20,224-76,935)
Total	201,932 (103,734-286,207)	217,579 (111,796-308,343)	244,261 (125,547-346,079)

Table 7. Deaths of obesity related diseases prevented from 2024 to 2060 in different scenarios of obesity reduction among children and adolescents by disease group in Brazil (Mean and Uncertainty Intervals 95%).

Deaths			
	2%	5%	10%
Ischaemic heart disease	7,767 (3,239-11,404)	8,345 (3,239-11,404)	9,322 (3,890-13,687)
Cerebrovascular diseases	15,072 (5,991-23,457)	16,218 (5,991-23,457)	18,164 (7,223-28,266)
Hypertensive heart disease	1,769 (766-2,447)	1,906 (766-2,447)	2,141 (927-2,960)
Diabetes	15,281 (7,136-19,741)	16,480 (7,136-19,741)	18,531 (8,658-23,941)
Cancers	134 (63-206)	144 (63-206)	162 (76-250)
Cirrhosis	10,900 (5,219-15,382)	11,751 (5,219-15,382)	13,197 (6,323-18,623)
Chronic kidney disease	7,767 (3,047-12,256)	8,345 (3,047-12,256)	9,322 (3,660-14,710)
Total	58,689 (25,461-84,893)	63,189 (25,461-84,893)	70,840 (30,757-102,438)

1 DISCUSSION

2 According to this study, if the current trends in childhood and adolescent obesity in Brazil
3 continue, the prevalence will significantly increase across different age subgroups and for
4 both sexes by 2060, potentially nearly doubling among children aged 10 to 14 and
5 adolescents. As a result, the costs attributable to childhood and adolescent obesity to the
6 SUS (Unified Health System) could reach R\$3.84 billion during this period. On the other
7 hand, if obesity prevalence among children and adolescents is reduced and remains
8 constant after 2040, these direct costs could be reduced by R\$1.05 to R\$1.27 billion by
9 2060.

10 Additionally, considering the impact of childhood and adolescent obesity on adulthood,
11 if the trends continue until 2060, obesity among adults will nearly double. However, in
12 scenarios where obesity prevalence in childhood and adolescence is reduced and then
13 maintained, there would be a significant slowdown in the rate of obesity increase among
14 adults. As a consequence of this reduction, across different scenarios studied (2% to 10%
15 reduction in childhood and adolescent obesity prevalence), up to 244,600 incident cases
16 and 70,800 deaths from obesity-related diseases could be prevented. Among the
17 outcomes, cardiovascular diseases, diabetes, and chronic kidney disease had the greatest
18 impact on the prevented cases and deaths due to reduced obesity prevalence in childhood
19 and adolescence.

20 These estimates demonstrate that reducing childhood and adolescent obesity is crucial to
21 lowering adult obesity and its consequences on the future morbidity and mortality of
22 Brazilians. Moreover, the impact on direct health costs is likely to grow even more in the
23 future if current trends continue, leading to a significant increase in direct and indirect
24 costs attributable to obesity in the adult population in the future.

25 Regarding the costs to the SUS attributable to obesity, it should be noted that they already
26 reached R\$1.42 million in 2018, and future projections indicate that the direct and indirect

costs of obesity could reach 6.09% of Brazil's Gross Domestic Product, approximately US\$243.23 billion.

If current trends in the growth of obesity and overweight among the Brazilian adult population continue, their prevalence will reach 29.6% and 68.1%, respectively, by 2030. During this period, 5.26 million new cases and 808,000 deaths from cardiovascular diseases, cancers, diabetes, chronic kidney diseases, among others, will be attributable to excess weight by the end of this decade. Additionally, another study using the same simulation model applied in this research estimated that, between 2020 and 2030, around R\$4.2 billion would be spent on direct expenses of NCDs if the current trend in the prevalence of excess weight among adults in Brazil continues, with indirect costs associated with premature mortality reaching approximately R\$45.5 billion in the same period.

These changes in obesity prevalence across all age groups, including children and adolescents, are strongly associated with the nutritional transition that Brazil is undergoing, with the gradual replacement of the traditional diet—composed of rice, beans, cereals, fruits, vegetables, and other food types—by ultra-processed products in all age groups, including children and adolescents. According to the National Study on Child Nutrition (Enani 2019), about 80% of children under 2 years old consumed ultra-processed products, despite the Ministry of Health's recommendation against this practice. It is even more concerning that, according to the 2017-2018 Household Budget Survey by the Brazilian Institute of Geography and Statistics (POF/IBGE), the highest proportion of ultra-processed foods in total dietary energy is among adolescents (reaching 26.7% of dietary calories) compared to adults and the elderly.

A growing body of scientific evidence reinforces that the consumption of ultra-processed products is directly related to a higher risk of overweight and obesity, as well as other noncommunicable diseases such as cardiovascular diseases, diabetes, and certain types

of cancer. In addition to studies based on cohorts and surveys, and meta-analyses derived from them, a recent randomized clinical trial demonstrated that a diet based on ultra-processed foods compared to a diet based on whole and minimally processed foods is associated with higher calorie intake and weight gain.

Childhood obesity is recognized as a risk factor for obesity in adulthood, with 55% of children with obesity likely to become adolescents with obesity, and about 80% of adolescents with obesity expected to have obesity by the age of 30. Furthermore, childhood obesity is associated with a moderate increase in the risk of morbidity in adulthood, including diabetes, cardiovascular diseases (such as ischemic heart disease and hypertension), and some types of cancer.

Moreover, childhood obesity is associated with higher healthcare costs, including hospitalizations, outpatient costs, medication expenses, and other costs to families, as well as longer hospital stays for various diseases exacerbated by obesity, which also entails significant direct and indirect costs among adults. In Brazil, the costs attributable to childhood and adolescent obesity for the treatment of diseases within the SUS up to the age of 19 exceed the total costs of hospitalizations for diseases classified under Chapter V of the ICD-10, which corresponds to endocrine, nutritional, and metabolic diseases.

The current study is based on methodologies already validated in other studies on the projection of the impact of childhood and adolescent obesity on adulthood, and the current adaptation of a multi-state life table model for the Brazilian population from age 5 onwards could serve as a useful tool for new studies on the burden of obesity-related diseases and for modeling the potential impact of policies and interventions on health and nutrition in childhood, adolescence, and adulthood, as well as evaluating their costs.

The results highlight the significant epidemiological and economic impact of childhood and adolescent obesity in Brazil, emphasizing the need and urgency for more cost-

effective policies to treat overweight and obesity within the SUS from childhood, along with the adoption of regulatory and fiscal policies that promote healthier food environments throughout the life cycle, especially in childhood and adolescence. These nutrition protection measures begin with the promotion and protection of breastfeeding and complementary feeding and continue through the regulation of food advertising, the regulation of ultra-processed food sales in schools, the improvement of nutritional labeling, and the taxation of ultra-processed products, along with subsidies for whole and minimally processed foods, as well as the strengthening of food and nutrition education actions for the population based on the Brazilian Dietary Guidelines and the Dietary Guidelines for Children Under 2 Years Old.

As strengths of this study, the direct cost estimation of obesity was based on modeling from a recent and robust meta-analysis on the costs of childhood and adolescent obesity worldwide and hospital expenses of the Brazilian public health system. For the prevalence of childhood and adolescent obesity, published data from a set of Brazilian cohorts were used, covering nutritional data for individuals aged 5 to 19 years up to 2019, including to model future trends in weight gain among children and adolescents. Finally, the modeling of the impact of changes in childhood obesity on adult health outcomes was adapted from a validated and published multi-state life table model. Additionally, the methods adapted and developed in this study could serve as a basis for new analyses and projections, including the potential impact of policies to reduce obesity among children and adolescents.

Among its limitations, we have the adoption of hospital cost estimates from primary studies in other countries, with different demographic, socioeconomic characteristics, and health systems compared to Brazil, possible limitations in the actual expenditures within the SUS, the assumptions adopted in incorporating the cohort's nutritional data into the

life table model, as well as the limitations already described regarding the multi-state life table for Brazil.

However, the assumptions adopted in the various parts of this study aimed to generate more conservative estimates for the outcomes, while also incorporating uncertainty analysis through Monte Carlo analysis for the projections of cases and deaths attributable in adulthood. Future analyses should further consider the economic evaluation of the impact of childhood and adolescent obesity and analyze new scenarios of changes in prevalence, as well as expand the model to years beyond 2060 when new population projections are generated by the IBGE.

In conclusion, the results of this study add to those of the previous study of this project, reinforcing that childhood obesity is a major public health problem and a significant epidemiological and economic issue for Brazilian society, considering its current and future impacts if nothing is done. Therefore, immediate and effective measures are essential to address it, including treatment and prevention policies.

Declarations

Ethics approval and consent to participate

Not applicable.

Availability of data and materials

All relevant data on the model are publicly available. Hospitalization data is available from <http://tabnet.datasus.gov.br/cgi/defthtm.exe?sih/cnv/niuf.def> and IHME

Competing interests

The authors declare that they have no competing interests.

Funding

The authors declare no funding.

Authors' contributions

EAFN conceptualised the idea and was responsible for the study design and data analysis and drafted the original manuscript. All authors revised and approved the final manuscript.

References

1. Ling J, Chen S, Zahry NR, Kao TSA. Economic burden of childhood overweight and obesity: A systematic review and meta-analysis. *Obes Rev* [Internet]. 2023 Feb 1 [cited 2023 Mar 7];24(2):e13535. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/obr.13535>
2. IHME. Global burden of disease 2019 [Internet]. 2020. Available from: <https://vizhub.healthdata.org/gbd-compare>
3. Briggs ADM, Cobiac LJ, Wolstenholme J, Scarborough P. PRIMEtime CE: A multistate life table model for estimating the cost-effectiveness of interventions affecting diet and physical activity. *BMC Health Serv Res* [Internet]. 2019 Jul 16 [cited 2021 Nov 19];19(1):1–19. Available from: <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-019-4237-4>
4. Cleghorn C, Blakely T, Nghiem N, Mizdrak A, Wilson N. Technical Report for BODE3 Diet Intervention and Multistate Lifetable Models - Technical Report Number 16 [Internet]. 2017. Available from: <https://www.otago.ac.nz/wellington/otago670797.pdf>
5. Ministério da Saúde. Vigitel Brazil 2019: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian sta [Internet]. Brasília, Brasil; 2020 [cited 2021 Sep 10]. 139 p. Available from: http://bvsms.saude.gov.br/bvs/publicacoes/vigitel_brasil_2019_vigilancia_fatores_risco.pdf
6. Barendregt JJ, van Oortmarssen GJ, Vos T, Murray CJL. A generic model for the assessment of disease epidemiology: The computational basis of DisMod II. *Popul Health Metr* [Internet]. 2003;1(1):4. Available from: <https://doi.org/10.1186/1478-7954-1-4>