

Fat burning capacity in a mixed macronutrient meal protocol does not reflect metabolic flexibility in women who are overweight or obese.

Mary M. Ahern¹⁺, Virginia M. Artegoitia²⁺, Rémy Bosviel³, John W. Newman^{2,3,4}, Nancy L. Keim^{2,4} Sridevi Krishnan^{1¥}

Affiliations

1 – School of Nutritional Sciences and Wellness, University of Arizona, Tucson AZ 85721

2 – United States Department of Agriculture, Agricultural Research Service, Western Human Nutrition Research Center, Davis, CA 95616, USA

3 – West Coast Metabolomics Center, Genome Center, University of California Davis, Davis, CA 95616, USA

4 – Department of Nutrition, University of California, Davis, Davis, CA 95616, USA

+ co-first authors

¥ corresponding author

Corresponding author information

Sridevi Krishnan: skrishnan@arizona.edu

Funding information: National Dairy Council, Campbell Soup Co., USDA- ARS Projects 2032-51530-022-00D and 2023-51530-025-00D. We acknowledge the West Coast Metabolomics Center Pilot and Feasibility study to N.K. and S.K. funded through National Institutes of Health Grant U24 DK097154.

Clinical Trials Information: [Clinicaltrials.gov: NCT02298725](https://clinicaltrials.gov/ct2/show/study/NCT02298725)

Key words: metabolic flexibility, mixed macronutrient challenge test, metabolites, lipidomics

Running title: Fat burning capacity and metabolic flexibility

Abstract

Introduction: Metabolic flexibility, the ability to switch from glucose to fat as a fuel source, is considered a marker of metabolic health. Higher fat oxidation is often associated with greater flexibility and insulin sensitivity, while lower fat oxidation is linked to metabolic inflexibility and insulin resistance. However, our study challenges the universal validity of this relationship, uncovering a more nuanced understanding of the complex interplay between fuel source switching and fat oxidation, especially in the presence of insulin resistance.

Methods: In an 8-week controlled feeding intervention, overweight to obese women with insulin resistance (as defined by McAuley's index) were randomized to consume either a diet based on the Dietary Guidelines for Americans 2010 (DGA) or a 'Typical' American Diet (TAD), $n = 22$ each. Participants were given a high-fat mixed macronutrient challenge test (MMCT) (60% fat, 28% carbohydrates, and 12% protein) at weeks 0, 2, and 8. Plasma lipids, metabolome, and lipidome were measured at 0, 0.5, 3, and 6h postprandial (PP); substrate oxidation measures were also recorded at 0, 1.5, 3, and 6h PP. Metabolic flexibility was evaluated as the change in fat oxidation from fasting to PP. Mixed model and multivariate analyses were used to evaluate the effect of diet on these outcomes, and to identify variables of interest to metabolic flexibility.

Results: Intervention diets (DGA and TAD) did not differentially affect substrate oxidation or metabolic flexibility, and equivalence tests indicated that groups could be combined for subsequent analyses. Participants were classified into three groups based on the % of consumed MMCT fat was oxidized in the 6h post meal period at weeks 0, 2 and 8. Low fat burners (LB, $n = 6$, burned <30% of fat in MMCT) and high fat burners (HB, $n = 7$, burned > 40% of fat in MMCT) at all weeks. Compared to LB, HB group had higher fat mass, total mass, lean mass, BMI, lower HDLc and lower RER ($p < 0.05$), but not different % body fat or % lean mass.

During week 0, at 1h PP, LB had an increase in % fat oxidation change from 0h compared to HB ($p < 0.05$), suggesting higher metabolic flexibility. This difference disappeared later in the PP phase, and we did not detect this beyond week 0. Partial least squares discriminant analysis (PLSDA (regular and repeated measures (sPLSDA))) models identified that LB group, in the late PP phase, was associated with higher rates of disappearance of acylcarnitines (AC) and lysophosphatidylcholines (LPC) from plasma (Q^2 : 0.20, R^2X : 0.177, R^2Y : 0.716).

Conclusion: In women with insulin resistance, a high fat burning capacity does not imply high metabolic flexibility, and not all women with insulin resistance are metabolically inflexible. LPCs and ACs are promising biomarkers of metabolic flexibility.

Introduction

Metabolic flexibility refers to the ability of an organism to adapt to the energetic push and pull and switching between fuel substrates allowing for an efficient metabolic response to physiological needs and opportunity¹. This flexibility is an important component of metabolic health, being a key metabolic adaptation that balances choice of fuel storage and oxidation. It can be measured as the shift in metabolism to use the more abundant fuel present in a given meal, e.g. fat, rather than defaulting to glucose².

The functional capacity of skeletal muscle tissue, the most metabolically active tissue, plays a key role in metabolic flexibility³. The muscle of lean individuals is considered metabolically flexible, as it readily adapts to available fuels in response to insulin⁴. In contrast, obese-insulin resistant individuals are metabolically inflexible, and experience stunted insulin-stimulated suppression of fat oxidation following a meal with glucose. Metabolic flexibility is also closely linked to fat oxidation rates. Lower rates of maximal fat oxidation are associated with reduced metabolic flexibility^{5,6}. Additionally, impaired insulin sensitivity and metabolic flexibility are linked to reduced fat oxidation under resting conditions in skeletal muscle⁶. Collectively, this suggests that skeletal muscle flexibility and the ability to oxidize lipids effectively are inextricably related to metabolic flexibility and this relationship ultimately impacts an individual's health.

Perturbations in metabolic flexibility are associated with insulin resistance, metabolic dysfunction-associated fatty liver disease (MAFLD), type 2 diabetes and cardiovascular disease⁷⁻¹². Accurate measurement of metabolic flexibility can allow for its use as a biomarker of health. Metabolic flexibility is measured systemically in humans using indirect calorimetry during a euglycemic-hyperinsulinemic clamp¹³. Oral glucose tolerance tests, fasting and re-feeding tests,

and oxygen restriction tests have also been used¹⁴. Currently, however, there is no consensus on when and how to measure metabolic flexibility.

Diet, especially fat composition, may impact metabolic flexibility. In mice, a high saturated fatty acid (SFA)-rich diet compared to a polyunsaturated fatty acid (PUFA)-rich diet reduced metabolic flexibility while increasing adiposity, liver damage and visceral fat deposits¹⁸. However, in humans, high-fat diets appear to be more useful as metabolic tests for long-term health¹⁷ and whole diet approaches, without weight loss, have not been successful at altering metabolic flexibility¹⁹. High-fat metabolic stress tests may be an effective way of measuring metabolic flexibility due to the stimulation of many metabolic systems simultaneously. Researchers have studied the association between dietary or meal fatty acid composition and subsequent systemic substrate oxidation^{12,20-22} but only a few studies have looked closely at circulating fatty acids and substrate oxidation^{23,24}. The rise of high throughput omic tools has made it possible to get molecular insight into metabolic states. Subsequently, recent studies have attempted to deconstruct metabolic flexibility using lipidomic and metabolomic tools especially after a high-fat meal challenge test^{16,25,26}.

This manuscript presents secondary outcomes from a randomized clinical trial that has been published²⁷. The clinical trial compared metabolic effects of following a diet based on the Dietary Guidelines for Americans (DGA) to a Typical American diet (TAD) for 8-weeks in women (n = 44, 22 each group) with insulin resistance, as defined by McAuley's index²⁸. The provided DGA diet was notably different in fat (less total fat, less saturated, more mono- and poly-unsaturated with more omega-3's) than the TAD diet²⁹. Here, we present our observations of metabolic rate parameters including fat oxidation and metabolic flexibility (calculated as the difference between measured fasting and postprandial fat oxidation³⁰) in response to high-fat

mixed macronutrient meal challenge tests (MMCT) conducted thrice during this diet intervention. In addition, fasting and postprandial lipidomic and metabolomic parameters in response to the MMCT were also measured. Our aims were to (a) identify if there was any effect of the diet intervention on metabolic rate, substrate oxidation and metabolic flexibility in response to the MMCT, and (b) glean further metabolic insight into fat oxidation and metabolic flexibility in women with insulin resistance using omic measures. As our intervention was designed to maintain body weight and physical activity during the testing period, we anticipated that our dietary intervention would have no effect on metabolic flexibility, especially given our small sample size³¹⁻³³. Instead, we explored the metabolic rate and substrate oxidation responses to the repeated MMCT, with molecular input from lipidomic and metabolomic measures. To our knowledge, this is the first report to include three repeats of the same MMCT across weeks, and with metabolomic and lipidomic measures to enable identification of “stable” substrate oxidation and metabolic flexibility characteristics in individuals. By leveraging these metabolomic and lipidomic measures, we hope to enhance our understanding of this population’s underlying metabolic response to a high fat challenge test, thereby identifying potential biomarkers or metabolic signatures for further exploration.

Methods

Study design and participants. To test the impact of diets meeting the Dietary Guidelines for Americans on cardiometabolic risk factors, the individual Metabolism and Physiological Signatures Study (iMAPS; ClinicalTrials.gov: NCT02298725) recruited women who were aged 20–65 y with BMIs between 25–39.9 kg/m². In addition, physical activity was limited to < 150 min/week, and they were insulin resistant based on screening tests to calculate McAuley’s insulin sensitivity index (values > 5.8 was considered insulin resistant³⁴)²⁷. Other inclusion criteria included resting blood pressure ≤ 140/90 mm Hg, impaired glucose homeostasis and/or elevated fasting TGs, maintenance of a sedentary lifestyle with activity monitoring for 7-day periods four times over the study period using waist-worn accelerometers (Respironics® Actical™; Philips North America Co, Cambridge MA). Body composition was determined at 0 and 8 weeks by dual-energy X-ray absorptiometry (DXA; Hologic Discovery QDR Series 84994; Hologic, Inc.). A consort diagram describing participants is shown in **Supplemental Figure 1**.

Mixed macronutrient challenge test. A high fat mixed macronutrient challenge vehicle contained 840 Kcals with 60% energy (calories) from fat, 28% from carbohydrates and 12% from protein was used in the mixed macronutrient challenge test (MMCT). The MMCT protocol was administered at week 0 prior to intervention, at intervention week 2, and intervention week 8 as reported³⁵. On the evening before the test day, participants consumed a provided standardized pretest dinner and began a 12h fast. The following morning, after obtaining a fasting blood sample and measuring resting metabolic rate, the MMCT ‘milkshake-like’ meal was provided,

and participants were given 10 minutes to consume it. After weighing back the residual left in the container, participants consumed 56 ± 2 g of palm oil, 59 ± 2 g of sucrose and 26 ± 1 g of egg white protein. The fatty acid composition by weight of the total fat content was 43% palmitate (16:0), 40% oleate (C18:1n9), 9% linoleate (18:2n6), 4% stearate (18:0), and < 1 % other detected fatty acid residues. The protocol included collecting four blood samples at 0 (fasting), and 0.5, 3 and 6h post meal challenge.

Indirect calorimetry and metabolic flexibility. Estimates of fuel utilization were generated based on indirect calorimetry measures using automated metabolic carts with an open circuit system (TrueOne 2400, ParvoMedics). Measurements were taken four times for intervals of 15-20 min; the time sequence was fasting 0h, consumption of challenge meal, then 0.75-1h, 3h, and 6h after the MMCT, closely coinciding with blood collection times. Respiratory exchange ratio (RER), a common indicator of carbohydrate vs. fat combustion, was calculated as the ratio of measured volume of carbon dioxide ($\dot{V}CO_2$) produced to volume of oxygen ($\dot{V}O_2$) consumed using the equation $\dot{V}CO_2/\dot{V}O_2$. The resting and postprandial energy expenditure (EE) were estimated using the Weir equation without urinary nitrogen correction: $EE = [(3.94 \times \dot{V}O_2) + (1.1 \times \dot{V}CO_2)]^{36}$. Rates of fat oxidation and carbohydrate oxidation were estimated using the Frayn equations³⁷. A urinary nitrogen correction was used with these equations based on the protein content of the controlled diet, assuming participants were in nitrogen balance²⁹. As mentioned earlier, metabolic flexibility was calculated as the change in postprandial fat oxidation compared to fasting, in response to the MMCT (i.e. postprandial – fasting fat oxidation)³⁰. In addition, we also calculated % change in fat oxidation postprandial compared to fasting and change and % change in RER postprandial compared to fasting at all weeks.

Plasma fatty acid analysis. Fatty acids were isolated in the presence of internal standards and quantified by gas chromatography-mass spectrometry as fatty acid methyl esters (FAMES) against authentic calibration standards. Samples were processed in a total of 12 batches, each containing blanks, replicates, and laboratory reference materials. Samples were prepared using standard extraction and derivatization methods which are explained in greater detail in the supplemental methods.

Except for the non-esterified fatty acids (NEFA), surrogate recoveries, replicate precision and blank levels were acceptable. For NEFA, a subset of saturated fatty acids were compromised and excluded from the analysis. Subtle batch specific differences in NEFA were removed by adjusting samples' means by laboratory reference material batch averages.

Kit based-targeted metabolomics. Plasma concentrations of acyl carnitines (n = 40), amino acids (n = 21), biogenic amines (n = 21), glycerophospholipids (n = 90), sphingomyelins (n = 15) and total hexoses were measured using AbsoluteIDQ® p180 kits (Biocrates Life Sciences, Innsbruck, Austria). Samples were prepared and data collected by UPLC tandem mass spectrometry on an API 6500 (Sciex, Framingham, MA) as per manufacturer's instructions.

Fat burner classification. The propensity of an individual to metabolically combust (i.e. burn) ingested fat was quantified from the MMCT indirect calorimetry data as both continuous and categorical variables at baseline, 2 and 8 weeks. The continuous variable, % fat burned (%FB), was determined with the following equation:

$$\% \text{ fat burned} = \frac{\text{fat oxidized}}{\text{fat consumed}} * 100$$

where,

fat oxidized = Frayn equation calculated sum of the 6h PP g fat oxidized

fat consumed = MMCT g fat ingested

Participants were classified based on the magnitude and stability of the % FB: high fat burners (HB) > 40% at all weeks (n = 6); low-fat burners (LB) < 30% at all weeks (n = 7); variable burners (VB) = a mix of HB and LB classification at different weeks (n = 31).

Calculation of appearance, disappearance, and area under the curve. To quantify the changes in metabolites from fasting to postprandial, a one-compartment oral bolus pharmacokinetic model non-linear curve fit was applied to all lipidomic and metabolomic data to estimate an “appearance” rate (i.e. rate at which the metabolite appeared in plasma), a “disappearance” rate (i.e. rate at which the metabolite disappeared from plasma) and an area under the curve (See **Supplemental Figure 2**). Physiologically, these could indicate changes happening in early postprandial (appearance) and late postprandial (disappearance) states. These data were used in subsequent analyses as described below.

Statistical tools. An overview of the analysis performed is presented in **Supplemental Figure 3**. All analyses were performed in JMP Pro 17.2 (SAS institute, Cary, NC) or R Statistical software³⁸, unless otherwise specified. Data were evaluated for distribution (normality) using Q-Q plots and Shapiro Wilk tests, followed by transformations, if necessary, using the Johnson family of normalizations. Data were evaluated for missingness using the Amelia package in R³⁹, and 3% of data were found to be missing. Missing data were imputed using singular value

decomposition (SVD) imputation in JMP Pro 16.1, after careful evaluation with several imputation tools (see **Supplemental Figure 4**).

Univariate statistics. Diet effects on parameter means were tested by analysis of covariance (ANCOVA) with the baseline values (week 0) used as a covariate. Energetic parameters tested included resting metabolic rate (RMR), respiratory exchange ratio (RER), substrate oxidation (carbohydrate and fat oxidation), and metabolic flexibility. Here, we calculated metabolic flexibility as the change in RER and fat oxidation between postprandial and fasting measures (360-0, 180-0, 30-0 mins). To evaluate the effect of the intervention on energy substrate parameters, a repeated measures mixed model was used with diet group, week and time as fixed effects, treatment group and week as an interaction and participant as a random effect, with week as the repeated measure. A two-one sided equivalence test (TOSTER package in R) was used to ensure that metabolic flexibility from the two intervention groups could be combined into a single population for analysis. This analysis tests whether an intervention had a statistically measurable impact on the primary outcome (RER) which is also clinically or physiologically relevant.

FB group differences in postprandial measures of metabolic flexibility, body composition, and circulating lipids over the course of the study were assessed using mixed models. These models included FB-group, week, time (mins) and interaction (FB-group*week) as fixed effects, and participant as a random effect with week as the repeated measure. Area under the curve was calculated based on Simpson's rule⁴¹ and Kruskal-Wallis non-parametric tests were used to compare the AUCs between FB-groups by week.

Multivariate statistics. To enhance interpretation of this highly dimensional data, metabolomic/lipidomic data (~230 parameters) were subjected to variable clustering using an implementation of the VARCLUS algorithm in JMP Pro v 17.2.0. To highlight the metabolomic features of interest associated with the fat burner group, partial least squares (PLS) analyses were performed using the cluster component scores and either as (a) continuous %FB or (b) as %FB categorical extremes (i.e. HB vs LB) as outcome variables. The non-linear iterative partial least squares (NIPALS) algorithm was used with leave-one-out cross validation to select the number of factors that minimize the Root Mean PRESS statistic. The Q² (goodness of prediction statistic) and R² (coefficient of multiple determination) for independent and dependent variables were used to evaluate the model fit. Cluster components with a variable importance in projection (VIP) score of > 1 were identified and interpreted as significant explanatory features for the %FB. In addition to running a PLS-DA, since we did the MMCT three times (week 0, 2 and 8), the mixOmics package in R was used to do a repeated measures (multi-level) sparse PLS-DA (sPLS-DA) to extract the loadings, scores, and VIP variables, to compare with the model developed by the NIPALS algorithm. Only VIP variables identified by both approaches were used for final interpretation.

Results

Forty- four women who were overweight or obese with insulin resistance were included in this study. Participant profiles have previously been published ²⁷. Briefly, at baseline the mean participant age was 47.1 ± 9.5 , range 21-64 y and mean BMI 32.4 ± 3.9 , range 25.2- 39.8 kg/m² ²⁷. Intervention groups were well matched by age and anthropomorphic data, and diet-dependent changes in body characteristics and metabolic rate measures were not observed (**Supplemental Table 1**)²⁷. Linear mixed models identified no significant differences between diet groups in fat oxidation ($p=0.47$), carbohydrate oxidation ($p = 0.53$), RMR ($p = 0.77$), RER ($p = 0.50$), % change in fat oxidation from fasting ($p = 0.46$) and % change in RER from fasting ($p = 0.72$) from minute 0 (before the MMCT) to minute 60, 180 or 360 (see **Figure 1**).

Since diet intervention groups were not statistically different in energy and substrate metabolic parameters and metabolic flexibility, this was further evaluated using equivalence tests to justify combining the intervention groups. The difference between RER from minute 0 (before the MMCT) to minute 360 (6 hours after the test) was used as a primary outcome to evaluate the equivalence of groups. A range of deltas for RER between ± 0.1 to ± 0.01 were tested as a change of ± 0.01 constituting a $< \sim 3\%$ change in RER which was deemed clinically irrelevant based on the American Heart Association report suggesting a 3% within participant measurement variability ⁴⁰. In our study, the maximum measured change in RER between the fasting and postprandial states was 0.294. Results for this test (**Supplemental Figure 5**) concluded that the changes in RER were significantly similar up to our pre-determined delta, and we could proceed in combining the diet groups and assessing all the data together. While this supplemental figure only shows these relationships for week 8, week 0 and week 2 were also tested with identical results.

Participants were classified into fat-burning groups based on the % of consumed fat from the MMCT burned over the six hours post-test. High burners (HB) burned 40% or more of the ingested fat, while low burners (LB) burned less than 30%. As shown in **Figure 2 panel A**, the VB participants showed inconsistent segregation into a low or high burner group. These characteristics needed to be consistent over the three test weeks of the study to be classified into either group. It is important to note that fat oxidation, RMR and RER were significantly different between burner groups, with HB consistently burning more fat than LB. However, %change from fasting in fat oxidation and RER showed no significant differences between the groups using linear mixed models (Figure 2 panel B). Upon visual inspection, however, it appears that the HB group reduces fat burned as a change from baseline compared to LB at weeks 0 and 8. **Supplemental Figure 6** shows % change from fasting in fat oxidation and RER only in HB and LB groups, where linear mixed model identified a significantly lower % change from fasting in fat oxidation at 60min postprandial in HB compared to LB ($p = 0.019$).

A summary of clinical outcome variables by fat burner group at baseline and the end of the intervention is presented in **Figure 3**. When separated by %FB group, there were no statistically significant differences in insulin ($p = 0.06$), glucose ($p = 0.61$), TG ($p = 0.55$), HDL ($p = 0.11$), LDL ($p = 0.52$), and total cholesterol ($p = 0.80$) at baseline between HB and LB and this largely persisted through week 8. However, HDLc AUC was significantly lower in HB compared to LB at weeks 2 and 8 ($p = 0.03$ and 0.01 respectively). Further, while the time course for triglycerides appears visually different between LB and HB in the late postprandial phase, we were not powered to detect the difference statistically. There were significant differences between HB and LB in BMI (**Table 1**) at week 0 ($p = 0.023$) which persisted through the end of the intervention (week 8 $p = 0.023$). The same was true for total mass (week 0: $p = 0.023$, week

8: 0.023), lean mass ($p = 0.023, 0.023$), fat mass ($p = 0.023, 0.023$), but not waist to hip ratio ($p = 0.49, 0.48$) or McAuley's ISI ($p = 0.749, 0.886$). There were also no significant group differences for % android fat ($p = 0.098, 0.098$), % gynoid fat ($p = 0.48, 0.86$), or age ($p = 0.098, 0.098$). Menopausal status was also not significantly different between the two groups, evaluated by Fishers exact tests ($p = 0.29$).

Variable Clustering dimension reduction

The metabolomic and lipidomic data included 236 variables (not including anthropometric, clinical, and metabolic variables). To better equip our analysis tools to detect metabolite predictors capable of differentiating between the FB groups, we used a dimension reduction algorithm that clusters variables. This dimension reduction tool generates components (like principal components analysis) that are a linear combination of variables. Based on these components, variables are placed into clusters of 'similar' variables (cluster components), such that the first cluster component (eigen vectors) within each cluster captures the most variance amongst those variables. In subsequent analyses, the cluster component scores for appearance, disappearance and area under the curve were used as independent predictors to identify differences between the fat burner group metabolic signatures. For appearance rates, AUC, and disappearance rates, clusters of 32, 31 and 36 variables were identified respectively.

Lipidome predictors of fat oxidation and metabolic flexibility

In our efforts to identify metabolites that were most predictive of %FB groups, only the disappearance rate cluster components (i.e. the 36 variable cluster components that were generated using the variable cluster algorithm in the previous step) resulted in a converged

model. **Table 2** lists select clusters and their component metabolites, their corresponding cluster identifier numbers, and eigen vectors (cluster components). Since the eigenvector directions (positive and negative) only translate to the scaled transformations to achieve the clustering, both positive and negative eigen vectors will be interpreted as positive integers in the next steps. The PLSDA models did not converge when the VB were included as an intermediate ordinal group, nor did they converge when we used PLSR to predict % FB as a continuous variable. Thus, our final PLSDA model only compared HB vs LB groups and converged with 2 minimizing factors which explained 17.7% of variation in X and 71.6% of variation in Y variables with a Q^2 of 0.20 (See **Figure 4**). While the Q^2 of 0.20 does not suggest complete discrimination between groups, our sPLSDA efforts showed strikingly similar outcomes, and scores and loadings values, indicating the robustness of identified differences. The scores plot of the PLSDA, inset into the loadings plot, shows only a small overlap between HB (in red) and LB (in blue) groups. The loadings plot, which depicts the corresponding cluster components shows which variables brought about the separation. Clusters with $VIP > 1$ are semi-synonymous to those with p-values < 0.05 , and therefore play a significant role in differentiating the HB and LB group. In addition, the results from the sPLSDA are presented in **Supplemental Figure 7** and show very similar separation of participants by burner classification, and VIP variables, with a total of 17% of X was explained by the first two loadings, like the PLSDA using NIPALS algorithm.

Table 2 lists all the VIP variables from the PLSDA and sPLSDA analyses. Only the commonly identified cluster components from both analyses will be used for interpretation. As shown in Figure 4 panel B, cluster components 10, 11, 22, 23 and 30 were predictors of the LB group, while components 6, 14, 16 and 17 were predictors of the HB group.

288 ***LB group metabolites:*** As seen in Table 2, cluster component 10 includes higher (faster)
 289 disappearance rates from plasma of total and non-esterified fatty acid ratios TFA C18:0/16:0,
 290 TFAC16:1n7/C16:0 (C16/18 elongase and Δ 9-desaturase), NEFA C20:4n6/C20:3n6 and
 291 TFAC18:1n9/C18:0. Cluster 11 includes lysophosphatidylcholines (LPC) with long chain
 292 saturated, monounsaturated, and polyunsaturated fatty acids. Cluster 22 and 23 are a mixture of
 293 acylcarnitines (acetylcarnitine and methylmalonylcarnitine), acetylmethionine, and phospholipids
 294 (PC.aa.C38.1). Cluster 30 is a mixture of acylcarnitines (octadecadienylcarnitine),
 295 sphingomyelins and alkyl-ethyl phosphatidylcholines.
 296

297 ***HB group metabolites:*** Cluster 6 includes higher (faster) disappearance rates from plasma of
 298 total NEFA saturated and monounsaturated fatty acids and the ratio of C16:1n9/C16. Cluster 14
 299 and 16 include all non-essential amino acids, while cluster 17 includes non-esterified fatty acid
 300 ratios (C18:0/C16:0, C22:4n6/C20:4n6 and C18:1n9/C18:0).

Discussion

In the current study, as expected, there was no differential effect of the diet interventions on our chosen measures of metabolic flexibility. Further, we identified individuals who were consistently different in how they handled the influx of energy substrates from the MMCT i.e. high fat burners (HB) and low-fat burners (LB). Women in the HB group consistently oxidized >40% of consumed fat, while the LB group consistently oxidized < 30% of consumed fat. Based on what is understood about fat oxidation, metabolic flexibility, and insulin resistance, overall low-fat burning is associated with reduced metabolic flexibility². However, in our cohort, women who burned less fat at fasting (LB group) switched to burning more fat during the 6h postprandial, and in the immediate postprandial phase after consuming the MMCT (with 60% fat, 28% carbohydrates and 12% protein) compared to the HB group. Our sample size was likely too small to detect this at week 2 and week 8, but did identify this difference at week 0, and can be visually seen at week 8. Further, the LB group had lower fat mass, lean mass, BMI and higher HDL compared to the HB group. However, proportions of lean and fat mass were not significantly different between groups. This suggests that in cases of high BMI and insulin resistance, matched proportional lean mass does not “rescue” the effect higher fat mass has on metabolic flexibility⁴². While women in the LB group were older and more postmenopausal, the HB group was not homogenously younger or premenopausal. While metabolic flexibility differences at week 0 were observed in the early postprandial phase, metabolomic and lipidomic profiles identified higher late-postprandial disappearance rates of lysophosphatidylcholines and acylcarnitines in LB group, both implying better metabolic health.

Women in the DGA group were given more whole fruits, vegetables, whole grains, seafood, and nuts, whereas women in the TAD group were given more refined grains, meat, and solid fats over the 8-week feeding intervention²⁷. As predicted, there was no impact of this diet

on metabolic flexibility, measured by RER or change in fat oxidation between fasting and postprandial time points. Similarly, Kardinaal et al.⁴³ evaluated RER changes in a group of healthy males given a high-fat meal over 4 weeks and did not see any changes in their RER. Likewise, in Fechner et al, a group of males and females randomized to six weeks of either a healthy diet or western diet, also did not experience changes in insulin sensitivity or RQ when measured by high fat meal challenge test¹⁹.

In the late postprandial phase, there was an increased rate of disappearance of Lysophosphatidylcholine (LPCs) and phosphatidylcholine (PCs) in the LB group of women when compared to HB. LPCs are phosphatidylcholines that have been cleaved by a phospholipase (see **Figure 5**), and more commonly function as lipid mediators⁴⁴. The composition of both PC and LPC are primarily saturated fatty acids (SFA) or monounsaturated fatty acids (MUFA) with varying degrees of length and position of unsaturated double bonds. While PCs are found in membranes in large quantities, higher concentrations of LPC have been associated with atherosclerosis through disruptions of mitochondrial integrity⁴⁴, while lower circulating concentrations are found in prediabetes and type 2 diabetes⁴⁵ particularly in the skeletal muscle⁴⁶. LPCs are known to inhibit hepatic fatty acid oxidation (among several other effects)⁴⁷. Hence, their clearance would have supported maximal fat oxidation in LB. LB women could have reduced plasma phospholipase2 (PLA2) activity, which generates LPC from PC. Higher PLA2 activity has been observed to be proatherogenic, irrespective of whether this is in plasma or in the endothelium⁴⁸. So, this higher rate of disappearance of LPC could be a largely metabolically favorable observation. Further, this suggests that relatively quicker disappearance of plasma LPC's following a high fat meal could be indicative of higher metabolic flexibility.

LB women also displayed a faster rate of disappearance of acylcarnitines from plasma. Higher circulating concentrations of acetylcarnitine have been reported in people with prediabetes and type 2 diabetes⁴⁹. Also, increased circulating octadecadienoylcarnitine has been associated with increased all-cause mortality and hospitalizations in heart failure patients⁴⁹. Together, these would suggest that a quicker disappearance of from plasma in LB women would lead to a metabolically healthier milieu compared to women in HB group.

In the late postprandial phase, especially following a meal that has carbohydrates in it (which our MMCT did), the initial rise followed by the post-absorptive drop in insulin concentration will result in an increase in adipose lipolysis and higher circulating NEFA concentrations⁵⁰. In this phase, the LB group had a reduction in $\Delta 9$ desaturase, $\Delta 5$ desaturase and C16/18 elongase activities affecting the complex lipid pool (fatty acids attached to triglycerides) likely resulting in more C16:0 and C18:0 acyl glycerol concentrations in plasma. Saturated TGs, such as C16:0 and C18:0, have been shown to suppress LDL-receptor activity, thereby increasing plasma LDL-c⁵¹, increasing cardiovascular disease risk. Thus far, this is the only less-than-ideal metabolic milieu that has been noted in women in the LB group.

Our initial assumption was that those who could burn a higher percentage of fat following the challenge meal would be more metabolically flexible, as they would demonstrate an effective switch of primary metabolic fuel sources, potentially leading to health benefits. However, our findings in this cohort do not support this idea, suggesting a more complex relationship between fat oxidation and metabolic flexibility. HB women had higher body mass and were in the obese BMI category compared to the LB group who were in the overweight BMI category. Obese insulin resistant individuals are known to have higher systemic fasting respiratory quotient^{2,52} but this is not true in our current cohort. Metabolic inflexibility to a high fat meal has been shown to

be a predictor for subsequent weight gain¹⁷, suggesting our HB group women are metabolically inclined to further gain body weight. Differences in their underlying physiology detected by omic analysis, such as faster clearance from circulation of LPCs and acylcarnitines to offset the higher saturated triglycerides, also support our observation that LB women may be on the “healthier” metabolic spectrum compared to HB women. Given our findings of higher fat burning not equaling higher metabolic flexibility, the definition of metabolic flexibility may need to be expanded beyond the ability to burn fat in a high-fat challenge meal. A follow up study that evaluates how much fat oxidation and fuel switching occurs following consumption of meals with serial increasing % dietary fat (0 – 100%) in a diverse population could shed further light on this relationship.

Strengths and Limitations

This investigation used a standard meal challenge test for all the women and repeated it three times, which affords robustness to our conclusions. Participants were categorized into burner groups based on consistent responses in all three challenge events, reducing the likelihood of measurement errors driving our sub-group determination. All participants were relatively sedentary, did not change physical activity during the intervention, and were given a pre-test dinner the night prior to consuming the MMCT to reduce variability from pre-meal dietary sources. Evaluating equivalence and robust approaches to data analyses (such as including a repeated measures PLSDA) also strengthen our interpretations. However, there were several limitations that must be acknowledged. Our categorized burner groups were small (HB, n = 6 and LB, n = 7). This small sample size may have been the reason why only one model PLSDA converged among several we tested. It is possible that the difference between the high fat burners and low-fat burners may be due to a genetic predisposition to be obligatory fat or carbohydrate

burners, rather than the flexibility or inflexibility of their metabolism. However, the study lacks SNP/genome sequence data that could corroborate this. The high inter-test and inter-individual variability in our VB group (n = 31, ~70% of participants) unfortunately rendered the group empirically inexplicable. The individuals in the VB group, however, are likely to be the majority in a population, highlighting the importance of being able to understand their underlying physiology. Future studies must use larger sample sizes to be able to overcome this failing. While we controlled the participants' food intake and physical activity, we did not control or account for changes in sleep, endocrine factors, stress, gut microbiome etc. which could have affected one of the three weeks' MMCT responses of women in the VB group.

Conclusion

In summary, we identified that a high fat burning capacity is unlikely to equal higher metabolic flexibility universally in women with insulin resistance. In addition, the HB group seemed to reap no clear metabolic or clinical rewards from this high burning capacity. Currently, there is a lack of consensus on the definition and standard metric for metabolic flexibility, particularly in the way it relates to dietary fat-induced metabolic flexibility. To fully understand the response of the metabolism to gauge flexibility, a mixed macronutrient challenge test offers more information than a standard oral glucose tolerance test. Additionally, we also highlighted the important inclusion of metabolomics, allowing investigators to “peer under the hood” of an individual's metabolism, corroborating results observed both clinically and physiologically. This may be especially true in insulin resistant individuals, as the relationship between insulin and lipid metabolism is complicated by the inherent involvement of body composition, dietary macro-nutrient composition, and several other endocrine and molecular mechanisms at play. As mentioned by other researchers, a comprehensive definition of metabolic flexibility to lipids is sorely needed, especially given the results mentioned above.

417

418 **Acknowledgements**

419 **Author Contributions**

420 Conceptualization SK, NLK, JWN; Methodology SK, NLK, JWN; Formal Analysis SK, VMA,
421 MMA; Investigation NLK, SK, VMA, MMA; Data Curation SK; Writing – Original Draft
422 VMA, SK, MMA; Writing – Review & Editing SK, NLM, JWN, MMA; Visualization SK,
423 MMA; Supervision SK; Funding Acquisition SK, NLK, JWN.

424 Declaration of Interests: The authors declare no competing interests.

425

426 Declaration of generative AI and AI-assisted technologies: SCITE and Elicit were used to search
427 for references to ensure comprehensive literature review to use in the writing of this manuscript.

428 However, no AI-assisted text was incorporated, and all writing was done by the authors.

429

430 Supplemental Information:

431 Document S1. Figures S1- S7 and Tables S1

432

References

1. Smith, R.L., Soeters, M.R., Wust, R.C.I., and Houtkooper, R.H. (2018). Metabolic Flexibility as an Adaptation to Energy Resources and Requirements in Health and Disease. *Endocr Rev* 39, 489-517. 10.1210/er.2017-00211.
2. Galgani, J.E., Moro, C., and Ravussin, E. (2008). Metabolic flexibility and insulin resistance. *Am J Physiol Endocrinol Metab* 295, E1009-1017. 10.1152/ajpendo.90558.2008.
3. Goodpaster, B.H., and Sparks, L.M. (2017). Metabolic Flexibility in Health and Disease. *Cell Metab* 25, 1027-1036. 10.1016/j.cmet.2017.04.015.
4. Kelley, D.E., and Mandarino, L.J. (2000). Fuel selection in human skeletal muscle in insulin resistance: a reexamination. *Diabetes* 49, 677-683. 10.2337/diabetes.49.5.677.
5. Ukropcova, B., McNeil, M., Sereda, O., de Jonge, L., Xie, H., Bray, G.A., and Smith, S.R. (2005). Dynamic changes in fat oxidation in human primary myocytes mirror metabolic characteristics of the donor. *J Clin Invest* 115, 1934-1941. 10.1172/JCI24332.
6. Robinson, S.L., Chambers, E.S., Fletcher, G., and Wallis, G.A. (2016). Lipolytic Markers, Insulin and Resting Fat Oxidation are Associated with Maximal Fat Oxidation. *Int J Sports Med* 37, 607-613. 10.1055/s-0042-100291.
7. Chakravarthy, M.V., and Neuschwander-Tetri, B.A. (2020). The metabolic basis of nonalcoholic steatohepatitis. *Endocrinol Diabetes Metab* 3, e00112. 10.1002/edm2.112.
8. van den Broek, T.J., Bakker, G.C.M., Rubingh, C.M., Bijlsma, S., Stroeve, J.H.M., van Ommen, B., van Erk, M.J., and Wopereis, S. (2017). Ranges of phenotypic flexibility in healthy subjects. *Genes Nutr* 12, 32. 10.1186/s12263-017-0589-8.
9. Muoio, D.M. (2014). Metabolic inflexibility: when mitochondrial indecision leads to metabolic gridlock. *Cell* 159, 1253-1262. 10.1016/j.cell.2014.11.034.
10. York, L.W., Puthalapattu, S., and Wu, G.Y. (2009). Nonalcoholic fatty liver disease and low-carbohydrate diets. *Annu Rev Nutr* 29, 365-379. 10.1146/annurev-nutr-070208-114232.
11. Gunaseelan, L., Khan, U.S., Khalid, F., and Hamid, M.A. (2021). Non-alcoholic Fatty Liver Disease and Carbohydrate Restricted Diets: A Case Report and Literature Review. *Cureus* 13, e18641. 10.7759/cureus.18641.
12. Soares, M.J., Cummings, S.J., Mamo, J.C., Kenrick, M., and Piers, L.S. (2004). The acute effects of olive oil v. cream on postprandial thermogenesis and substrate oxidation in postmenopausal women. *Br J Nutr* 91, 245-252. 10.1079/BJN20031047.
13. Rynders, C.A., Blanc, S., DeJong, N., Bessesen, D.H., and Bergouignan, A. (2018). Sedentary behaviour is a key determinant of metabolic inflexibility. *J Physiol* 596, 1319-1330. 10.1113/JP273282.
14. Shi, W., Hegeman, M.A., van Dartel, D.A.M., Tang, J., Suarez, M., Swarts, H., van der Hee, B., Arola, L., and Keijer, J. (2017). Effects of a wide range of dietary nicotinamide riboside (NR) concentrations on metabolic flexibility and white adipose tissue (WAT) of mice fed a mildly obesogenic diet. *Mol Nutr Food Res* 61. 10.1002/mnfr.201600878.
15. van Dijk, S.J., Mensink, M., Esser, D., Feskens, E.J., Muller, M., and Afman, L.A. (2012). Responses to high-fat challenges varying in fat type in subjects with different metabolic risk phenotypes: a randomized trial. *PLoS One* 7, e41388. 10.1371/journal.pone.0041388.
16. Yu, E.A., Le, N.A., and Stein, A.D. (2021). Measuring Postprandial Metabolic Flexibility to Assess Metabolic Health and Disease. *J Nutr* 151, 3284-3291. 10.1093/jn/nxab263.
17. Begaye, B., Vinales, K.L., Hollstein, T., Ando, T., Walter, M., Bogardus, C., Krakoff, J., and Piaggi, P. (2020). Impaired Metabolic Flexibility to High-Fat Overfeeding Predicts Future Weight Gain in Healthy Adults. *Diabetes* 69, 181-192. 10.2337/db19-0719.

18. Nagai, N., Sakane, N., and Moritani, T. (2005). Metabolic responses to high-fat or low-fat meals and association with sympathetic nervous system activity in healthy young men. *J Nutr Sci Vitaminol (Tokyo)* 51, 355-360. 10.3177/jnsv.51.355.
19. Fechner, E., Bilet, L., Peters, H.P.F., Hiemstra, H., Jacobs, D.M., Op 't Eyndt, C., Kornips, E., Mensink, R.P., and Schrauwen, P. (2020). Effects of a whole diet approach on metabolic flexibility, insulin sensitivity and postprandial glucose responses in overweight and obese adults - A randomized controlled trial. *Clin Nutr* 39, 2734-2742. 10.1016/j.clnu.2019.12.010.
20. Jones, P.J., Jew, S., and AbuMweis, S. (2008). The effect of dietary oleic, linoleic, and linolenic acids on fat oxidation and energy expenditure in healthy men. *Metabolism* 57, 1198-1203. 10.1016/j.metabol.2008.04.012.
21. Casas-Agustench, P., Lopez-Uriarte, P., Bullo, M., Ros, E., Gomez-Flores, A., and Salas-Salvado, J. (2009). Acute effects of three high-fat meals with different fat saturations on energy expenditure, substrate oxidation and satiety. *Clin Nutr* 28, 39-45. 10.1016/j.clnu.2008.10.008.
22. Piers, L.S., Walker, K.Z., Stoney, R.M., Soares, M.J., and O'Dea, K. (2002). The influence of the type of dietary fat on postprandial fat oxidation rates: monounsaturated (olive oil) vs saturated fat (cream). *Int J Obes Relat Metab Disord* 26, 814-821. 10.1038/sj.ijo.0801993.
23. Jones, P.J., Pencharz, P.B., and Clandinin, M.T. (1985). Absorption of ¹³C-labeled stearic, oleic, and linoleic acids in humans: application to breath tests. *J Lab Clin Med* 105, 647-652.
24. DeLany, J.P., Windhauser, M.M., Champagne, C.M., and Bray, G.A. (2000). Differential oxidation of individual dietary fatty acids in humans. *Am J Clin Nutr* 72, 905-911. 10.1093/ajcn/72.4.905.
25. Miao, H., Li, B., Wang, Z., Mu, J., Tian, Y., Jiang, B., Zhang, S., Gong, X., Shui, G., and Lam, S.M. (2022). Lipidome Atlas of the Developing Heart Uncovers Dynamic Membrane Lipid Attributes Underlying Cardiac Structural and Metabolic Maturation. *Research* 2022. 10.34133/research.0006.
26. Ghosh, A., and Nishtala, K. (2017). Biofluid lipidome: a source for potential diagnostic biomarkers. *Clin Transl Med* 6, 22. 10.1186/s40169-017-0152-7.
27. Krishnan, S., Adams, S.H., Allen, L.H., Laugero, K.D., Newman, J.W., Stephensen, C.B., Burnett, D.J., Witbracht, M., Welch, L.C., Que, E.S., and Keim, N.L. (2018). A randomized controlled-feeding trial based on the Dietary Guidelines for Americans on cardiometabolic health indexes. *Am J Clin Nutr* 108, 266-278. 10.1093/ajcn/nqy113.
28. McAuley, K.A., Williams, S.M., Mann, J.I., Walker, R.J., Lewis-Barned, N.J., Temple, L.A., and Duncan, A.W. (2001). Diagnosing insulin resistance in the general population. *Diabetes Care* 24, 460-464. 10.2337/diacare.24.3.460.
29. Krishnan, S., Lee, F., Burnett, D.J., Kan, A., Bonnel, E.L., Allen, L.H., Adams, S.H., and Keim, N.L. (2020). Challenges in Designing and Delivering Diets and Assessing Adherence: A Randomized Controlled Trial Evaluating the 2010 Dietary Guidelines for Americans. *Curr Dev Nutr* 4, nzaa022. 10.1093/cdn/nzaa022.
30. He, S., Le, N.A., Ramirez-Zea, M., Martorell, R., Narayan, K.M.V., and Stein, A.D. (2021). Metabolic flexibility differs by body composition in adults. *Clin Nutr ESPEN* 46, 372-379. 10.1016/j.clnesp.2021.09.730.
31. Lepine, G., Tremblay-Franco, M., Boudier, S., Dimina, L., Fouillet, H., Mariotti, F., and Polakof, S. (2022). Investigating the Postprandial Metabolome after Challenge Tests to Assess Metabolic Flexibility and Dysregulations Associated with Cardiometabolic Diseases. *Nutrients* 14. 10.3390/nu14030472.
32. Weinisch, P., Fiamoncini, J., Schraner, D., Raffler, J., Skurk, T., Rist, M.J., Romisch-Margl, W., Prehn, C., Adamski, J., Hauner, H., et al. (2022). Dynamic patterns of postprandial metabolic responses to three dietary challenges. *Front Nutr* 9, 933526. 10.3389/fnut.2022.933526.

33. LaBarre, J.L., Singer, K., and Burant, C.F. (2021). Advantages of Studying the Metabolome in Response to Mixed-Macronutrient Challenges and Suggestions for Future Research Designs. *J Nutr* 151, 2868-2881. 10.1093/jn/nxab223.
34. Gutch, M., Kumar, S., Razi, S.M., Gupta, K.K., and Gupta, A. (2015). Assessment of insulin sensitivity/resistance. *Indian J Endocrinol Metab* 19, 160-164. 10.4103/2230-8210.146874.
35. Newman, J.W., Krishnan, S., Borkowski, K., Adams, S.H., Stephensen, C.B., and Keim, N.L. (2022). Assessing Insulin Sensitivity and Postprandial Triglyceridemic Response Phenotypes With a Mixed Macronutrient Tolerance Test. *Front Nutr* 9, 877696. 10.3389/fnut.2022.877696.
36. Weir, J.B. (1949). New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol* 109, 1-9. 10.1113/jphysiol.1949.sp004363.
37. Frayn, K.N. (1983). Calculation of substrate oxidation rates in vivo from gaseous exchange. *J Appl Physiol Respir Environ Exerc Physiol* 55, 628-634. 10.1152/jappl.1983.55.2.628.
38. Team, R.C. (2021). A language and environment for statistical computing.
39. Honaker J, K.G., Blackwell M (2011). Amelia II: A Program for Missing Data. *Journal of Statistical Software* 45, 1 - 47. doi:10.18637/jss.v045.i07.
40. Balady, G.J., Arena, R., Sietsema, K., Myers, J., Coke, L., Fletcher, G.F., Forman, D., Franklin, B., Guazzi, M., Gulati, M., et al. (2010). Clinician's Guide to cardiopulmonary exercise testing in adults: a scientific statement from the American Heart Association. *Circulation* 122, 191-225. 10.1161/CIR.0b013e3181e52e69.
41. Tuah, A.N., Ibrahim, A.B., Dzulkifly, S., Mohammad Yusof, F., Awang Nor, R., & Ariffin, R. (2022). Analysis of the Area Under a Curve (AUC) using C-programming: Trapezium and Simpson rules techniques. *Journal of ICT in Education* 9, 143-153. <https://doi.org/10.37134/jictie.vol9.1.12.2022>.
42. Sparks, L.M., Ukropcova, B., Smith, J., Pasarica, M., Hymel, D., Xie, H., Bray, G.A., Miles, J.M., and Smith, S.R. (2009). Relation of adipose tissue to metabolic flexibility. *Diabetes Res Clin Pract* 83, 32-43. 10.1016/j.diabres.2008.09.052.
43. Kardinaal, A.F., van Erk, M.J., Dutman, A.E., Stroeve, J.H., van de Steeg, E., Bijlsma, S., Kooistra, T., van Ommen, B., and Wopereis, S. (2015). Quantifying phenotypic flexibility as the response to a high-fat challenge test in different states of metabolic health. *FASEB J* 29, 4600-4613. 10.1096/fj.14-269852.
44. Bellot, P., Moia, M.N., Reis, B.Z., Pedrosa, L.F.C., Tasic, L., Barbosa, F., Jr., and Sena-Evangelista, K.C.M. (2023). Are Phosphatidylcholine and Lysophosphatidylcholine Body Levels Potentially Reliable Biomarkers in Obesity? A Review of Human Studies. *Mol Nutr Food Res* 67, e2200568. 10.1002/mnfr.202200568.
45. Barber, M.N., Risis, S., Yang, C., Meikle, P.J., Staples, M., Febbraio, M.A., and Bruce, C.R. (2012). Plasma lysophosphatidylcholine levels are reduced in obesity and type 2 diabetes. *PLoS One* 7, e41456. 10.1371/journal.pone.0041456.
46. Diamanti, K., Cavalli, M., Pan, G., Pereira, M.J., Kumar, C., Skrtic, S., Grabherr, M., Riserus, U., Eriksson, J.W., Komorowski, J., and Wadelius, C. (2019). Intra- and inter-individual metabolic profiling highlights carnitine and lysophosphatidylcholine pathways as key molecular defects in type 2 diabetes. *Sci Rep* 9, 9653. 10.1038/s41598-019-45906-5.
47. Law, S.H., Chan, M.L., Marathe, G.K., Parveen, F., Chen, C.H., and Ke, L.Y. (2019). An Updated Review of Lysophosphatidylcholine Metabolism in Human Diseases. *Int J Mol Sci* 20. 10.3390/ijms20051149.
48. Hurt-Camejo, E., Camejo, G., Peilot, H., Oorni, K., and Kovanen, P. (2001). Phospholipase A(2) in vascular disease. *Circ Res* 89, 298-304. 10.1161/hh1601.095598.
49. Dambrova, M., Makrecka-Kuka, M., Kuka, J., Vilskersts, R., Nordberg, D., Attwood, M.M., Smesny, S., Sen, Z.D., Guo, A.C., Oler, E., et al. (2022). Acylcarnitines: Nomenclature, Biomarkers,

574 Therapeutic Potential, Drug Targets, and Clinical Trials. *Pharmacol Rev* 74, 506-551.
575 10.1124/pharmrev.121.000408.
576 50. Jelic, K., Hallgreen, C.E., and Colding-Jorgensen, M. (2009). A model of NEFA dynamics with focus
577 on the postprandial state. *Ann Biomed Eng* 37, 1897-1909. 10.1007/s10439-009-9738-6.
578 51. Woollett, L.A., Spady, D.K., and Dietschy, J.M. (1992). Saturated and unsaturated fatty acids
579 independently regulate low density lipoprotein receptor activity and production rate. *J Lipid Res*
580 33, 77-88.
581 52. Kelley, D.E., Goodpaster, B., Wing, R.R., and Simoneau, J.A. (1999). Skeletal muscle fatty acid
582 metabolism in association with insulin resistance, obesity, and weight loss. *Am J Physiol* 277,
583 E1130-1141. 10.1152/ajpendo.1999.277.6.E1130.
584




Figure 1. No substantial effect of diet on parameters of metabolic flexibility over the course of the study. Mixed models (with treatment group (TAD or DGA), week, and time as fixed effects, participant as random effect, week as a repeated measure and treatment group by week interaction) were used to evaluate the effect of intervention, and weeks on intervention in (A) fat oxidation rate, (B) carbohydrate oxidation rate, (C) Resting metabolic rate (RMR), (D) respiratory exchange ratio (RER), (E) change in fat oxidation, represented as percentage and (F) change in RER also represented as a percentage in TAD (n = 22) and DGA (n = 22). The p-values inset in the figures indicates the lack of a week effect within that group, and the week x group interaction was not significant in these parameters (data not shown).




Figure 2. Fat burning groups show significant differences in metabolic features: (A) Fat burning groups, high burners (HB), low burners (LB), and variable burners (VB) were determined based on the % of fat from the challenge meal that was burned over the 6 hours under observation. Participants were separated into fat burner groups (HB burned > 40% of consumed fat, LB burned < 30% of consumed fat) if they sustained their burner classification at weeks 0, 2 and 8 of the intervention. Once categorized, variables relating to metabolic function were assessed by group. (B) p-values in figure were derived via mixed model, with fat burning group (HB or LB), fat burning group*time interaction, and time as fixed effects, and participant as a random effect, and time as a repeated measure. The p-values represent group differences in postprandial metabolic response of all FB-groups. HB (n = 6), LB (n = 7), and VB (n = 31).




Figure 3. (Left panel) Clinical measures demonstrated subtle differences in key health indicators between fat burner groups: p-values for clinical time course differences over the intervention period were determined through a mixed model approach with fat burner group, and time as fixed effects, participant as random effect, time as repeated measure, along with FB group and time interaction. Horizontal dashed lines represent what is ‘within normal limits’ (WNL for females based on NCEP-STEPIII guidelines) for each measure. Total cholesterol < 200 mg/dL. HDL-c > 50 mg/dL. LDL-c < 100 mg/dL. Glucose < 100 mg/dL (fasted). Insulin < 1010 pg/mL (fasted). p-values reported are between all groups (HB, LB, and VB). (Right panel) Area under the curve, calculated using Simpsons rule for numerical integration, along with p-values inset, based on non-parametric van der Weerdan’s tests comparing LB vs HB groups.

Table 1. Select anthropometric and clinical measurements at week 0 and week 8 for low fat burner and high fat burner groups.

Variable	Week 0			Week 8		
	LB	HB	p-value	LB	HB	p-value
	mean (SD)	mean (SD)		mean (SD)	mean (SD)	
Age	57.0 (6.4)	42.7 (13.9)	0.980	57.0 (6.4)	42.7 (13.9)	0.980
BMI (kg/m ²)	29.26 (1.65)	35.15 (2.96)	0.023	28.67 (2.03)	34.22 (2.95)	0.023
Total Mass (kg)	74.24 (11.79)	98.67 (9.24)	0.023	72.99 (12.56)	96.38 (8.08)	0.023
Fat Mass (kg)	31.17 (4.80)	42.53 (4.70)	0.023	41.89 (5.25)	30.53 (4.84)	0.023
Lean Mass (kg)	41.19 (7.07)	53.81 (4.72)	0.023	40.59 (7.83)	52.13 (3.45)	0.023
*Lean mass %	57.94 (2.50)	56.95 (1.36)	0.821	58.10 (3.46)	56.62 (1.78)	0.792
Body Fat %	42.06 (2.50)	48.05 (1.36)	0.280	41.90 (3.48)	43.37 (1.79)	0.480
Android Fat % of Body Fat	44.11 (2.37)	47.87 (3.22)	0.098	43.39 (3.63)	47.60 (2.55)	0.098
Gynoid Fat % of Body Fat	43.31 (3.77)	44.68 (3.45)	0.480	43.69 (3.72)	44.70 (4.05)	0.860
Waist to Hip Ratio	0.82 (0.03)	0.86 (0.07)	0.490	0.81 (0.04)	0.85 (0.06)	0.480
McAuleys ISI	9.76 (0.86)	9.93 (1.24)	0.749	9.71 (1.06)	9.78 (0.57)	0.886
Menopausal Status						
Pre- (count)	4	2	0.29**			
Post- (count)	2	5				

Table 1. For anthropometric and clinical time course differences over the intervention period, p-values were determined through a mixed model approach with fat burner group, week, and time as fixed effects, participant as random effect. P-values reported are for the relationship between HB and LB groups, where the addition of VB to the model, resulted in no significant differences identified (all p-value > 0.05). Distribution of pre- and post-menopausal status between burner group is also shown.

*lean mass includes bone mineral content

**Fisher's exact test demonstrated no significant association between HB and LB and menopausal status at baseline.

Table 2. List of each metabolite by cluster for the Partial Least Squares Discriminant Analysis (PLSDA).

Cluster	Cluster Name on PLSDA	Members	Species name	Cluster Component	Group HB or LB
6	NEFA	NEFA.C18.0	Non-esterified fatty acids	0.5043	HB
6	NEFA	NEFA.S.SFA	Non-esterified fatty acids	0.4721	HB
6	NEFA	NEFA.C16.0	Non-esterified fatty acids	0.4451	HB
6	NEFA	NEFA.C18.1n9	Non-esterified fatty acids	0.4112	HB
6	NEFA	NEFA.C16.1n7.C16.0	Non-esterified fatty acid ratio	-0.3944	HB
8	Not shown	<i>NEFA.C20.4n6</i>	<i>Non-esterified fatty acids</i>	<i>0.4614</i>	<i>HB</i>
8	Not shown	<i>NEFA.S.n.3.PUFA</i>	<i>Non-esterified fatty acid ratio</i>	<i>0.4607</i>	<i>HB</i>
8	Not shown	<i>NEFA.C22.6n3</i>	<i>Non-esterified fatty acids</i>	<i>0.4365</i>	<i>HB</i>
8	Not shown	<i>NEFA.C20.3n6</i>	<i>Non-esterified fatty acids</i>	<i>0.4036</i>	<i>HB</i>
8	Not shown	<i>NEFA.C20.5n3</i>	<i>Non-esterified fatty acids</i>	<i>0.3421</i>	<i>HB</i>
8	Not shown	<i>NEFA.C22.4n6</i>	<i>Non-esterified fatty acids</i>	<i>0.3232</i>	<i>HB</i>
10	TFA	TFA.C18.0.C16.0	Total fatty acid ratio	0.5452	LB
10	TFA	TFA.C16.1n7.C16.0	Total fatty acid ratio	0.5238	LB
10	TFA	NEFA.C20.4n6.C20.3n6	Total fatty acid ratio	-0.3739	LB
10	TFA	TFA.C18.1n9.C18.0	Total fatty acid ratio	-0.5372	LB
11	Lysophosphatidylcholine	lysoPC.a.C20.4	Lysophosphatidylcholine	0.3848	LB
11	Lysophosphatidylcholine	lysoPC.a.C16.0	Lysophosphatidylcholine	0.3793	LB
11	Lysophosphatidylcholine	lysoPC.a.C18.1	Lysophosphatidylcholine	0.3704	LB
11	Lysophosphatidylcholine	lysoPC.a.C18.0	Lysophosphatidylcholine	0.3602	LB
11	Lysophosphatidylcholine	lysoPC.a.C16.1	Lysophosphatidylcholine	0.3557	LB
11	Lysophosphatidylcholine	lysoPC.a.C18.2	Lysophosphatidylcholine	0.3254	LB
11	Lysophosphatidylcholine	lysoPC.a.C20.3	Lysophosphatidylcholine	0.3173	LB
11	Lysophosphatidylcholine	lysoPC.a.C17.0	Lysophosphatidylcholine	0.3002	LB
11	Lysophosphatidylcholine	Glu	Amino acid	-0.1334	LB
14	Amino Acids	Pro	Amino acid	0.5291	HB
14	Amino Acids	Gln	Amino acid	0.4993	HB
14	Amino Acids	Gly	Amino acid	0.4957	HB
14	Amino Acids	Ala	Amino acid	0.4744	HB
16	Amino Acids	Lys	Amino acid	0.4396	HB
16	Amino Acids	Arg	Amino acid	0.4354	HB
16	Amino Acids	Ser	Amino acid	0.4139	HB
16	Amino Acids	His	Amino acid	0.4082	HB
16	Amino Acids	H1	Amino acid	0.3906	HB
16	Amino Acids	Met.SO	Biogenic amines (Methionine sulfoxide)	0.3559	HB
17	NEFA	NEFA.C18.0.C16.0	Non-esterified fatty acid ratio	0.5889	HB
17	NEFA	NEFA.C22.0	Non-esterified fatty acids	0.5454	HB
17	NEFA	NEFA.C22.4n6.C20.4n6	Non-esterified fatty acid ratio	-0.3137	HB
17	NEFA	NEFA.C18.1n9.C18.0	Non-esterified fatty acid ratio	-0.5072	HB

19	<i>Not shown</i>	<i>PC.ae.C30.0</i>	<i>Phosphatidylcholine (Alkyl ethyl)</i>	0.5070	HB
19	<i>Not shown</i>	<i>PC.aa.C30.0</i>	<i>Phosphatidylcholine (Alkyl alkyl)</i>	0.4548	HB
19	<i>Not shown</i>	<i>PC.aa.C42.6</i>	<i>Phosphatidylcholine (Alkyl alkyl)</i>	0.4156	HB
19	<i>Not shown</i>	<i>PC.ae.C44.3</i>	<i>Phosphatidylcholine (Alkyl ethyl)</i>	0.4154	HB
19	<i>Not shown</i>	<i>total.DMA</i>	<i>Biogenic amines (dimethyl arginine)</i>	0.3287	HB
19	<i>Not shown</i>	<i>NEFA.C22.2n6</i>	<i>Non-esterified fatty acids</i>	0.2877	HB
22	Mixed species	PC.aa.C38.1	Phosphatidylcholine (Alkyl ethyl)	0.7071	LB
22	Mixed species	C2	Acylcarnitine (Acetyl)	0.7071	LB
23	Mixed species	Ac.Orn	AcetylOrnithine	0.7071	LB
23	Mixed species	C5-OH (C3-DC-M)	Acylcarnitine (Hydroxyvaleryl carnitine)	-0.7071	LB
			Methylmalonylcarnitine)		
29	<i>Not shown</i>	<i>NEFA.Total.Fatty.Acids</i>	<i>Non-esterified fatty acids</i>	0.5053	HB
29	<i>Not shown</i>	<i>NEFA.C18.2n6</i>	<i>Non-esterified fatty acids</i>	0.5028	HB
29	<i>Not shown</i>	<i>NEFA.C14.0</i>	<i>Non-esterified fatty acids</i>	0.4706	HB
29	<i>Not shown</i>	<i>NEFA.C18.1n7</i>	<i>Non-esterified fatty acids</i>	0.3912	HB
29	<i>Not shown</i>	<i>NEFA.S.n.6.S.n.3</i>	<i>Non-esterified fatty acid ratio</i>	0.3426	HB
30	Phosphatidylcholine	PC.aa.C38.5	Phosphatidylcholine	0.3284	LB
30	Phosphatidylcholine	SM.C24.1	Sphingomyelin	0.3225	LB
30	Phosphatidylcholine	PC.ae.C36.3	Phosphatidylcholine	0.3213	LB
30	Phosphatidylcholine	PC.ae.C36.4	Phosphatidylcholine	0.3188	LB
30	Phosphatidylcholine	PC.ae.C44.5	Phosphatidylcholine	0.3145	LB
30	Phosphatidylcholine	PC.ae.C42.5	Phosphatidylcholine	0.3125	LB
30	Phosphatidylcholine	PC.ae.C42.4	Phosphatidylcholine	0.3063	LB
30	Phosphatidylcholine	PC.ae.C42.3	Phosphatidylcholine	0.3041	LB
30	Phosphatidylcholine	PC.ae.C38.3	Phosphatidylcholine	0.2901	LB
30	Phosphatidylcholine	SM.C20.2	Sphingomyelin	0.2654	LB
30	Phosphatidylcholine	C18:2	Acylcarnitine (Octadecadienyl carnitine)	0.2143	LB

Table 2. Table only includes clusters that reached the threshold for VIP (>1.5 in regular PLSDA and >1.0 in all 10 components in sPLSDA with repeated measures) and were included in the subsequent PLS analysis. *Clusters that have been italicized were identified by PLSDA but not by sPLSDA and were therefore not included in our interpretation.* NEFA: non-esterified fatty acids. TFA: total fatty acids. lysoPC: Lysophosphatidylcholine. PC: Phosphatidylcholine. aa: acyl-acyl. ae: acyl-acetyl. C2, C5-OH, and C18:2 are all acylcarnitines.




Figure 4. PLSDA demonstrates clustering of various lipids that are strongly associated with fat burning groups. Cluster components (CC) from variable clustering of disappearance rates of metabolites were used as input predicting fat burner groups HB and LB. Inclusion of VB did not allow for model convergence, so the presented model is only comparing HB and LB. (A) Highlights the clusters that were associated with each burner group. HB (n = 6), LB (n = 7), and VB (n = 31). Partial Least Squares Discriminant Analysis using NIPALS algorithm converged on a model with 2 minimizing factors, where 17.7% of X and 71.6% of Y variables were explained with a Q^2 of 0.20 (B) Variable importance plot (VIP) scores were chosen for analysis if they were over 1, with blue indicating association with LB and red for HB. TFA: Total Fatty acids, NEFA: Non-esterified fatty acids




Figure 5: A summary of late postprandial (disappearance) metabolite changes that delineated HB and LB phenotypes. A and B summarize metabolic milieu in LB and C and D do the same in HB group women. The green '+' and red '-' signs denote the speculated increase and decrease of metabolic processes based on observed disappearance trends. Metabolites in yellow oval/boxes were measured parameters, and light blue boxes are upstream or downstream metabolites not measured and hypothesized based on known biosynthesis and degradation pathways.