

Human cardiac stromal cells exposed to SARS-CoV-2 evolve into hyper-inflammatory/*pro*-fibrotic phenotype and produce infective viral particles depending on the levels of ACE2 receptor expression

Supplementary Information

Ethical Information

Derivation of cells from human atrial appendage was approved by the Ethical Committee at Centro Cardiologico Monzino, IRCCS; Milan. Each patient, admitted for routine cardiac surgery interventions signed an informed consent to authorize collection of the material.

Table S1. Patients demographic and risk characteristics, with an indication of the anti-hypertensive therapy. Data in the columns on the right indicate the normalized ACE2/GAPDH protein and the ACE2 mRNA (Δ Ct values) expression levels, as detected by western analysis and RT-qPCR, respectively. In blue, red and green colors, the subjects that were chosen as ACE2 ‘Lo’, ‘Mid’ and ‘Hi’ cST-Cs donors, respectively.

Patient	Sex	Age range	Diabetes	Dyslipidemia	Hypertension	Therapy	ACE2 normalized level (WB)	ACE2 mRNA (Δ Ct)
1	M	70-80	Y	Y	Y	Perindopril	0.334	24,984
2	M	70-80	N	Y	Y	Enalapril	0.989	16.022
3	M	70-80	Y	N	Y	-	0.854	15.215
4	M	70-80	N	Y	Y	Losartan	0.989	15.794
5	M	70-80	Y	N	Y	Olmesartan	1.270	14.113
6	M	70-80	N	N	Y	-	0.854	17.388
7	M	60-70	N	Y	Y	Perindopril	0.632	19.443
8	M	50-60	Y	Y	Y	Ramipril	1.147	16.815
9	M	70-80	Y	Y	Y	Losartan	1.173	14.607
10	F	50-60	N	N	Y	Perindopril	0.918	19.749

Preparation of cSt-Cs and antigenic/proteomic characterization

For preparation of human cardiac stromal cells, human right atrial appendage biopsies, obtained from patients undergoing cardiac surgery, were subjected to mechanical fragmentation and enzymatic digestion, and cultured for 4 weeks in Iscove’s modified Dulbecco’s medium (IMDM) supplemented with 20% Hyclone characterized Fetal Bovine Serum (FBS), 1% penicillin-streptomycin, 1% L-glutamine and 0.1mM2-mercaptoethanol. Explant-derived cells were plated on poly-D-lysine coated wells in the following media to obtain primary cardiospheres: 35% IMDM/65% DMEM/F-12 Mix, 3.5% Hyclone characterized FBS, 1% penicillin-streptomycin, 1% L-glutamine, 0.1 mM 2-mercaptoethanol, 1 unit/ml thrombin, 1:50 B-27, 80 ng/ml bFGF, 25 ng/ml EGF, and 4 ng/ml cardiotrophin-1 (all Peprotech). cSt-Cs were then derived by dissociating cardiospheres followed by plating onto Fibronectin-coated dishes. For antigenic characterization, cST-Cs suspensions (passages 1-4) were incubated with a combination of fluorochrome-conjugated antibodies against CD29, CD31, CD44, CD90, CD105 and CD144 (or the corresponding Isotype antibodies; all from BD Biosciences) for 15 minutes at room temperature (RT). Cells were analyzed with Gallios Flow Cytometer (Beckman Coulter) equipped with Kaluza analysis software (Beckman Coulter).

Western blot

Cells were lysed in a buffer containing 10 mM Tris-Cl, pH 7.4, 150 mM NaCl, 5 mM EDTA, 1% (v/v) Triton X-100, 1% (w/v) sodium deoxycholate, 0.1% (w/v), sodium dodecyl sulfate and 1% (v/v), protease and phosphatase inhibitor mixture (Sigma-Aldrich). Whole cell lysates were sonicated, centrifuged for 15 min at 14 000 g; cell supernatants were then collected. Proteins were quantified by BCA protein assay kit (Pierce Chemical Co). Cell lysates (30 μ g per lane) were diluted in Laemli sample buffer, heated at 95 °C for 5 minutes,

run onto 4-12% gradient SDS polyacrylamide gels (Invitrogen), and transferred to nitrocellulose membranes. The blots were blocked with Tris Buffered-saline containing 5% (w/v) BSA at RT for 1 hour. Overnight incubation at 4 °C with ACE2 and GAPDH primary antibodies (Novus Biologicals; Santa Cruz Biotechnology) was performed. Membranes were incubated with appropriate secondary antibodies for 20 minutes. Images were acquired with LI-COR Odyssey and band intensities were quantified using ImageJ software.

Preparation of SARS-CoV-2 isolates

Viral stock used to infect cSt-Cs was the first Italian SARS-Cov-2 isolate (named 2019-nCoV/Italy-INMI1 and available from EVAg, Ref-SKU: 008V-03893) obtained from a patient hospitalized at National Institute for Infectious Diseases “Lazzaro Spallanzani” (INMI) in Rome. Virus was expanded on Vero E6 cells (ATCC) maintained in Modified Eagle Medium (MEM) supplemented with 10% heat inactivated FBS, 2 mM L-glutamine, and 1% penicillin/streptomycin solution at 37°C, in a humidified atmosphere of 5% CO₂. Virus titer was determined in culture supernatants as fifty-percent tissue culture infective dose (TCID₅₀/mL), according to Reed and Muench method¹⁶.

SARS-CoV-2 infection of cSt-Cs

For infection with SARS-CoV-2, the cSt-Cs lines were seeded in 6-well culture plates. After reaching cell confluence, they were exposed to Vero E6 supernatant containing SARS-CoV-2 at the multiplicity of infection (MOI) of 0.1, 1 and 10 for 2 hour at 37°C. Supernatant was then removed and, after three washes with DPBS, fresh culture medium was added for 3 days. cSt-Cs supernatants were collected at 2, 24 and 72h post infection to assess the kinetics of viral replication. Supernatants were in part lysed in TRIZOL for molecular detection of the virus by RT-qPCR and digital PCR immediately and in part frozen at -80 °C for further investigation. At the time of supernatant collection, cells were lysed in TRIZOL (Invitrogen) to recover cellular RNAs and perform the analyses on cellular and viral RNAs (see below).

Titration of SARS-CoV-2 produced by the cST-Cs in Vero E6 cells

The infectivity of the virus produced by cSt-Cs collected at 2h, 24h and 72h post infection was measured by Reed and Muench method on Vero E6 cells¹⁶. Vero E6 cells were seeded in 96-wells culture plates. Ten fold dilutions of cellular supernatants were prepared in culture medium supplemented with 2% heat-inactivated FBS and 0,05ml of each dilution was added to the cells. The plates were incubated for 48-72h at 37 °C. The kinetics of virus infection was followed by PFU for 72h and expressed as fifty-percent tissue culture infective dose (TCID₅₀/mL).

RT-qPCR in cellular lysates

Total RNA from infected cells at various times with different MOI was extracted using TRIZOL (Invitrogen), quantified with NanoDrop-1000 spectrophotometer (Thermo Fisher Scientific). Superscript III (Thermo Fisher Scientific) was used for reverse transcription. Quantitative real-time PCR analysis were performed with Power SYBR Green PCR Master Mix (Applied Biosystems) in an ABI 7900 Fast thermal cycler to detect *ACE2*, inflammatory cytokines and *pro*-fibrotic gene amplification products (primers details in **Table S2**). The reported expression levels were calculated relative to GAPDH mRNA, used as an internal standard control. The fold change of the genes in infected cells vs. control samples was calculated as $2^{-\Delta\Delta CT}$ and the statistical analysis was done on the ΔCT values.

Table S2. List of primers used to perform RT-qPCR analyses on cellular extracts.

Gene	Sequence
<i>hACE2</i>	Forward: 5'- GCCACTGCTCAACTACTTTG -3' Reverse: 5'- GCTTATCCTCACTTTGATGCTTTG -3'
<i>hIL-1β</i>	Forward: 5'- CAGCCAATCTTCATTGCTCAAG -3' Reverse: 5'- GAACAAGTCATCCTCATTGCC -3'

<i>hIL-6</i>	Forward: 5'- GCAGATGAGTACAAAAGTCCTGA -3' Reverse: 5'- TTCTGTGCCTGCAGCTTC -3'
<i>hCCL2</i>	Forward: 5'- AGCAGCCACCTTCATTCC -3' Reverse: 5'- GCCTCTGCACTGAGATCTTC -3'
<i>hACTA2</i>	Forward: 5'- AGAGTTACGAGTTGCCTGATG -3' Reverse: 5'- CTGTTGTAGGTGGTTTCATGGA -3'
<i>hCTGF</i>	Forward: 5'- AGGAGTGGGTGTGTGACGA -3' Reverse: 5'- CCAGGCAGTTGGCTCTAATC -3'
<i>hCOL1A1</i>	Forward: 5'- GGACACAGAGGTTTCAGTGG -3' Reverse: 5'- CCAGTAGCACCATCATTTCC -3'
<i>hCOL3A</i>	Forward: 5'- AGCTACGGCAATCCTGAACT -3' Reverse: 5'- GGGCCTTCTTTACATTTCCA-3'
<i>hGAPDH</i>	Forward: 5'- AATCCCATCACCATCTTCCAG -3' Reverse: 5'- AAATGAGCCCCAGCCTTC -3'

mRNA analysis with gene arrays

Human Cardiotoxicity RT2 Profiler PCR Array (PAHS-095ZA, Qiagen) and Human NF- κ B Signalling pathway plus RT2 Profiler PCR Array (PAHS-025YE, Qiagen) were performed on an Applied Biosystems (ABI) 7900 according to the manufacturer's instructions using 400 ng of each RNA preparation. Data were expressed as fold changes compared to the control condition, calculated with the $2^{-\Delta\Delta C_t}$ method (i.e., using the cells harvested before infection as internal calibrator). Differential and clustering analyses were performed using algorithms incorporated in the open-source software MultiExperiment Viewer (MeV) version 4.9. Differences in gene expression were identified by one-way ANOVA: P-values computed from the F-distribution <0.05 were deemed as significant. Hierarchical cluster analysis was performed to visualize the similarities and differences in gene expression among the time points. Normalized gene expression values were mean-centered, scaled and clustered by Pearson's correlation and average linkage.

Quantification of SARS-CoV-2 in cST-Cs supernatants and cell extracts by conventional and digital PCR methods

For an initial evaluation of the virus released by infected cST-Cs in culture supernatant during the time course (2, 24 and 72h p.i.) we used a diagnostic RT-qPCR assay (SimplexaTM COVID-19 Direct assay, Diasorin S.p.A) targeting *Spike* (S) and *ORF1ab* SARS-CoV-2 coding sequences. Data were expressed as Ct values of each gene as detected in the supernatant of the cells at each time point.

For viral RNA quantification in cellular extracts and supernatants, a one-step chip-based digital PCR was performed on a QuantStudio 3D Digital PCR System platform composed by the QuantStudio 3D Instrument, the Dual Flat Block GeneAmp PCR System 9700 and the QuantStudio 3D Digital PCR Chip Loader (Thermo Fisher Scientific). TaqMan Fast Virus 1-Step Master Mix (Applied Biosystems) was used in combination with TaqMan primers and FAM probe for viral RNA (**Supplementary Table S3**), purchased from Integrated DNA Technologies.

Regarding the supernatant viral RNA quantification, 5 μ L of the extracted extracellular RNA were analyzed with primers and probe for viral N gene. Concerning the intracellular viral RNA quantification, 20 ng of extracted cellular RNA were analyzed with primers and probes for viral N, E, and RdRp genes. Human RNase P gene was used as internal amplification control. Reverse transcription was performed at 50 °C for 10 minutes, followed by inactivation/initial denaturation at 96 °C for 5 minutes. PCR cycling consisted of a denaturation step at 98 °C for 30 seconds followed by annealing/extension at 56 °C for 1 minute (40 cycles) and a final extension at 60 °C for 5 minutes. Analyses were executed with the online version of the QuantStudio 3D

AnalysisSuite (Thermo Fisher Scientific Cloud) following the Manufacturer's instructions.

Table S3. List of primers and probes used for the viral RNA evaluation with dPCR.

Gene	Type	Concentration	Number
nCOV_N	Forward Primer	50 nmol	10006824
nCOV_N	Reverse Primer	50 nmol	10006825
nCOV_N	Probe	25 nmol	10006826
E_Sarbeco	Forward Primer	50 nmol	10006888
E_Sarbeco	Reverse Primer	50 nmol	10006890
E_Sarbeco	Probe	25 nmol	10006892
RdRP_SARSr	Forward Primer	50 nmol	10006860
RdRP_SARSr	Reverse Primer	50 nmol	10006881
RdRP_SARSr	Probe	25 nmol	10006886
RNase P	Forward Primer	50 nmol	10006827
RNase P	Reverse Primer	50 nmol	10006828
RNase P	Probe	25 nmol	10006829

Immunofluorescence

After fixation with 4% paraformaldehyde, cells were permeabilized for 1 hour at RT with PBS containing 3% (w/v) BSA and 1% (v/v) Triton X-100, followed by incubation with COVID-19 patient derived-serum and the primary antibodies against α SMA (DAKO) and Coll (Rockland) at 4°C overnight. Cells were then incubated for 1 hour at RT with the appropriate secondary antibodies, Phalloidin-TRITC and DAPI nuclear dye. Digital images were obtained using an LSM710 confocal scanning microscope (Carl Zeiss) equipped with the AiryScan super resolution system.




Figure S1. VERO E6 cells were exposed to culture supernatants of cSt-Cs infected with SARS-CoV-2 to perform viral titration. The picture shows the cytopathic effect induced by SARS-CoV-2 contained in the ACE2 'Hi' cells supernatant after infection (1 MOI) for 72 hours.

Table S4. List of the transcripts significantly affected in cSt-Cs in response to SARS-CoV-2 exposure. Data are expressed as fold change of each transcript in the given cell/infection condition/time vs. the levels of the same gene expressed in uninfected cells. The statistical significance (P-value) of the data has been calculated with **ANOVA**.

Gene	Lo 2h	Mid 2h	Hi 2h	Lo 24h	Mid 24h	Hi 24h	Lo 72h	Mid 72h	Hi 72h	Mean 2h	Mean 24h	Mean 72h	P-value
PDK4	1,56	1,16	1,06	0,61	0,19	0,13	0,40	0,27	0,13	1.26 ± 0.26	0.31 ± 0.26	0.27 ± 0.14	0,0029
EGR1	0,52	1,45	0,92	0,21	0,60	0,12	1,77	2,46	1,80	0.96 ± 0.47	0.31 ± 0.26	2.01 ± 0.39	0,0045
STAT1	0,79	1,08	0,94	0,78	0,73	0,66	1,44	2,72	2,09	0.94 ± 0.15	0.72 ± 0.06	2.08 ± 0.64	0,0097
NEXN	1,20	0,70	1,01	1,80	1,46	1,22	2,52	2,32	1,75	0.97 ± 0.25	1.49 ± 0.29	2.20 ± 0.40	0,0097
BCL2A1	0,62	0,84	0,54	0,11	0,16	0,40	0,22	0,29	0,37	0.67 ± 0.16	0.22 ± 0.16	0.29 ± 0.08	0,0138
ZNF23	1,06	0,95	0,99	1,88	1,59	1,85	1,45	2,24	2,37	1.00 ± 0.06	1.77 ± 0.16	2.02 ± 0.50	0,0148
ITPR2	0,76	1,02	1,05	1,39	1,76	2,04	1,54	2,01	1,50	0.94 ± 0.16	1.73 ± 0.33	1.68 ± 0.28	0,0189
ICAM1	0,87	1,62	0,82	1,01	0,52	0,78	2,00	1,60	1,74	1.10 ± 0.45	0.77 ± 0.25	1.78 ± 0.20	0,0209
SERPINE1	0,78	0,88	0,74	1,01	1,46	2,28	1,89	3,08	3,78	0.80 ± 0.07	1.58 ± 0.64	2.92 ± 0.96	0,0218
HSPH1	1,12	0,95	1,42	1,03	1,14	0,97	0,23	0,77	0,59	1.16 ± 0.24	1.05 ± 0.09	0.53 ± 0.27	0,0246
CCL7	0,99	1,54	1,05	1,47	0,46	1,16	4,15	1,91	4,78	1.19 ± 0.30	1.03 ± 0.52	3.61 ± 1.51	0,0258
PVR	1,25	0,94	1,03	1,03	1,35	2,01	1,85	2,03	2,35	1.07 ± 0.16	1.46 ± 0.50	2.08 ± 0.25	0,0288
RND1	0,55	1,72	1,12	3,27	2,45	3,15	2,65	2,35	1,18	1.13 ± 0.59	2.96 ± 0.44	2.06 ± 0.78	0,0307
FHL1	1,06	0,74	1,21	0,65	0,40	1,13	1,72	1,26	1,91	1.00 ± 0.24	0.73 ± 0.37	1.63 ± 0.33	0,0338
ZNF148	1,03	0,96	1,12	1,66	1,99	1,51	0,93	1,60	1,31	1.04 ± 0.08	1.72 ± 0.25	1.28 ± 0.34	0,0369
FOSL1	0,91	1,20	1,01	0,30	0,49	0,95	0,20	0,45	0,55	1.04 ± 0.15	0.58 ± 0.33	0.40 ± 0.18	0,0382
HMOX1	0,95	2,64	1,41	0,29	0,31	0,35	0,14	0,93	0,48	1.67 ± 0.87	0.32 ± 0.03	0.52 ± 0.40	0,0492