

Body mass index and risk of COVID-19 diagnosis, hospitalisation, and death: a population-based multi-state cohort analysis including 2,524,926 people in Catalonia, Spain

Martina Recalde, *PhD student*^{1,2}, Andrea Pistillo, *statistician*¹, Sergio Fernandez-Bertolin, *data scientist*¹, Elena Roel, *PhD student*^{1,2}, Maria Aragon, *SIDIAP coordinator*¹, Heinz Freisling, *scientist*³, Daniel Prieto-Alhambra, *Professor of pharmaco- and device epidemiology*⁴, Edward Burn, *post-doctoral researcher*^{1,4*}, Talita Duarte-Salles, *senior epidemiologist*^{1*}

*Joint senior authorship

1. Fundació Institut Universitari per a la recerca a l'Atenció Primària de Salut Jordi Gol i Gurina (IDIAPJGol), Barcelona, Spain
2. Universitat Autònoma de Barcelona, Spain
3. International Agency for Research on Cancer (IARC-WHO), 150 Cours Albert Thomas, 69008 Lyon, France
4. Centre for Statistics in Medicine, NDORMS, University of Oxford

CORRESPONDING AUTHORS

Daniel Prieto-Alhambra
Botnar Research Centre, Windmill Road
OX37LD, Oxford, United Kingdom
daniel.prietoalhambra@ndorms.ox.ac.uk

Talita Duarte-Salles
Fundació Institut Universitari per a la recerca a l'Atenció Primària de Salut Jordi Gol i Gurina (IDIAPJGol)
Gran Via Corts Catalanes, 587 àtic
08007 Barcelona - Spain
tduarte@idiapjgol.org

Appendix

Figure S1. Flowchart with the inclusion and exclusion criteria of the study population

Figure S2. Directed Acyclic Graph for the possible causal effect of obesity on COVID-19 outcomes used to adjust the Cox proportional hazard models

Table S1. Characteristics of individuals excluded from the study due to missing information on BMI, smoking status and/or the MEDEA deprivation index

Table S2. Characteristics of the study participants by multi-state transition

Figure S3. Association between body mass index and the risk of COVID-19 outcomes with different types of model-adjustment, allowing for non-linear effects, with 95% CIs

Table S3. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, for the unadjusted, adjusted for age and sex and fully adjusted models

Table S4. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, stratified by age and sex

Figure S4. Association between body mass index and the risk of COVID-19 diagnosis, allowing for non-linear effects, with 95% CIs, stratified by calendar month

Table S5. Hazard Ratios of the risk of COVID-19 diagnosis related to body mass index, allowing for non-linear effects, with 95% CIs, stratified by calendar month

Figure S5. Hazard Ratios of COVID-19 outcomes related to body mass index in categories, with 95% CIs


Figure S6. Association between body mass index and the risk of COVID-19 outcomes, allowing for non-linear effects, with 95% CIs, after multiple imputation of missing data on BMI, smoking status, and/or the MEDEA deprivation index

Table S6. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, after multiple imputation of missing data on BMI, smoking status, and/or the MEDEA deprivation index

Figure S7. Association between body mass index and the risk of COVID-19 outcomes, allowing for non-linear effects, with 95% CIs among people with a BMI measurement not older than five years prior to the index date


Table S7. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs among people with a BMI measurement not older than five years prior to the index date

Figure S1. Flowchart with the inclusion and exclusion criteria of the study population



Abbreviations: BMI: Body Mass Index; COVID-19: Coronavirus Disease 2019; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”; SIDIAP: Information System for Research in Primary Care.

Figure S2. Directed Acyclic Graph for the possible causal effect of obesity on COVID-19 outcomes used to adjust the Cox proportional hazard models



Notes: Green lines indicate a possible pathway between the exposure and outcome of interest.

Abbreviations: COVID-19: Coronavirus Disease 2019; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”; SES: Socioeconomic status.

Table S1. Characteristics of individuals excluded from the study due to missing information on BMI, smoking status and/or the MEDEA deprivation index

	Overall	BMI recorded and complete data on covariates	Missing data on BMI ^a or covariates ^b	SMD	Missing BMI ^a	BMI recorded but missing data on covariates ^b
N	4,619,841	2,524,926	2,094,915		1,357,553	737,362
BMI, median (IQR)	26.5 (23.5-30.0)	26.5 (23.5-29.9)	26.6 (23.6-30.1)	0.260	-	26.6 (23.6-30.1)
Age, median (IQR)	48 (36.0-63.0)	52 (39.0-67.0)	44 (32.0-57.0)	0.394	40 (27.0-50.0)	56 (43.0-71.0)
Age, n (%)				0.371		
18 to 39	1,437,297 (31.1)	633,408 (25.1)	803,889 (38.4)		678,297 (50.0)	125,592 (17.0)
40 to 59	1,785,852 (38.7)	958,492 (38.0)	827,360 (39.5)		535,200 (39.4)	292,160 (39.6)
60 to 69	615,538 (13.3)	405,640 (16.1)	209,898 (10.0)		94,137 (6.9)	115,761 (15.7)
70 to 79	468,662 (10.1)	325,948 (12.9)	142,714 (6.8)		32,370 (2.4)	110,344 (15.0)
80 or older	312,492 (6.8)	201,438 (8.0)	111,054 (5.3)		17,549 (1.3)	93,505 (12.7)
Female sex, n (%)	2,361,933 (51.1)	1,386,678 (54.9)	975,255 (46.6)	0.168	545,827 (40.2)	429,428 (58.2)
Smoking status, n (%)				1.841		
Never smoker	712918 (15.4)	517558 (20.5)	195360 (9.3)		159987 (11.8)	-
Former smoker	773325 (16.7)	663383 (26.3)	109942 (5.2)		66878 (4.9)	-
Current smoker	1835150 (39.7)	1343985 (53.2)	491165 (23.4)		406087 (29.9)	-
Missing	1298448 (28.1)	-	1298448 (62.0)		724601 (53.4)	-
MEDEA deprivation index, n (%)				0.574		
Quintile 1 (least deprived)	714440 (15.5)	394503 (15.6)	319937 (15.3)		248387 (18.3)	-
Quintile 2	704177 (15.2)	399883 (15.8)	304294 (14.5)		215825 (15.9)	-

Quintile 3	697299 (15.1)	405747 (16.1)	291552 (13.9)		201562 (14.8)	-
Quintile 4	693046 (15.0)	410440 (16.3)	282606 (13.5)		189614 (14.0)	-
Quintile 5 (most deprived)	687266 (14.9)	410231 (16.2)	277035 (13.2)		189495 (14.0)	-
Rural	832483 (18.0)	504122 (20.0)	328361 (15.7)		219066 (16.1)	-
Missing	291130 (6.3)	-	291130 (13.9)		93604 (6.9)	-
Comorbidities, n (%)						
Autoimmune condition	259,383 (5.6)	170,240 (6.7)	89,143 (4.3)	0.109	36,253 (2.7)	52,890 (7.2)
Chronic kidney disease	201,456 (4.4)	141,921 (5.6)	59,535 (2.8)	0.138	6,006 (0.4)	53,529 (7.3)
COPD	119,669 (2.6)	86,723 (3.4)	32,946 (1.6)	0.119	4,112 (0.3)	28,834 (3.9)
Heart disease	516,553 (11.2)	363,012 (14.4)	153,541 (7.3)	0.228	31,384 (2.3)	122,157 (16.6)
Hyperlipidemia	505,365 (10.9)	357,572 (14.2)	147,793 (7.1)	0.232	43,730 (3.2)	104,063 (14.1)
Hypertension	687,824 (14.9)	514,533 (20.4)	173,291 (8.3)	0.351	35,457 (2.6)	137,834 (18.7)
Malignant neoplasm	289,827 (6.3)	197,171 (7.8)	92,656 (4.4)	0.142	27,211 (2.0)	65,445 (8.9)
Type 2 diabetes	317,254 (6.9)	236,253 (9.4)	81,001 (3.9)	0.222	6,025 (0.4)	74,976 (10.2)

Notes: a) Includes individuals who do not have a BMI recorded in their electronic health records and those with a BMI measurement recorded before being 18 years of age. b) Includes individuals without information on smoking or socioeconomic status. Malignant neoplasm does not include non-melanoma skin cancer.

Abbreviations: BMI: Body Mass Index; COPD: Chronic Obstructive Pulmonary Disease; IQR: Interquartile range; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”; SMD: Standardized mean difference.

Table S2. Characteristics of the study participants by multi-state transition


	General population	From general population			From diagnosed with COVID-19		From hospitalised with COVID-19
		To diagnosed with COVID-19	To hospitalised with COVID-19	To death	To hospitalised with COVID-19	To death	To death
N	2,524,926	57,443	5,191	5,276	5,671	1,166	1,301
BMI, median (IQR)	26 (23.5-29.9)	27 (23.5-30.3)	29 (25.9-32.1)	27 (23.6-30.1)	28 (25.7-32.0)	27 (24.1-30.4)	29 (25.6-31.9)
BMI in WHO categories, n (%)							
Normal or underweight	951,280 (37.7)	20,931 (36.4)	939 (18.1)	1,907 (36.1)	1,093 (19.3)	378 (32.4)	269 (20.7)
Overweight	952,479 (37.7)	21,369 (37.2)	2,233 (43.0)	2,024 (38.4)	2,421 (42.7)	470 (40.3)	541 (41.6)
Obesity	621,167 (24.6)	15,143 (26.4)	2,019 (38.9)	1,345 (25.5)	2,157 (38.0)	318 (27.3)	491 (37.7)
Age, median (IQR)	52 (39.0-67.0)	48 (38.0-60.0)	70 (58.0-79.0)	85 (75.0-90.0)	60 (50.0-72.0)	86 (79.0-91.0)	79 (72.0-86.0)
Age, n (%)							
18 to 39	633,408 (25.1)	16,502 (28.7)	264 (5.1)	19 (0.4)	505 (8.9)	1 (0.1)	5 (0.4)
40 to 59	958,492 (38.0)	26,455 (46.1)	1,140 (22.0)	295 (5.6)	2,203 (38.8)	29 (2.5)	69 (5.3)
60 to 69	405,640 (16.1)	6,927 (12.1)	1,100 (21.2)	526 (10.0)	1,176 (20.7)	69 (5.9)	163 (12.5)
70 to 79	325,948 (12.9)	4,198 (7.3)	1,490 (28.7)	974 (18.5)	1,083 (19.1)	195 (16.7)	446 (34.3)
80 or older	201,438 (8.0)	3,361 (5.9)	1,197 (23.1)	3,462 (65.6)	704 (12.4)	872 (74.8)	618 (47.5)
Female sex, n (%)	1,386,678 (54.9)	35,236 (61.3)	2,168 (41.8)	2,379 (45.1)	2,568 (45.3)	632 (54.2)	471 (36.2)
Smoking status, n (%)							
Never smoker	517,558 (20.5)	10,103 (17.6)	328 (6.3)	509 (9.6)	346 (6.1)	64 (5.5)	58 (4.5)

Former smoker	663,383 (26.3)	15,886 (27.7)	2,174 (41.9)	2,268 (43.0)	2,103 (37.1)	502 (43.1)	635 (48.8)
Current smoker	1,343,985 (53.2)	31,454 (54.8)	2,689 (51.8)	2,499 (47.4)	3,222 (56.8)	600 (51.5)	608 (46.7)
MEDEA deprivation index, n (%)							
Quintile 1 (least deprived)	394,503 (15.6)	8,755 (15.2)	759 (14.6)	873 (16.5)	783 (13.8)	239 (20.5)	200 (15.4)
Quintile 2	399,883 (15.8)	9,423 (16.4)	835 (16.1)	750 (14.2)	943 (16.6)	162 (13.9)	191 (14.7)
Quintile 3	405,747 (16.1)	9,624 (16.8)	909 (17.5)	736 (13.9)	1,019 (18.0)	189 (16.2)	228 (17.5)
Quintile 4	410,440 (16.3)	10,077 (17.5)	977 (18.8)	690 (13.1)	1,037 (18.3)	111 (9.5)	219 (16.8)
Quintile 5 (most deprived)	410,231 (16.2)	9,942 (17.3)	1,018 (19.6)	644 (12.2)	1,027 (18.1)	106 (9.1)	190 (14.6)
Rural	504,122 (20.0)	9,622 (16.8)	693 (13.4)	1,583 (30.0)	862 (15.2)	359 (30.8)	273 (21.0)
Comorbidities, n (%)							
Autoimmune condition	170,240 (6.7)	4,025 (7.0)	541 (10.4)	615 (11.7)	496 (8.7)	125 (10.7)	152 (11.7)
Chronic kidney disease	141,921 (5.6)	2,500 (4.4)	890 (17.1)	1,783 (33.8)	585 (10.3)	402 (34.5)	404 (31.1)
COPD	86,723 (3.4)	1,622 (2.8)	513 (9.9)	717 (13.6)	326 (5.7)	126 (10.8)	179 (13.8)
Heart disease	363,012 (14.4)	7,083 (12.3)	1,674 (32.2)	2,733 (51.8)	1,228 (21.7)	561 (48.1)	625 (48.0)
Hyperlipidemia	357,572 (14.2)	7,439 (13.0)	1,040 (20.0)	786 (14.9)	1,036 (18.3)	184 (15.8)	260 (20.0)
Hypertension	514,533 (20.4)	9,923 (17.3)	1,726 (33.2)	1,940 (36.8)	1,609 (28.4)	422 (36.2)	481 (37.0)
Malignant neoplasm	197,171 (7.8)	3,588 (6.2)	861 (16.6)	1,828 (34.6)	641 (11.3)	272 (23.3)	326 (25.1)
Type 2 diabetes	236,253 (9.4)	4,327 (7.5)	1,153 (22.2)	1,313 (24.9)	927 (16.3)	288 (24.7)	386 (29.7)

Notes: BMI categories: underweight or normal weight (BMI <18.5 kg/m² and between ≥18.5 and <25 kg/m²), overweight (BMI ≥25 and <30 kg/m²) and obesity (BMI ≥30 kg/m²). Malignant neoplasm does not include non-melanoma skin cancer.

Abbreviations: BMI: Body Mass Index; COPD: Chronic Obstructive Pulmonary Disease; COVID-19: Coronavirus Disease 2019; IQR: Interquartile range; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”; WHO: World Health Organization.

Figure S3. Association between body mass index and the risk of COVID-19 outcomes with different types of model-adjustment, allowing for non-linear effects, with 95% CIs



Notes: the fully adjusted models are adjusted for age, sex, smoking status and the MEDEA deprivation index.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.

Table S3. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, for the unadjusted, adjusted for age and sex and fully adjusted models

	From general population						From diagnosed with COVID-19						From hospitalised with COVID-19		
	to diagnosed with COVID-19			to hospitalised with COVID-19			to hospitalised with COVID-19			to death			to death		
BMI values (kg/m ²)	Un-adjusted	Adjusted for age and sex	Fully adjusted	Un-adjusted	Adjusted for age and sex	Fully adjusted	Un-adjusted	Adjusted for age and sex	Fully adjusted	Un-adjusted	Adjusted for age and sex	Fully adjusted	Un-adjusted	Adjusted for age and sex	Fully adjusted
16	1.01 (0.98-1.03)	0.80 (0.78-0.82)	0.81 (0.79-0.84)	0.26 (0.23-0.29)	0.54 (0.49-0.60)	0.58 (0.53-0.64)	0.29 (0.26-0.32)	0.48 (0.43-0.53)	0.51 (0.46-0.57)	0.90 (0.85-0.96)	1.27 (1.07-1.51)	1.28 (1.07-1.52)	1.07 (1.01-1.14)	1.17 (1.01-1.36)	1.20 (1.02-1.42)
19	1.00 (0.99-1.02)	0.89 (0.88-0.90)	0.90 (0.89-0.91)	0.51 (0.48-0.54)	0.75 (0.72-0.78)	0.77 (0.74-0.81)	0.53 (0.51-0.56)	0.69 (0.65-0.73)	0.71 (0.68-0.75)	0.95 (0.92-0.98)	1.13 (1.03-1.23)	1.13 (1.04-1.23)	1.04 (1.00-1.07)	1.08 (1.00-1.17)	1.08 (1.00-1.16)
22	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference
25	1.00 (0.99-1.01)	1.11 (1.10-1.12)	1.10 (1.09-1.11)	1.85 (1.76-1.95)	1.30 (1.26-1.34)	1.27 (1.22-1.31)	1.77 (1.69-1.85)	1.41 (1.35-1.48)	1.37 (1.31-1.43)	1.05 (1.02-1.09)	0.91 (0.84-0.98)	0.90 (0.84-0.97)	0.97 (0.93-1.00)	0.92 (0.86-1.00)	0.97 (0.91-1.03)
28	1.02 (1.00-1.04)	1.20 (1.18-1.22)	1.17 (1.15-1.19)	2.79 (2.58-3.01)	1.64 (1.55-1.74)	1.56 (1.47-1.66)	2.60 (2.42-2.79)	1.83 (1.70-1.97)	1.74 (1.61-1.87)	1.11 (1.05-1.18)	0.88 (0.78-0.99)	0.88 (0.78-0.99)	0.93 (0.87-0.99)	0.89 (0.78-1.02)	0.97 (0.88-1.08)
31	1.06 (1.04-1.08)	1.24 (1.22-1.27)	1.22 (1.19-1.24)	3.34 (3.08-3.63)	2.01 (1.87-2.17)	1.88 (1.75-2.03)	3.06 (2.83-3.30)	2.14 (1.98-2.31)	2.01 (1.86-2.18)	1.17 (1.07-1.29)	0.93 (0.82-1.06)	0.93 (0.82-1.05)	0.90 (0.82-0.99)	0.94 (0.80-1.10)	1.02 (0.89-1.17)
34	1.11 (1.09-1.13)	1.27 (1.25-1.30)	1.24 (1.22-1.26)	3.58 (3.30-3.88)	2.40 (2.21-2.60)	2.22 (2.04-2.41)	3.19 (2.96-3.44)	2.37 (2.19-2.56)	2.22 (2.06-2.40)	1.24 (1.09-1.40)	1.03 (0.90-1.18)	1.02 (0.89-1.17)	0.87 (0.76-0.99)	1.08 (0.92-1.26)	1.11 (0.95-1.31)
37	1.16 (1.13-1.19)	1.29 (1.26-1.33)	1.26 (1.23-1.29)	3.77 (3.46-4.11)	2.78 (2.54-3.03)	2.54 (2.33-2.78)	3.26 (3.01-3.53)	2.59 (2.39-2.81)	2.43 (2.24-2.64)	1.30 (1.12-1.52)	1.15 (0.98-1.36)	1.14 (0.97-1.34)	0.84 (0.71-0.98)	1.28 (1.09-1.51)	1.26 (1.06-1.51)
40	1.21 (1.18-1.25)	1.32 (1.28-1.36)	1.28 (1.25-1.32)	3.97 (3.61-4.37)	3.12 (2.84-3.44)	2.85 (2.58-3.13)	3.32 (3.03-3.63)	2.83 (2.59-3.10)	2.66 (2.43-2.91)	1.38 (1.14-1.66)	1.29 (1.05-1.58)	1.27 (1.03-1.56)	0.81 (0.67-0.98)	1.52 (1.26-1.84)	1.49 (1.23-1.81)
43	1.27 (1.23-1.32)	1.35 (1.30-1.40)	1.31 (1.26-1.36)	4.18 (3.75-4.67)	3.41 (3.04-3.82)	3.11 (2.77-3.49)	3.38 (3.05-3.75)	3.10 (2.80-3.44)	2.91 (2.62-3.23)	1.45 (1.17-1.80)	1.44 (1.11-1.85)	1.42 (1.10-1.83)	0.78 (0.62-0.98)	1.81 (1.45-2.27)	1.83 (1.47-2.29)
47	1.35 (1.29-1.42)	1.38 (1.32-1.45)	1.34 (1.28-1.40)	4.48 (3.92-5.12)	3.66 (3.12-4.31)	3.37 (2.87-3.96)	3.47 (3.06-3.94)	3.50 (3.08-3.97)	3.27 (2.88-3.72)	1.56 (1.21-2.01)	1.67 (1.20-2.30)	1.64 (1.18-2.27)	0.74 (0.57-0.97)	2.29 (1.72-3.04)	2.56 (1.92-3.41)
50	1.42 (1.35-1.50)	1.41 (1.33-1.48)	1.36 (1.29-1.44)	4.72 (4.05-5.50)	3.74 (3.01-4.64)	3.48 (2.81-4.31)	3.54 (3.06-4.10)	3.83 (3.31-4.43)	3.58 (3.09-4.15)	1.64 (1.23-2.19)	1.86 (1.27-2.72)	1.82 (1.24-2.68)	0.72 (0.53-0.97)	2.72 (1.95-3.80)	3.45 (2.41-4.96)

Notes: the fully adjusted models are adjusted for age, sex, smoking status and the MEDEA deprivation index.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.

Table S4. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, stratified by age and sex**A. Age**

	From general population						From diagnosed with COVID-19						From hospitalised with COVID-19		
	to diagnosed with COVID-19			to hospitalised with COVID-19			to hospitalised with COVID-19			to death			to death		
BMI values (kg/m ²)	Aged 18 to 59	Aged 60 to 79	Aged 80 or older	Aged 18 to 59	Aged 60 to 79	Aged 80 or older	Aged 18 to 59	Aged 60 to 79	Aged 80 or older	Aged 18 to 59	Aged 60 to 79	Aged 80 or older	Aged 18 to 59	Aged 60 to 79	Aged 80 or older
16	0.80 (0.77-0.82)	0.97 (0.95-0.99)	1.23 (1.11-1.36)	0.39 (0.33-0.47)	0.79 (0.76-0.83)	0.84 (0.77-0.90)	0.44 (0.38-0.52)	0.81 (0.77-0.85)	0.91 (0.83-0.99)	6.37 (2.22-18.23)	1.95 (1.39-2.73)	1.05 (0.97-1.14)	0.64 (0.52-0.80)	1.27 (0.99-1.63)	0.98 (0.88-1.09)
19	0.89 (0.88-0.91)	0.98 (0.97-1.00)	1.11 (1.05-1.17)	0.64 (0.60-0.69)	0.89 (0.87-0.91)	0.91 (0.88-0.95)	0.67 (0.62-0.72)	0.90 (0.88-0.92)	0.95 (0.91-1.00)	2.52 (1.49-4.26)	1.40 (1.18-1.65)	1.03 (0.98-1.07)	0.80 (0.72-0.89)	1.10 (0.98-1.23)	0.99 (0.94-1.04)
22	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference
25	1.10 (1.09-1.12)	1.02 (1.00-1.03)	0.91 (0.87-0.96)	1.49 (1.40-1.58)	1.12 (1.10-1.15)	1.09 (1.05-1.14)	1.45 (1.36-1.55)	1.11 (1.08-1.14)	1.05 (1.00-1.10)	0.47 (0.30-0.74)	0.72 (0.61-0.85)	0.97 (0.94-1.02)	1.25 (1.12-1.39)	0.95 (0.87-1.04)	1.01 (0.96-1.07)
28	1.18 (1.16-1.20)	1.03 (1.01-1.06)	0.90 (0.83-0.97)	2.12 (1.91-2.35)	1.26 (1.21-1.32)	1.20 (1.11-1.29)	1.91 (1.72-2.12)	1.24 (1.18-1.30)	1.10 (1.01-1.21)	0.36 (0.18-0.73)	0.59 (0.44-0.78)	0.95 (0.87-1.03)	1.55 (1.25-1.92)	0.95 (0.81-1.12)	1.02 (0.92-1.14)
31	1.23 (1.20-1.25)	1.05 (1.01-1.08)	0.98 (0.91-1.06)	2.88 (2.52-3.29)	1.42 (1.33-1.52)	1.31 (1.17-1.47)	2.26 (2.02-2.53)	1.37 (1.27-1.48)	1.16 (1.01-1.33)	0.46 (0.22-0.98)	0.61 (0.44-0.84)	0.93 (0.82-1.05)	1.93 (1.40-2.66)	1.00 (0.81-1.23)	1.03 (0.88-1.22)
34	1.26 (1.23-1.29)	1.07 (1.02-1.11)	1.13 (1.04-1.23)	3.74 (3.21-4.35)	1.60 (1.46-1.75)	1.43 (1.23-1.67)	2.55 (2.28-2.85)	1.53 (1.38-1.68)	1.22 (1.01-1.46)	0.77 (0.37-1.58)	0.74 (0.54-1.02)	0.90 (0.77-1.07)	2.41 (1.57-3.69)	1.10 (0.86-1.40)	1.05 (0.84-1.30)
37	1.29 (1.26-1.32)	1.08 (1.02-1.15)	1.31 (1.18-1.46)	4.64 (3.95-5.45)	1.80 (1.60-2.02)	1.56 (1.29-1.89)	2.84 (2.54-3.18)	1.70 (1.50-1.92)	1.28 (1.02-1.61)	1.33 (0.65-2.75)	0.96 (0.68-1.35)	0.88 (0.72-1.08)	3.00 (1.76-5.12)	1.27 (0.97-1.65)	1.06 (0.81-1.39)
40	1.32 (1.28-1.37)	1.10 (1.03-1.18)	1.52 (1.33-1.74)	5.50 (4.64-6.52)	2.02 (1.76-2.32)	1.71 (1.36-2.15)	3.17 (2.80-3.59)	1.88 (1.62-2.19)	1.34 (1.02-1.77)	2.32 (1.08-4.97)	1.23 (0.82-1.85)	0.86 (0.67-1.10)	3.74 (1.97-7.09)	1.53 (1.15-2.04)	1.07 (0.77-1.48)
43	1.35 (1.30-1.41)	1.12 (1.03-1.21)	1.76 (1.49-2.08)	6.24 (5.18-7.51)	2.27 (1.93-2.67)	1.87 (1.43-2.44)	3.54 (3.07-4.08)	2.09 (1.76-2.49)	1.41 (1.02-1.94)	4.04 (1.75-9.33)	1.58 (0.96-2.59)	0.84 (0.63-1.12)	4.66 (2.20-9.83)	1.95 (1.42-2.66)	1.08 (0.74-1.58)
47	1.40 (1.33-1.47)	1.14 (1.04-1.25)	2.15 (1.74-2.66)	6.87 (5.43-8.70)	2.65 (2.19-3.22)	2.11 (1.53-2.90)	4.09 (3.45-4.85)	2.41 (1.96-2.96)	1.51 (1.03-2.20)	8.46 (3.18-22.52)	2.20 (1.17-4.13)	0.81 (0.57-1.14)	6.24 (2.56-15.2)	2.88 (1.95-4.24)	1.10 (0.70-1.73)
50	1.43 (1.35-1.52)	1.16 (1.04-1.29)	2.49 (1.95-3.19)	7.01 (5.18-9.48)	2.98 (2.41-3.70)	2.30 (1.61-3.29)	4.56 (3.76-5.55)	2.68 (2.13-3.38)	1.58 (1.03-2.42)	14.72 (4.86-44.52)	2.83 (1.35-5.91)	0.79 (0.54-1.16)	7.77 (2.87-21.06)	4.08 (2.53-6.60)	1.11 (0.67-1.85)


B. Sex

	From general population				From diagnosed with COVID-19				From hospitalised with COVID-19	
	to diagnosed with COVID-19		to hospitalised with COVID-19		to hospitalised with COVID-19		to death		to death	
BMI values (kg/m ²)	Females	Males	Females	Males	Females	Males	Females	Males	Females	Males
16	0.84 (0.82-0.87)	0.74 (0.70-0.77)	0.51 (0.45-0.58)	0.77 (0.73-0.8)	0.49 (0.43-0.57)	0.54 (0.46-0.62)	0.98 (0.90-1.07)	0.93 (0.82-1.06)	1.25 (1.01-1.56)	0.93 (0.84-1.02)
19	0.92 (0.90-0.93)	0.86 (0.84-0.88)	0.73 (0.68-0.77)	0.88 (0.86-0.9)	0.7 (0.66-0.75)	0.73 (0.68-0.79)	0.99 (0.95-1.03)	0.97 (0.91-1.03)	1.12 (1.00-1.25)	0.96 (0.92-1.01)
22	reference	reference	reference	reference	reference	reference	reference	reference	reference	reference
25	1.08 (1.07-1.10)	1.16 (1.13-1.18)	1.34 (1.28-1.40)	1.14 (1.12-1.17)	1.38 (1.3-1.47)	1.35 (1.26-1.45)	1.01 (0.97-1.05)	1.03 (0.97-1.10)	0.90 (0.81-1.00)	1.04 (0.99-1.09)
28	1.14 (1.12-1.17)	1.28 (1.23-1.32)	1.74 (1.60-1.88)	1.3 (1.24-1.37)	1.74 (1.59-1.92)	1.73 (1.54-1.93)	1.02 (0.93-1.11)	1.07 (0.94-1.22)	0.85 (0.70-1.03)	1.08 (0.98-1.19)
31	1.19 (1.16-1.21)	1.33 (1.28-1.37)	2.19 (1.97-2.43)	1.49 (1.39-1.59)	2.01 (1.82-2.23)	2.01 (1.78-2.27)	1.03 (0.90-1.17)	1.11 (0.92-1.34)	0.90 (0.71-1.14)	1.12 (0.96-1.30)
34	1.22 (1.19-1.24)	1.34 (1.29-1.39)	2.67 (2.38-3.01)	1.7 (1.55-1.86)	2.21 (1.99-2.45)	2.25 (2.00-2.52)	1.04 (0.87-1.23)	1.15 (0.89-1.48)	1.07 (0.84-1.36)	1.16 (0.95-1.42)
37	1.24 (1.21-1.28)	1.35 (1.29-1.41)	3.17 (2.80-3.60)	1.94 (1.73-2.18)	2.39 (2.15-2.66)	2.49 (2.20-2.81)	1.04 (0.84-1.29)	1.19 (0.86-1.63)	1.35 (1.06-1.72)	1.21 (0.94-1.55)
40	1.27 (1.23-1.32)	1.36 (1.29-1.43)	3.65 (3.19-4.17)	2.21 (1.93-2.54)	2.59 (2.3-2.92)	2.76 (2.40-3.17)	1.05 (0.82-1.36)	1.23 (0.84-1.80)	1.74 (1.34-2.26)	1.26 (0.93-1.69)
43	1.30 (1.24-1.36)	1.37 (1.28-1.46)	4.08 (3.51-4.74)	2.53 (2.15-2.97)	2.81 (2.45-3.23)	3.06 (2.60-3.59)	1.06 (0.79-1.43)	1.27 (0.82-1.98)	2.23 (1.66-3.00)	1.30 (0.92-1.85)
47	1.34 (1.27-1.42)	1.38 (1.27-1.50)	4.51 (3.69-5.50)	3.02 (2.49-3.66)	3.13 (2.64-3.7)	3.51 (2.88-4.27)	1.08 (0.75-1.54)	1.33 (0.78-2.25)	3.11 (2.16-4.48)	1.37 (0.90-2.08)
50	1.37 (1.28-1.46)	1.39 (1.26-1.53)	4.69 (3.62-6.08)	3.44 (2.77-4.27)	3.39 (2.79-4.11)	3.89 (3.10-4.88)	1.09 (0.73-1.62)	1.38 (0.76-2.48)	4.00 (2.62-6.10)	1.42 (0.89-2.27)

Notes: the models are adjusted for age, sex, smoking status and the MEDEA deprivation index.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.

Figure S4. Association between body mass index and the risk of COVID-19 diagnosis, allowing for non-linear effects, with 95% CIs, stratified by calendar month



Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index. These models were stratified by calendar month to account for the violation of the assumption of proportionality for age in the first transition.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.


Table S5. Hazard Ratios of the risk of COVID-19 diagnosis related to body mass index, allowing for non-linear effects, with 95% CIs, stratified by calendar month

	From general population to diagnosed with COVID-19		
BMI values (kg/m²)	Overall	March	April
16	0.81 (0.79-0.84)	0.78 (0.75-0.81)	0.86 (0.83-0.90)
19	0.90 (0.89-0.91)	0.88 (0.87-0.90)	0.93 (0.91-0.95)
22	reference	reference	reference
25	1.10 (1.09-1.11)	1.12 (1.10-1.14)	1.07 (1.05-1.09)
28	1.17 (1.15-1.19)	1.21 (1.18-1.24)	1.13 (1.10-1.16)
31	1.22 (1.19-1.24)	1.25 (1.22-1.28)	1.17 (1.14-1.20)
34	1.24 (1.22-1.26)	1.27 (1.24-1.31)	1.20 (1.16-1.24)
37	1.26 (1.23-1.29)	1.29 (1.25-1.33)	1.23 (1.19-1.28)
40	1.28 (1.25-1.32)	1.31 (1.26-1.36)	1.26 (1.21-1.32)
43	1.31 (1.26-1.36)	1.33 (1.26-1.39)	1.30 (1.23-1.37)
47	1.34 (1.28-1.40)	1.35 (1.27-1.43)	1.34 (1.25-1.44)
50	1.36 (1.29-1.44)	1.37 (1.27-1.47)	1.38 (1.27-1.50)

Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index. These models were stratified by calendar month to account for the violation of the assumption of proportionality for age in the first transition.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.


Figure S5. Hazard Ratios of COVID-19 outcomes related to body mass index in categories, with 95% CIs



Notes: Hazard ratios are relative to normal or underweight (BMI <18.5 kg/m² and between ≥18.5 and <25 kg/m²). The overweight category includes BMIs between ≥25 and <30 kg/m² and for the obesity category BMIs ≥30 kg/m². Models are adjusted for age, sex, smoking status and the MEDEA deprivation index.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019; HR: Hazard ratio.

Figure S6. Association between body mass index and the risk of COVID-19 outcomes, allowing for non-linear effects, with 95% CIs, after multiple imputation of missing data on BMI, smoking status, and/or the MEDEA deprivation index



Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index. For the multiple imputations we used predictive mean matching, with 5 imputations drawn.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”.


Table S6. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs, after multiple imputation of missing data on BMI, smoking status, and/or the MEDEA deprivation index

	From General Population		From diagnosed with COVID-19		From hospitalised with COVID-19
BMI values (kg/m ²)	To diagnosed with COVID-19	To hospitalised with COVID-19	To hospitalised with COVID-19	To death	To death
16	0.84 (0.79-0.88)	0.58 (0.53-0.64)	0.49 (0.44-0.53)	1.34 (1.18-1.52)	1.09 (0.96-1.24)
19	0.91 (0.89-0.94)	0.77 (0.74-0.81)	0.70 (0.67-0.73)	1.16 (1.09-1.23)	1.03 (0.97-1.09)
22	reference	reference	reference	reference	reference
25	1.09 (1.06-1.11)	1.26 (1.22-1.30)	1.39 (1.34-1.45)	0.88 (0.84-0.93)	0.99 (0.95-1.04)
28	1.15 (1.12-1.18)	1.55 (1.48-1.63)	1.77 (1.67-1.88)	0.85 (0.78-0.92)	1.01 (0.94-1.10)
31	1.20 (1.16-1.23)	1.86 (1.75-1.98)	2.04 (1.91-2.17)	0.88 (0.81-0.96)	1.06 (0.95-1.17)
34	1.23 (1.19-1.27)	2.18 (2.04-2.34)	2.23 (2.09-2.37)	0.96 (0.87-1.05)	1.13 (1.00-1.28)
37	1.26 (1.20-1.33)	2.50 (2.32-2.69)	2.41 (2.26-2.57)	1.05 (0.93-1.18)	1.24 (1.08-1.42)
40	1.30 (1.20-1.39)	2.78 (2.55-3.04)	2.61 (2.42-2.81)	1.14 (0.98-1.33)	1.40 (1.20-1.62)
43	1.33 (1.21-1.46)	3.03 (2.69-3.41)	2.82 (2.59-3.08)	1.25 (1.04-1.51)	1.61 (1.35-1.92)
47	1.38 (1.22-1.56)	3.26 (2.71-3.93)	3.14 (2.83-3.49)	1.41 (1.11-1.79)	2.02 (1.60-2.55)
50	1.41 (1.22-1.64)	3.35 (2.61-4.31)	3.40 (3.01-3.84)	1.54 (1.17-2.04)	2.47 (1.84-3.31)

Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index. For the multiple imputations we used predictive mean matching, with 5 imputations drawn.

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019; MEDEA: “Mortalidad en áreas pequeñas Españolas y Desigualdades Socioeconómicas y Ambientales”.

Figure S7. Association between body mass index and the risk of COVID-19 outcomes, allowing for non-linear effects, with 95% CIs among people with a BMI measurement not older than five years prior to the index date



Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index
 Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.

Table S7. Hazards ratios of COVID-19 outcomes related to body mass index, with 95% CIs among people with a BMI measurement not older than five years prior to the index date

	From General Population		From diagnosed with COVID-19		From hospitalised with COVID-19
BMI values (kg/m ²)	To diagnosed with COVID-19	To hospitalised with COVID-19	To hospitalised with COVID-19	To death	To death
16	0.81 (0.79-0.84)	0.61 (0.55-0.67)	0.55 (0.49-0.61)	1.32 (1.11-1.58)	1.20 (1.01-1.43)
19	0.90 (0.89-0.92)	0.79 (0.75-0.82)	0.74 (0.70-0.78)	1.15 (1.05-1.26)	1.08 (0.99-1.16)
22	reference	reference	reference	reference	reference
25	1.10 (1.09-1.11)	1.25 (1.20-1.29)	1.33 (1.27-1.40)	0.89 (0.82-0.96)	0.97 (0.91-1.03)
28	1.18 (1.16-1.21)	1.52 (1.43-1.62)	1.68 (1.55-1.81)	0.85 (0.75-0.96)	0.97 (0.87-1.08)
31	1.23 (1.21-1.26)	1.82 (1.68-1.96)	1.95 (1.79-2.13)	0.89 (0.77-1.02)	1.01 (0.88-1.17)
34	1.26 (1.24-1.29)	2.12 (1.95-2.32)	2.17 (1.99-2.36)	0.99 (0.85-1.14)	1.10 (0.93-1.30)
37	1.29 (1.26-1.33)	2.43 (2.22-2.67)	2.38 (2.18-2.59)	1.12 (0.93-1.33)	1.24 (1.03-1.49)
40	1.32 (1.28-1.36)	2.73 (2.47-3.02)	2.61 (2.37-2.87)	1.26 (1.01-1.58)	1.45 (1.18-1.78)
43	1.35 (1.30-1.40)	3.00 (2.66-3.38)	2.86 (2.56-3.20)	1.42 (1.08-1.88)	1.76 (1.40-2.22)
47	1.39 (1.32-1.46)	3.29 (2.79-3.89)	3.24 (2.83-3.71)	1.67 (1.17-2.39)	2.43 (1.80-3.28)
50	1.42 (1.34-1.50)	3.45 (2.76-4.30)	3.55 (3.05-4.15)	1.89 (1.25-2.87)	3.24 (2.22-4.71)

Notes: Models are adjusted for age, sex, smoking status and the MEDEA deprivation index

Abbreviations: BMI: Body mass index; CI: Confidence interval; COVID-19: Coronavirus Disease 2019.