

Table S2. Multiple linear regression models assessing the unique contribution of R1 and FA to reading outcome (Oral Reading Index) at age 8 years in A. the left superior longitudinal fasciculus (SLF-L). B. the right uncinate fasciculus (UF-R), and C. the left inferior longitudinal fasciculus (ILF-L).

A. SLF nodes 1-5	Model A1	Model A2
R1	0.42	0.39
FA		0.068
ΔR^2		0.003 (p = 0.74)
Total R ²	0.18 (p = 0.02)	0.18 (p = 0.07)
Adj R ²	0.15	0.12
B. UF-R nodes 13-19	Model B1	Model B2
R1	0.45	0.40
FA		0.17
ΔR^2		0.02 (p = 0.37)
Total R ²	0.20 (p = 0.015)	0.23 (p = 0.04)
Adj R ²	0.17	0.17
C. ILF-L nodes 20-23	Model C1	Model C2
R1	0.53	0.56
FA		-0.31
ΔR^2		0.09 (p = 0.06)
Total R ²	0.28 (p = 0.003)	0.37 (p = 0.002)
Adj R ²	0.25	0.32

Data are standardized coefficients. R² change (ΔR^2) values in Model 2 reflect the increase in variance accounted for by FA in addition to R1 (Model 1).