





Supplemental Tables and Figures—


High Throughput Screening for Expanded CTG Repeats in Myotonic Dystrophy Type 1 Using Melt Curve Analysis. RJ Butterfield et al.

Supplemental Figure 1. Template for scoring TP-PCR melt curves.


Four examples of typical melt curves as reviewed by the blinded reviewers. Each sample classified into one of six categories based on visual inspection of the MCP: Normal, Intermediate, Premutation, Expanded, Uncertain, and Fail. The melt curve for the DBS samples (black) is plotted with averaged melt curves from the four positive (expanded) controls with (CTG)_n ≥ 50 [5::75 (green), 5::80 (purple), 5::480 (light red), and 14:2530 (light blue)].



Supplemental Figure 2. Melt curve profiles from all samples classified as ‘expanded’ by blinded scoring. Each sample was reviewed by 4 blinded reviewers, with 16 samples resulting in a ‘expanded’ score based on the consensus of the reviewers. The melt curve for the DBS sample (black) is plotted with averaged melt curves from four positive (expanded) controls with (CTG)_n ≥ 50: 5::75 (green), 5::80 (purple), 5::480 (light red), and 14:2530 (light blue). Repeat numbers for Allele 1:: Allele 2 by direct PCR or the Amplidex® DM1 Dx Kit are shown above each plot. Samples with true expansions are highlighted in yellow.



Supplemental Figure 3. Individual scores of melt curve profiles and validated CTG repeat size from all samples classified as ‘premutation’ by blinded scoring. Heatmap of individual calls from the four blinded reviewers and final CTG repeat numbers for Allele 1 :: Allele 2 by direct PCR or the Amplidex® DM1 Dx Kit are shown for each sample, with the true premutations boxed in yellow.



Supplemental Table 1. Individual allele sizes and blinded scores from for control samples.

Sample ID	Diplotype Group [†]	Allele 1 (CTG) _n	Allele 2 (CTG) _n	Consensus Score from Blinded Review
1	(-+-)	5	5	Normal
2	(-+-)	5	5	Normal
3	(-+-)	5	5	Normal
4	(-+-)	5	5	Normal
5	(-+-)	5	5	Normal
6	(-+-)	5	5	Normal
7	(-+-)	5	5	Normal
8	(-+-)	5	5	Normal
9	(-+-)	5	5	Normal
10	(-+-)	5	5	Normal
11	(-+-)	5	5	Normal
12	(-+-)	5	5	Normal
13	(-+-)	5	5	Normal
14	(-+-)	5	5	Normal
15	(-+-)	5	5	Normal
16	(+++)	8	13	Normal
17	(+++)	11	11	Normal
18	(+++)	12	12	Normal
19	(+++)	12	12	Normal
20	(+++)	13	13	Normal
21	(+++)	11	13	Normal
22	(+++)	12	13	Normal
23	(+++)	13	13	Normal
24	(+++)	12	13	Normal
25	(+++)	11	13	Normal
26	(+++)	12	13	Normal
27	(+++)	12	13	Normal
28	(+++)	12	13	Normal
29	(+++)	11	13	Normal
30	(+++)	14	14	Normal
31	(---)	14	14	Normal
32	(---)	14	17	Normal
33	(---)	14	21	Intermediate
34	(---)	14	22	Intermediate
35	(---)	22	23	Intermediate
36	(---)	14	25	Intermediate
37	(---)	14	25	Intermediate

38	(---)	10	25	Intermediate
39	(---)	14	26	Intermediate
40	(---)	24	26	Intermediate
41	(---)	27	28	Intermediate
42	(---)	26	29	Premutation
43	(---)	12	37	Premutation
44	(---)	14	37	Premutation
45	(---)	14	40	Premutation

† Expected range for (CTG)_n in the (-+-) haplotype=5, in the (+++) haplotype= 8-17 and in the (---) haplotype = 18-35.