## **Supplementary Methods**

## Mindfulness-based Intervention

The MBI program used in this study was specifically adapted to adolescents. For each session one theme was addressed, such as attention and the stabilisation of the focus of attention, bodily sensations, breath, emotions, thoughts, compassion, stress, stress reactivity and coping strategies. Different formal meditation practices were introduced, such as sitting mediation with different objects of attention, body scan, walking meditation and mindful movement. Participants had the opportunity to share experiences with their peers and the instructors during each session, allowing the recognition of individual patterns of behaviour, and at the same time, the appreciation of the shared experiences as a community. They were also invited to practice between 5 to 20 minutes per day at home, using guided mediation recorded by the instructors, see Siffredi\*, Liverani\* & al., under revision, provided in supplementary file.

## Supplementary Table S1. Details of the RCT outcome measures

Domains	Modalities	Measures	Description	Scores
Executive	competences			
	Parent quest	ionnaire		
		Behaviour l	Rating Inventory of Executive Function, parent version (BRIEF, Gioia, Isquith, Guy, and Kenworthy (2000))	
			The BRIEF parent questionnaire provides an index of attention, hyperactivity and impulsivity in everyday	BRIEF CEG
			life. The BRIEF comprises 86 items over two standardised subscales: (i) Behavioural Regulation Index (BRI)	BRIEF BRI
			comprising 3 subscores including, Inhibit, Shift, Emotional Control; (ii) Metacognition index (MCI)	BRIEF MCI
			comprising 5 subscores including, Initiate, Working Memory, Plan/Organise, Organisation of Materials,	

Neuropsychological tests	
Letter-Number Sequencing (WISC-IV, Wechsler (2003))	
	Letter-number
participant, and he/she is then asked to re-sequence the numbers in numerical order from lowest to highest	sequencing
and then to sequence the letters in alphabetical order. Standardised scores were used.	
Tempo Test Rekenen (De Vos, 1992)	
	Tempo test
five rows (one row with addition, one row with subtraction, one row with division, one row with	
multiplication, and one mixed problem row). Within each row, the problems increase in difficulty. Participant	
are asked to solve as many items as possible within 1 min per row. The total raw score was age-adjusted for	
each participant using the procedure described in the main statistical analyses section.	
Neurocognitive computerised tasks	
Flanker Visual Filtering Task (Christ, Kester, Bodner, & Miles, 2011)	
The Flanker Visual Filtering Task was used to assess attentional control and information processing speed.	-Flanker
Each trial showed a horizontal row of five fish. The participant was asked to respond as quickly as possible to	processing spee
whether the central fish was facing to the left or right. Congruent trials were the ones with all five fish in the	-Flanker inhibit
horizontal row pointing in the same direction and incongruent trials were the ones with the four distracting	
fishes pointing in the opposite direction of the central target fish. Reaction time of the congruent condition	
and of the incongruent condition were used to assess information processing speed, and the inhibition score	
(reaction time in incongruent conditions – reaction time in congruent conditions) was used as a measure of	
attentional control.	
Reality Filtering Task (Liverani et al., 2017; Schnider, 2018)	
	Reality filtering
	TCC
but arranged in different order. Accuracy of the second run (D2) and Temporal Context Confusion index	
(TCC as defined by Schnider, 2018) measures reality filtering.	
Behaviour and socio-emotional competences	
Parent questionnaire	
Strength and Difficulties Questionnaire, parent version (SDQ, Goodman (2001))	an a
	SDQ total
inattention, peer relationship problems, and prosocial behaviour. It rates participant's behaviour over the	
previous 6 months. The SDQ is scored on a Likert scale and includes 25 items, providing a Total Difficulties	
score. Self-reported questionnaires	
KIDSCREEN-27 (Robitail et al., 2007)	

children and adolescents. This instrument scored on a Likert scale and includes 27 items, providing a total score.

Social Goal Scale (SGS, Patrick, Hicks, and Ryan (1997))

The SGS is a self-reported questionnaire providing an index of social responsiveness and of goals setting which ultimately gets you involve with some social work. This instrument scored on a Likert scale and includes 11 items providing one score.

Self-Compassion Scale – Short form (SCS, Raes, Pommier, Neff, and Van Gucht (2011))

The SCS is a self-reported questionnaire comprising 12 items, which produces a total global score that can also be classified into two subscores: negative behaviours toward the self (6 items) or positive behaviours towards the self (6 items).

Self-compassion

Affect recognition

Neuropsychological tests

Affect Recognition (NEPSY-II, Korkman, Kirk, and Kemp (2007)

The affect recognition subtest assesses the ability to recognise facial emotional expressions (happy, sad, anger, fear, disgust, and neutral) from photographs of children's faces in several matching tasks. In the first task, the participant selected one of the four faces that depicted the same emotion as a child's face at the top of the page. In a second task, the participant selected two photographs of faces that displayed the same affect from a selection of four photographs. Finally, the participant examined a photograph of a child's face for 5 seconds, and then from memory, selected two photographs that matched the same emotion as the face previously shown. Standardised scores were used.

Theory of Mind (NEPSY-II, Korkman et al. (2007))

The theory of mind subtest measures understanding of mental functions and other people's perspectives. In the first task, questions are asked to the participant about different verbal scenarios measuring understanding of beliefs, intentions, others' thoughts, ideas and comprehension of figurative language. In the second task, participants have to match facial emotional expressions, from photographs of children's faces, to a scenario. The total raw score was age-adjusted for each participant using the procedure described in the main statistical analyses section.

Theory of mind

Supplementary Table S2. Randomised controlled trial timing

	RCT, n=56				
	Intervention group (IG), n=29	Waiting group (WG), n=27	Group comparison (IG vs WG)		
Difference between Time 1 and Time 2, mean (SD) in days	95.27 (19.9)	90 (30.9)	t(34.73)=0.688, p=0.496		
Difference between Time 2 and Time 3, mean (SD) in days	81.23 (21.1)	84.64 (13.4)	t(46.98)= -0.654, p=0.502		

Note: Independent-sample t-test was used to compare the randomised groups.

**Supplementary Table S3**. Linear model results of the intervention (IG) and wait (WG) for the delta  $\Delta 1$  and  $\Delta 2$  for the Intent-to-Treat Sample. Delta mean and standard deviation (SD) are also showed.

Outcomes measures	Interaction effect time (Δ1 and Δ2) by group (IG and WG)	Planned contrast: MBI VS treatment as	Planned contrast: long-term VS	Planned contrasts Mean Δ (SD)			
		usual	treatment as usual	Treatment as usual	МВІ	Long-term	
Executive competences							
BRIEF CEG	F(3,80)=4.441, p=0.006, d=0.5, q=0.026	t(80)=-3.195,p=0.002	t(80)= -0.589, p=0.558	5.16 (17.68)	-10.79 (19.43)	1.87 (14.89)	
BRIEF MI	F(3,80)=6.72, p<0.001, d=0.615, q=0.006	t(80)= -3.985, p<0.001	t(80)=-0.852, p=0.397	4.58 (12.39)	-8.31 (11.99)	1.48 (10.49)	
BRIEF BRI	F(3,80)=1.47, p=0.229, d=0.288, q=0.368	t(80)= -1.327,p=0.188	t(80)=- 0.075, p=0.941	0.58 (6.93)	-2.48 (9.17)	0.39 (6.96)	
Letter-Number sequencing	F(3,92)=0.155, p=0.962, d=0.093, q=0.962	t(92)=0.332, p=0.74	t(92)=0.674, p=0.502	0.41 (2.36)	0.58 (1.76)	0.81 (2.19)	
Tempo test	F(3,91)=0.667, p=0.574, d=0.194, q=0.662	t(91)=0.511, p=0.611	t(91)= -0.729, p=0.468	4.55 (8.52)	5.83 (8.77)	2.42 (12.87)	
Flanker processing speed	F(3,90)=4.83, p=0.004, d=0.521, q=0.022	t(90)=-3.789, p<0.001	t(90)=-2.620, p=0.01	116.02 (256)	-140.8 (290.02)	-82.44 (180.99)	
Flanker inhibition	F(3,90)=0.57, p=0.636, d=0.179, q=0.707	t(90)=-0.335, p=0.739	t(90)= 0.713, p=0.478	-12.77 (97.33)	-26.61 (171.34)	18.69 (141.77)	
Reality filtering, TCC	F(3,86)=0.691, p=0.560, d=0.15, q=0.368	t(86)=-0.795, p=0.422	t(86)= -1.394, p=0.167	0.07 (0.22)	0.04 (0.14)	0.01 (0.1)	
Socio-emotional competend	ees						
SDQ total	F(3,88)=3.518, p= 0.018, d=0.445, q=0.048	t(88)=-2.423, p=0.017	t(88)=-0.178, p=0.859	0.59 (3.86)	-2.02 (4.21)	0.38 (4.18)	
KIDSCREEN total	F(3,84)=0.401, p=0.752, d=0.15, q=0.806	t(84)=-0.045 p=0.964	t(84)=0.854, p=0.396	0.52 (13.37)	0.39 (10.21)	3.48 (11.64)	
Social goal	F(3,82)=3.812, p=0.013, d=0.463, q=0.044	t(82)=-0.353, p=0.725	t(82)=0.890, p=0.376	0.08 (0.31)	0.04 (0.52)	0.19 (0.41)	
Self-compassion	F(3,82)=2.693, p=0.051, d=0.389, q=0.110	t(82)=2.181, p=0.032	t(82)=2.763, p=0.007	-0.18 (0.43)	0.15 (0.55)	0.29 (0.66)	
Affect recognition	F(3,92)=0.951, p=0.419, d=0.231, q=0.571	t(92)=-0.248, p=0.804	t(92)=-1.157, p=0.25	1.32 (3.31)	1.15 (3.49)	0.19 (3.14)	
Theory of mind	F(3,89)=1.061, p=0.37, d=0.244, q=0.529	t(89)=0.927, p=0.356	t(89)=-0.546, p=0.586	0.52 (1.91)	1.02 (2.08)	0.2 (1.89)	

Abbreviations: intervention group (IG), waiting group (WG); delta scores: difference between Time 1 and Time 2 ( $\Delta$ 1), and between Time 2 and Time 3 ( $\Delta$ 2). Note: d shows Cohen's d effect sizes. All p-values of linear models that survived false discovery rate (FDR) correction according to Benjamini and Hochberg (1995) are indicated in bold (q < 0.05). Negative  $\Delta$  indicate a reduction of the scores between the 2 time points whereas positive  $\Delta$  indicate an increase of the scores between the 2 time points.

## Supplementary Table S4. Planned contrast MBI VS long-term of the significant planned contrast: MBI VS treatment as usual

Outcomes measures	Significant planned contrast: MBI VS treatment as usual (reported in supplementary table 3)	Planned contrast: MBI VS long-term		
BRIEF CEG	t(80)=-3.195,p=0.002	t(80)= -2.702, p=0.008		
BRIEF MI	t(80)= -3.985, p<0.001	t(80)=-3.229, p=0.002		

Flanker processing speed	t(90)=-3.789, p<0.001	t(90)= -0.926, p=0.3571
SDQ total	t(88)=-2.423, p=0.017	t(88)=-2.285, p=0.025

**Supplementary Table S5**. Subgrouping "prematurity": linear model results of the intervention (IG) and wait (WG) for the delta  $\Delta 1$  and  $\Delta 2$  for the Intent-to-Treat Sample, as well as delta mean and standard deviation (SD)

Outcomes measures	Interaction effect: time ( $\Delta 1$ and $\Delta 2$ ) by	Planned contrast: MBI VS treatment as usual		Planned contrast: long-term VS treatment as usual		Planned contrasts Mean Δ (SD)			Planned contrasts Mean Δ (SD)		
Outcomes measures	group (IG and WG) by subgrouping	Moderate-risk group	High-risk group	Moderate-risk group	High-risk group	Moderate-risk group			High-risk group		
	prematurity (high and moderate-risk)					Treatment as usual	MBI	Long term	Treatment as usual	MBI	Long term
Executive competences							•				
BRIEF CEG	F(7,76)=2.622, d= 0.384, p=0.018, q=0.048	t(76)=-1.967, p=0.053	t(76)=-2.625, p=0.011	t(76)=-0.911, p=0.365	t(76)=-1.920, p=0.059	4.1 (14.22)	-9.55 (22.94)	-3.2 (11.22)	6.33 (21.74)	-11.91 (16.06)	5.77 (16.55)
BRIEF MI	F(7,76)= 3.228, d=0.426, p=0.005,q=0.024	t(76)=-3.098,p=0.003	t(76)=-2.462,p=0.016	t(76)=-1.568, p=0.121	t(76)=-2.216, p=0.03	6.5 (10.89)	-7.7 (14.02)	-1.8 (6.49)	2.44 (14.22)	-8.86 (10.11)	4 (12.42)
BRIEF BRI	F(7,76)=2.484, d=0.374, p=0.024,q=0.055	t(76)=0.184, p=0.855	t(76)=-2.318, p=0.023	t(76)=0.290, p=0.773	t(76)=-1.058, p=0.293	-2.4 (4.72)	-1.85 (10.59)	-1.4 (7.56)	3.89 (7.72)	-3.05 (7.88)	1.77 (6.42)
Letter-Number sequencing	F(7,88)=1.567, d=0.297, p=0.156, q=0.275	t(88)= 2.169, p=0.0328	t(88)= -1.512, p=0.134	t(88)=2.724, p=0.008	t(88)=0.383, p=0.702	-0.8 (1.55)	0.82 (1.56)	1.5 (2.71)	1.42 (2.5)	0.38 (1.92)	0.21 (1.48)
Tempo test	F(7,87)=2.944, d=0.407, p=0.008, q=0.031	t(87)= 1.986, p=0.0501	t(87)= -1.065, p=0.3	t(87)=-1.558, p=0.123	t(87)=-1.881, p=0.063	2.9 (9.87)	10.19 (6.8)	-3.33 (15.29)	5.92 (7.38)	2.31 (8.69)	7.36 (8)
Flanker processing speed	F(7,86)=3.481, d=0.443, p=0.003, q=0.019	t(86)= -3.354, p=0.001	t(86)= -2.142, p=0.035	t(86)=-3.341, p=0.001	t(86)=-1.246, p=0.216	250.57 (265.49)	-73.35 (219.92)	-108.78 (221.17)	-6.29 (181.86)	-200.16 (333.25)	-59.87 (142.88)
Flanker inhibition	F(7,86)=0.49, d=0.166, p=0.84, q=0.869	t(86)= 0.409, p=0.683	t(86)=-0.831, p=0.408	t(86)=0.313, p=0.755	t(86)=-1.516, p=0.133	-27.43 (72.64)	-5.26 (109.19)	-7.02 (67.63)	0.56 (117.42)	-45.39 (212.25)	40.73 (183.44)
Reality filtering, TCC	F(7,82)= 0.433, d=0.215, p=0.879, q=0.662	t(82)= -0.615, p=0.540	t(82)= -0.499, p=0.619	t(82)=-1.538, p=0.128	t(82)=-0.297, p=0.767	0.06 (0.19)	0.03 (0.14)	0.03 (0.06)	0.08 (0.26)	0.05 (0.15)	-0.02 (0.12)
Socio-emotional competences											
SDQ total	F(7,84)= 1.837, d=0.321, p=0.091, q=0.170	t(84)=-1.908, p=0.06	t(84)=-1.509, p=0.1350	t(84)=-0.966, p=0.337	t(84)=-2.182, p=0.032	1.2 (2.53)	-1.9 (4.45)	-0.55 (4.06)	0.08 (4.76)	-2.12 (4.09)	1.15 (4.28)
KIDSCREEN total	F(7,80)=1.246, d=0.265, p=0.288, q=0.432	t(80)=-1.454, p=0.15	t(80)=1.161, p=0.249	t(80)=0.076, p=0.94	t(80)=0.097, p=0.923	6.89 (13.25)	0.38 (9.7)	7.27 (11.43)	-4.25 (11.79)	0.39 (10.87)	0 (11.17)
Social goal	F(7,78)=1.989, d=0.334, p=0.067, q=0.134	t(78)= -0.482, p=0.631	t(78)= -0.085, p=0.932	t(78)=0.756, p=0.452	t(78)=0.421, p=0.675	0.16 (0.39)	0.09 (0.58)	0.31 (0.32)	0.01 (0.22)	-0.01 (0.46)	0.09 (0.46)
Self-compassion	F(7,78)=6.135, d=0.587, p<0.001, q<0.001	t(78)= 0.540, p=0.591	t(78)= 3.015, p=0.004	t(78)=3.165, p=0.002	t(78)=2.743, p=0.008	-0.02 (0.38)	0.07 (0.64)	0.67 (0.64)	-0.32 (0.43)	0.22 (0.46)	-0.06 (0.48)
Affect recognition	F(7,88)=0.823, d=0.215, p=0.571, q=0.662	t(88)=0.459, p=0.648	t(88)=-0.751, p=0.454	t(88)=-1.060, p=0.292	t(88)=0.447, p=0.656	0.7 (2.36)	1.32 (2.03)	-0.83 (1.7)	1.83 (3.97)	1 (4.41)	1.07 (3.83)
Theory of mind	F(7,85)= 0.887, d=0.223, p=0.521, q=0.662	t(85)=0.910, p=0.365	t(85)=0.457, p=0.649	t(85)=-0.682, p=0.497	t(85)=0.282, p=0.779	0.6 (2.55)	1.33 (2.11)	0 (1.48)	0.45 (1.21)	0.77 (2.07)	0.36 (2.21)

Abbreviations: intervention group (IG), waiting group (WG); delta scores: difference between Time 1 and Time 2 ( $\Delta$ 1), and between Time 2 and Time 3 ( $\Delta$ 2). Note: d shows Cohen's d effect sizes. All p-values of linear models that survived false discovery rate (FDR) correction according to Benjamini and Hochberg (1995) are indicated in bold (q < 0.05). Negative  $\Delta$  indicate a reduction of the scores between the 2 time points whereas positive  $\Delta$  indicate an increase of the scores between the 2 time points.

Supplementary Table S6. Subgrouping "prematurity": Planned contrast MBI VS long-term of the significant planned contrast: MBI VS treatment as usual

Outcomes measures	Significant planned contrast: (reported in supple		Planned contrast: MBI VS long-term		
	Moderate-risk group	High-risk group	Moderate-risk group	High-risk group	
BRIEF CEG	t(76)=-1.967, p=0.053	t(76)=-2.625, p=0.011		t(76)= -2.879, p=0.005	
BRIEF MI	t(76)=-3.098,p=0.003	t(76)=-2.462,p=0.016	t(76)= -1.287 0.2019	t(76)= -3.159, p=0.002	
Flanker processing speed	t(86)= -3.354, p=0.001	t(86)= -2.142, p=0.035	t(86)= 0.416 0.6782	t(86)= -1.682, p=0.096	
Self-compassion	t(78)= 0.540, p=0.591	t(78)= 3.015, p=0.004		t(78)= 1.602, p=0.113	