

Table S1: Attachment variables, corresponding to the three questions asked at the three different ages. This includes the participants who answered with “never apart”. Sample is based on the core ALSPAC sample (n=14,732).

Age (months)	Question 1: Avoids contact with mother/carer				Question 2: Pushes mother/carer away				Question 3: Runs for hug			
	n (%)				n (%)				n (%)			
	Hardly ever	Sometimes	Always/ usually*	Never apart	Hardly ever	Sometimes	Always/ usually*	Never apart	Always/ Usually*	Sometimes	Hardly ever	Never apart
18	8115 (86.56)	601 (6.41)	61 (0.65)	598 (6.38)	8233 (88.1)	474 (5.07)	43 (0.46)	598 (6.40)	8947 (81.2)	1283 (11.7)	188 (1.71)	598 (5.43)
30	9708 (94.9)	505 (4.94)	15 (0.15)	n/a**	9681 (94.7)	529 (5.18)	9 (0.09)	n/a**	7695 (75.5)	2380 (23.3)	124 (1.22)	n/a**
42	9564 (95.68)	328 (3.28)	5 (0.05)	99 (0.99)	9571 (95.75)	318 (3.18)	8 (0.08)	99 (0.99)	7505 (75.1)	2257 (22.6)	135 (1.35)	99 (0.99)

Note: * on 18 months questionnaire, the category ‘usually’ was asked, on the 30 and 42 months questionnaire it was re-worded to ‘always’; ** “never apart” was not an option on the questionnaire at 30 months; categories were collapsed into the above one if the number of observations was <15.

Table S2: Structural equation modelling of attachment (standardised coefficients)

Latent Variables		
First order model	Questionnaire item*	DWLS
Attachment at 18 months	Frequency infant avoids contact with mother after separation	1.000
	Frequency infant pushes mother away after separation	0.867
	Frequency infant cuddles mother after separation**	0.893
Attachment at 30 months	Frequency infant avoids contact with mother after separation	0.976
	Frequency infant pushes mother away after separation	0.873
	Frequency infant cuddles mother after separation**	1.000
Attachment at 42 months	Frequency infant avoids contact with mother after separation	1.002
	Frequency infant pushes mother away after separation	0.969
	Frequency infant cuddles mother after separation**	1.000
Second order model	First order model latent variable	

Attachment over the three

ages

Attachment at 18 months	0.965
Attachment at 30 months	1.292
Attachment at 42 months	1.000

Note: * Based on questions: when you and your toddler meet again after being apart for an hour or more, how often

((a) always/usually, (b) sometimes, or (c) hardly ever) do they: (1) move away or avoid looking at you (2) push you

away (3) run to you for a hug or cuddle?; **reverse coded so remains consistent with other measures that higher

scores = poor attachment; DWLS estimate = diagonally weighted least squares.

Fig S1: Flowchart of ALSPAC participants showing attrition.




Table S3 : Selective attrition for participants who attended the 18-year clinic.

		Available/ attended clinic (n= 4878*)	Not available/ did not attend clinic (n= 9854*)	OR (95% CI)
		n (%)	n (%)	
Gender				
	Female	2743 (56.2)	4079 (41.4)	0.61 (0.57, 0.66)
Housing tenure				
High	Owned/mortgaged	3978 (81.5)	5894 (59.8)	Ref
	Private rented	216 (4.43)	762 (7.73)	2.38 (2.04, 2.78)
Low	Council/housing association rented	408 (8.36)	1757 (17.8)	2.91 (2.59, 3.26)
Parental social class				
High	I	800 (16.4)	738 (7.49)	Ref
	II	2055 (42.1)	2773 (28.1)	1.46 (1.30, 1.64)
	III (non-manual)	1042 (21.4)	1904 (19.3)	1.98 (1.75, 2.25)
	III (manual)	408 (8.36)	1155 (11.7)	3.07 (2.64, 3.57)
	IV	162 (3.32)	421 (4.27)	2.82 (2.29, 3.46)
Low	V	24 (0.49)	77 (0.78)	3.48 (2.18, 5.56)
Maternal education				
High	Degree	872 (17.9)	737 (7.48)	Ref
	A level	1308 (26.8)	1485 (15.1)	1.34 (1.19, 1.52)
	O level	1611 (33.0)	2712 (27.5)	1.99 (1.77, 2.24)
	Vocational	355 (7.28)	874 (8.87)	2.91 (2.49, 3.41)

Low	Secondary education	554 (11.4)	1971 (20.0)	4.21 (3.67, 4.82)
Maternal alcohol usage in pregnancy				
	Never	2063 (42.3)	3932 (39.9)	Ref
	< 1 glass per week	1958 (40.1)	3141 (31.9)	0.84 (0.78, 0.91)
	1 glass per day	647 (13.3)	1186 (12.0)	0.96 (0.86, 1.07)
	1-2 glasses per day	72 (1.48)	140 (1.42)	1.02 (0.76, 1.36)
	≥3 glasses per day	15 (0.31)	27 (0.27)	0.94 (0.50, 1.78)
Maternal smoking in pregnancy				
	Yes	793 (16.3)	2570 (26.1)	2.15 (1.97, 2.35)
Parental separation or divorce				
	Yes	228 (4.67)	493 (5.00)	1.55 (1.32, 1.83)
		Mean (SD)	Mean (SD)	
Maternal age	(linear scale)	29.7 (4.58)	28.0 (4.9)	0.93 (0.92, 0.94)
Number of children in household < 15 years old	(linear scale)	1.75 (0.99)	1.90 (0.96)	1.18 (1.13, 1.23)
Maternal depression	(linear scale 0-29)	5.10 (4.49)	5.63 (4.82)	1.02 (1.02, 1.03)
Partner affection	(linear scale 0-49)	21.45 (7.06)	21.72 (7.38)	1.01 (0.99, 1.01)

Note: *from core ALSPAC sample n=14732; highest of either parent's social class based on 5 employment categories: I = professional; II = managerial or technical, III = skilled manual or non-manual, IV = unskilled manual, V = casual and lowest grade work; maternal depression as measured by the validated scale Edinburgh Postnatal Depression Score at 8 months post-natal;

partner affection score (created from individual questions assessing affection toward partner) where higher scores indicate higher levels of affection.

Table S4: Associations between attachment and (a) depression (b) self-harm at age 18. Results are based on complete case analysis (n=2397).

	Depression		Self-Harm	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Model 1	1.07 (0.88, 1.26)	0.49	1.02 (0.86,1.18)	0.82
Model 2	1.11 (0.92, 1.31)	0.26	1.08 (0.91, 1.25)	0.37
Model 3	1.13 (0.93, 1.32)	0.24	1.08 (0.91, 1.24)	0.41
Model 4	1.13 (0.94, 1.33)	0.21	1.05 (0.88, 1.23)	0.54
Model 5	1.14 (0.94, 1.34)	0.21	1.05 (0.88, 1.22)	0.58

Note: **Model 1** = unadjusted; **Model 2** = adjusted for sex; **Model 3** = further adjusted for parental social class, housing tenure, maternal education; **Model 4** = further adjusted for maternal age, maternal depression, maternal smoking and alcohol consumption in pregnancy, number of children <15 years-of-age in household; **Model 5** = further adjusted for parental affection score and whether parents separated/divorced.