

Supplementary Material

Linear Mixed Effect Models

Model

Below is model used to fit the data to each diffusion metric (FA, MD, NDI, ODI and TF) in each ROI (n = 15). Random effects were separated by group, with random intercepts and slopes allowed to be correlated (i.e. unstructured covariance structure) for the majority of models:

$$y_{ijk} = (\beta_{0k} + \beta_{1k}b_{ij}) + (\beta_{2k} + \beta_{3k}d_i^2 + \beta_{4k}d_i) + \epsilon_{ijk}$$

$$\begin{pmatrix} \beta_{0k} \\ \beta_{1k} \end{pmatrix} \sim N \left(0, \begin{pmatrix} \sigma_{\beta_0}^2 & \sigma_{\beta_1}^2 \\ \sigma_{\beta_0\beta_1} & \sigma_{\beta_1}^2 \end{pmatrix} \right), \epsilon_{ijk} \sim N(0, \sigma_{\epsilon}^2)$$

Where, for the kth (k=1,2,3,4) measurement for the jth participant (j = 1...n_{ij}) in the ith group (i=1 YOAD, i=2 control):

y_{ijk} = dMRI metric

β_{0k} = Fixed distance (mm) sampled from the GM/WM boundary. Moving from GM towards the SWM/DWM:

- GM (+1mm) was set as $d_1=0$
- GM/WM boundary (0mm) as $d_2=1$
- SWM (-1mm) as $d_3=2$
- SWM/DWM (-2mm) as $d_4=3$

ϵ_{ijk} = cortical thickness (mm) as a constant over k

β_{0k} = Fixed intercepts for the two groups

β_{1k} = Fixed linear effects for the two groups

β_{2k} = Fixed quadratic effects for the two groups

β_{3k} = Fixed cortical thickness effect

β_{4k} = Random intercept for the jth participant in the ith group

b_{ij} = Random slope for the jth participant in the ith group

ϵ_{ijk} = Residual

In regions and dMRI metrics where the model was too complex (6/75 models), we fit the same model but removed the control group's random slope.

Example R code for fitting models

The above model was implemented in R using the `lme4` library. Separate models were run for each combination of dMRI metric and ROI. Group-specific random effects were achieved by creating separate intercept and distance vectors (e.g. (0 + control + distance_control | Participant)). Below shows example code for modelling NDI measures across the GM/WM boundary in the parahippocampal cortex (phippo_ndi) where random intercepts and slopes are included for both control and YOAD groups.

```
# Set tolerance for convergence to resolve convergence problems at cost of slower fit (see lme4 '?convergence' help file)
strict_tol <- lmerControl(optCtrl=list(xtol_abc=1e-8, ftol_abc=1e-8))

# Run model
phippo_ndi_model <- lmer(DWI ~ distance + I(distance^2)* group + distance*group + I(distance^2)*group + cortical_thick +
  (0 + control + distance_control | Participant) +
  (0 + AD + distance_AD | Participant),
  data=phippo_ndi,
  REML=TRUE,
  control = strict_tol)

# View model results
summary(phippo_ndi_model)
```

Extracting Average Marginal Effects (AMEs)

Marginal effects were estimated from the LME models to assess differences between YOAD and control groups at each sampled distance along the cortical profile. This was done using the `margins` library which emulates the average marginal effects command used in STATA.


```
# Extract marginal effects at each distance (0:3 = 1mm to -2mm) across groups (0 = control, 1 = YOAD)
phippo_ndi_marg <- as.data.frame(summary(margins(phippo_ndi_model, at = list(distance = 0:3, group = 0:1))))

# View marginal effects between groups
phippo_ndi_marg %>% filter(factor == "group")
```

The output provides the average marginal effect (AME) between groups at each distance, standard error, z value and p value. All AME p-values were then adjusted using false discovery rate (FDR) using the `p.adjust` package, where $pFDR < 0.05$ indicates statistical significance at our chosen threshold of 0.05 after correcting for multiple comparisons.


Average Marginal Effects Plots

Supplementary Figure 1: DTI AMEs for all ROIs Along the Cortical Profile




Supplementary Figure 1: Plots showing DTI AMEs of each ROI across the GM/WM boundary from GM to SWM/DWM. AMEs are shown in full colour when $pFDR < 0.05$ and transparent when $pFDR > 0.05$.

Supplementary Figure 2: NODDI AMEs for all ROIs Along the Cortical Profile



Supplementary Figure 2: Plots showing NODDI AMEs of each ROI across the GM/WM boundary from GM to SWM/DWM. AMEs are shown in full colour when $pFDR < 0.05$ and transparent when $pFDR > 0.05$.

Supplementary Figure 3: TF AMEs for all ROIs Along the Cortical Profile



Supplementary Figure 3: Plots showing TF AMEs of each ROI across the GM/WM boundary from GM to SWM/DWM. AMEs are shown in full colour when $pFDR < 0.05$ and transparent when $pFDR > 0.05$.

Tables

Supplementary Table 1: DTI AMEs for all ROIs and Distances

ROI	Measure	GM	GM/WM	SWM	SWM/DWM
Cuneus	FA	-0.071 (0.025, 0.025)	-0.011 (0.025, 0.025)	-0.004 (0.025, 0.025)	0.006 (0.025, 0.025)
	MD	0.247 (0.108, 0.330)	0.103 (0.025, 0.142)	0.013 (0.027, 0.027)	-0.254 (0.027, 0.027)
Entorhinal	FA	-0.023 (0.025, 0.025)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	-0.022 (0.025, 0.025)
	MD	0.148 (0.108, 0.223)	0.068 (0.025, 0.148)	0.014 (0.025, 0.025)	0.027 (0.025, 0.025)
Fusiform	FA	-0.004 (0.025, 0.025)	-0.003 (0.025, 0.025)	-0.003 (0.025, 0.025)	0.004 (0.025, 0.025)
	MD	0.178 (0.124, 0.223)	0.044 (0.025, 0.066)	-0.011 (0.025, 0.025)	-0.018 (0.025, 0.025)
Inferioparietal	FA	-0.022 (0.027, 0.027)	-0.013 (0.025, 0.025)	-0.009 (0.025, 0.025)	-0.007 (0.027, 0.027)
	MD	0.341 (0.247, 0.414)	0.123 (0.066, 0.178)	0.002 (0.044, 0.044)	-0.004 (0.066, 0.066)
Inferiotemporal	FA	-0.006 (0.025, 0.025)	-0.002 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.004 (0.025, 0.025)
	MD	0.156 (0.114, 0.190)	0.047 (0.024, 0.078)	-0.005 (0.027, 0.027)	-0.005 (0.024, 0.024)
Laterooccipital	FA	-0.016 (0.026, 0.026)	-0.012 (0.025, 0.025)	-0.005 (0.025, 0.025)	0.001 (0.025, 0.025)
	MD	0.302 (0.234, 0.373)	0.144 (0.046, 0.198)	0.006 (0.024, 0.024)	-0.011 (0.026, 0.026)
Middletemporal	FA	-0.011 (0.027, 0.027)	-0.005 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.004 (0.025, 0.025)
	MD	0.214 (0.161, 0.266)	0.097 (0.026, 0.150)	-0.014 (0.042, 0.042)	-0.001 (0.026, 0.026)
Paracentral	FA	-0.016 (0.026, 0.026)	-0.014 (0.024, 0.024)	-0.009 (0.025, 0.025)	0.001 (0.025, 0.025)
	MD	0.144 (0.063, 0.204)	0.066 (0.017, 0.090)	-0.001 (0.027, 0.027)	-0.001 (0.027, 0.027)
Parahippocampal	FA	-0.013 (0.027, 0.027)	-0.024 (0.026, 0.026)	-0.005 (0.025, 0.025)	-0.001 (0.027, 0.027)
	MD	0.177 (0.108, 0.227)	0.062 (0.047, 0.117)	0.008 (0.026, 0.026)	0.001 (0.026, 0.026)
Postcentral	FA	-0.012 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.008 (0.026, 0.026)
	MD	0.208 (0.144, 0.276)	0.074 (0.028, 0.117)	0.007 (0.025, 0.025)	-0.004 (0.026, 0.026)
Posterocingulate	FA	-0.009 (0.026, 0.026)	-0.008 (0.025, 0.025)	-0.007 (0.025, 0.025)	0.004 (0.027, 0.027)
	MD	0.118 (0.018, 0.198)	0.064 (0.026, 0.082)	0.003 (0.025, 0.025)	0.004 (0.027, 0.027)
Precentral	FA	-0.015 (0.026, 0.026)	-0.003 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.001 (0.026, 0.026)
	MD	0.148 (0.071, 0.221)	0.062 (0.025, 0.086)	-0.011 (0.024, 0.024)	-0.001 (0.026, 0.026)
Precuneus	FA	-0.016 (0.026, 0.026)	-0.002 (0.025, 0.025)	-0.002 (0.025, 0.025)	0.001 (0.026, 0.026)
	MD	0.269 (0.207, 0.322)	0.081 (0.045, 0.130)	-0.010 (0.042, 0.042)	-0.018 (0.026, 0.026)
Superiorparietal	FA	-0.009 (0.026, 0.026)	-0.018 (0.025, 0.025)	-0.002 (0.025, 0.025)	0.000 (0.026, 0.026)
	MD	0.306 (0.276, 0.414)	0.154 (0.066, 0.218)	0.007 (0.024, 0.024)	0.001 (0.026, 0.026)
Superiortemporal	FA	-0.008 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.004 (0.026, 0.026)	0.001 (0.026, 0.026)
	MD	0.143 (0.101, 0.188)	0.028 (0.014, 0.056)	-0.001 (0.025, 0.025)	-0.008 (0.026, 0.026)

Notes: Values reflect Average Marginal Effects (AME) and associated 95% confidence intervals. Colours indicate FDR adjusted p-value thresholds. Red = $pFDR < 0.001$, orange = $pFDR < 0.01$, blue = $pFDR < 0.05$.

Supplementary Table 2: NODDI AMEs for all ROIs and Distances

ROI	Measure	GM	GM/WM	SWM	SWM/DWM
Cuneus	NDI	-0.010 (0.025, 0.025)	-0.007 (0.025, 0.025)	-0.005 (0.025, 0.025)	-0.004 (0.025, 0.025)
	ODI	-0.020 (0.046, 0.074)	-0.003 (0.025, 0.025)	0.003 (0.025, 0.025)	0.003 (0.025, 0.025)
Entorhinal	NDI	-0.004 (0.025, 0.025)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)
	ODI	0.008 (0.016, 0.016)	0.031 (0.016, 0.042)	0.010 (0.016, 0.016)	0.012 (0.016, 0.016)
Fusiform	NDI	-0.002 (0.025, 0.025)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.001 (0.025, 0.025)
	ODI	-0.007 (0.016, 0.016)	0.007 (0.016, 0.016)	0.007 (0.016, 0.016)	0.007 (0.016, 0.016)
Inferioparietal	NDI	-0.001 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.009 (0.046, 0.024)	-0.003 (0.025, 0.025)	-0.003 (0.024, 0.024)	-0.003 (0.046, 0.024)
Inferiotemporal	NDI	-0.011 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.002 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.014 (0.026, 0.026)	-0.003 (0.025, 0.025)	-0.009 (0.025, 0.025)	0.004 (0.026, 0.026)
Laterooccipital	NDI	-0.012 (0.026, 0.026)	-0.011 (0.025, 0.025)	-0.010 (0.025, 0.025)	-0.008 (0.026, 0.026)
	ODI	-0.004 (0.041, 0.041)	-0.003 (0.018, 0.018)	-0.006 (0.016, 0.016)	-0.009 (0.026, 0.026)
Middletemporal	NDI	-0.012 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.003 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.014 (0.026, 0.026)	-0.001 (0.025, 0.025)	0.001 (0.018, 0.018)	0.001 (0.016, 0.016)
Paracentral	NDI	-0.012 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.016 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.009 (0.025, 0.025)	0.012 (0.026, 0.026)
Parahippocampal	NDI	-0.011 (0.026, 0.026)	-0.001 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.012 (0.026, 0.026)
	ODI	0.001 (0.016, 0.016)	0.001 (0.016, 0.016)	0.001 (0.016, 0.016)	0.001 (0.016, 0.016)
Postcentral	NDI	-0.007 (0.026, 0.026)	-0.003 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.004 (0.026, 0.026)
	ODI	-0.028 (0.046, 0.024)	-0.011 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.002 (0.026, 0.026)
Posterocingulate	NDI	-0.006 (0.026, 0.026)	-0.008 (0.025, 0.025)	-0.008 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.006 (0.046, 0.024)	-0.008 (0.024, 0.024)	0.001 (0.016, 0.016)	0.001 (0.016, 0.016)
Precentral	NDI	-0.004 (0.026, 0.026)	-0.002 (0.025, 0.025)	-0.002 (0.025, 0.025)	0.001 (0.026, 0.026)
	ODI	-0.019 (0.016, 0.016)	0.002 (0.016, 0.016)	0.001 (0.016, 0.016)	0.001 (0.016, 0.016)
Precuneus	NDI	-0.015 (0.026, 0.026)	-0.009 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.002 (0.041, 0.041)
	ODI	-0.029 (0.046, 0.024)	-0.011 (0.025, 0.025)	-0.003 (0.025, 0.025)	-0.001 (0.026, 0.026)
Superiorparietal	NDI	-0.003 (0.046, 0.024)	-0.002 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.003 (0.026, 0.026)
	ODI	-0.005 (0.046, 0.024)	-0.002 (0.025, 0.025)	-0.001 (0.025, 0.025)	0.007 (0.026, 0.026)
Superiortemporal	NDI	0.003 (0.026, 0.026)	0.003 (0.025, 0.025)	0.003 (0.025, 0.025)	0.003 (0.026, 0.026)
	ODI	-0.006 (0.046, 0.024)	0.004 (0.024, 0.024)	0.008 (0.016, 0.016)	0.010 (0.026, 0.026)

Notes: Values reflect Average Marginal Effects (AME) and associated 95% confidence intervals. Colours indicate FDR adjusted p-value thresholds. Red = $pFDR < 0.001$, orange = $pFDR < 0.01$, blue = $pFDR < 0.05$.

Supplementary Table 3: Tissue Fraction AMEs for all ROIs and Distances

ROI	Measure	GM	GM/WM	SWM	SWM/DWM
Cuneus	TF	-0.148 (0.181, -0.112)	-0.086 (0.108, -0.062)	-0.042 (0.024, 0.026)	-0.041 (0.022, -0.024)
	MD	-0.104 (0.194, -0.066)	-0.044 (0.074, -0.016)	-0.001 (0.028, 0.016)	-0.004 (0.022, 0.024)
Entorhinal	TF	-0.117 (0.172, -0.108)	-0.065 (0.088, -0.048)	-0.021 (0.045, 0.022)	-0.009 (0.028, 0.028)
	MD	-0.083 (0.148, -0.011)	-0.031 (0.027, -0.027)	-0.001 (0.025, 0.025)	-0.009 (0.028, 0.028)
Fusiform	TF	-0.207 (0.245, -0.165)	-0.069 (0.127, -0.071)	-0.021 (0.027, -0.015)	-0.009 (0.024, 0.024)
	MD	-0.117 (0.143, -0.091)	-0.022 (0.027, -0.022)	-0.015 (0.025, 0.025)	-0.001 (0.024, 0.024)
Inferioparietal	TF	-0.187 (0.165, -0.198)	-0.060 (0.086, -0.082)	-0.014 (0.028, 0.028)	0.000 (0.012, 0.012)
	MD	-0.088 (0.118, -0.058)	-0.041 (0.061, -0.021)	-0.012 (0.023, 0.023)	-0.014 (0.023, 0.023)
Laterooccipital	TF	-0.086 (0.198, -0.122)	-0.041 (0.061, -0.041)	-0.012 (0.023, 0.023)	0.000 (0.012, 0.012)
	MD	-0.119 (0.143, -0.091)	-0.041 (0.061, -0.041)	-0.021 (0.023, 0.023)	-0.021 (0.023, 0.023)
Middletemporal	TF	-0.102 (0.128, -0.076)	-0.041 (0.058, -0.041)	-0.021 (0.023, 0.023)	-0.004 (0.016, 0.016)
	MD	-0.094 (0.128, -0.066)	-0.041 (0.057, -0.022)	-0.013	