

Supplemental material to:

## **Pneumococcal exposure routes for infants, a nested cross-sectional survey in Nha Trang, Vietnam**

Authors: George Qian<sup>1\*</sup>, Michiko Toizumi<sup>2\*</sup>, Sam Clifford<sup>1</sup>, Lien Thuy Le<sup>3</sup>, Tasos Papastylia<sup>4</sup>, Billy Quilty<sup>1</sup>, Chihiro Iwasaki<sup>2</sup>, Noriko Kitamura<sup>2</sup>, Mizuki Takegata<sup>2</sup>, Trang Minh Nguyen<sup>5</sup>, Hien Anh Thi Nguyen<sup>6</sup>, Duc Anh Dang<sup>6</sup>, Albert Jan van Hoek<sup>7</sup>, Lay Myint Yoshida<sup>2†</sup>, Stefan Flasche<sup>1†</sup>

Institutions:

<sup>1</sup> Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene and Tropical Medicine, London, UK

<sup>2</sup> Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan

<sup>3</sup> Department of Bacteriology, the Pasteur Institute in Nha Trang, Nha Trang, Vietnam

<sup>4</sup> School of Computer Science and Electronic Engineering, University of Essex, Colchester, UK

<sup>5</sup> Khanh Hoa Health Service, Nha Trang, Vietnam

<sup>6</sup> National Institute of Hygiene and Epidemiology, Hanoi, Vietnam

<sup>7</sup> Centre for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, The Netherlands

### *Logistic Regression Equation*

The equation for the logistic regression model predicting carriage in infants (see the ‘Statistical Analyses’ part of the Methods section in the main text) is:

$$\text{logit}(p_i) = b_0 + C_i + b_1 E_i + b_2 A_i \quad (1)$$

where:

- $p_i$  is the probability that the  $i$ th infant in the study is a pneumococcal carrier
- $b_0$  is the model intercept under a single-level regression model (fixed effects only)
- $C_i$  is the random effect on the intercept due to the influence of the commune to which the  $i$ th individual belongs
- $b_1$  is the coefficient corresponding to the effect of PEI
- $E_i$  is the value of PEI, for the  $i$ th individual
- $b_2$  is the coefficient corresponding to the effect of the infant's age

- $A_i$  is the age of the  $i$ th individual in months

### Model Comparison

We compared the model in Equation 2 with other logistic regression models with different covariants:

- Carriage ~ Number of Contacts (traditionally the covariate used for predicting carriage)
- Carriage ~ PEI (to justify the inclusion of the other covariates)
- Carriage ~ PEI + Infants' Age
- Carriage ~ PEI + Commune
- Carriage ~ [(PEI + Infants' Age) with commune as groups] | longer duration contacts only (including only long-duration contacts)

These models were compared using the Deviance Information Criterion (DIC) values, a generalisation of the Akaike Information Criterion that is suitable for comparison of Bayesian models, provided that the covariates approximately follow a Gaussian distribution; we found that this was indeed the case for the covariates in our model.

The results are shown in Table 5. The model with PEI as the lone covariate appears to outperform the corresponding model with the number of contacts alone, based on DIC values. The set of models with PEI together with covariates such as mobility and motorcycle share similar DIC values and appear to be the best-performing models; this set includes the proposed model (Equation 2). Including only longer-duration contacts does not appear to reduce the DIC value.

**Table S3:** Effect of each infant's characteristic on having contact outside of the commune of residence, estimated using logistic regression model.

Characteristics	N	% Infants with contacts in other communes	Odds ratio	Adjusted odds ratio*	Adjusted odds ratio**
Total	1583	154 (9.7)			
Demographics					
Sex					
Male	871	85 (9.8)	reference	reference	
Female	712	69 (9.7)	0.99 (0.71-1.39)	0.96 (0.66-1.41)	
Age (months)					
<8 months	616	44 (7.1)	reference		reference
8-13 months	967	110 (11.4)	1.67 (1.16-2.40)		1.41 (1.05-1.90)***
Family					
Siblings in the household					
No sibling	666	66 (9.9)	reference	reference	
One or more	917	88 (9.6)	0.97 (0.69-1.35)	0.96 (0.74-1.23)	

siblings						
Number of people in the household						
2-4	611	62 (10.2)	reference	reference		
>4	972	92 (9.5)	0.93 (0.66-1.3)	0.91 (0.62-1.36)		
Caretaker currently in a paid employment						
No	1056	86 (8.1)	reference	reference	reference	
Yes	527	68 (12.9)	1.67 (1.19-2.34)	1.60 (1.13-2.27)	1.08 (0.71-1.65)	
Highest level of education in the household						
Secondary school or lower	268	27 (10.1)	reference	reference		
Degree	1315	127 (9.7)	0.95 (0.62-1.48)	0.94 (0.65-1.36)		
Infant's activity						
Sit						
Yes	1334	139 (10.4)	1.81 (1.05-3.15)	1.36 (0.70-2.62)		
No	249	15 (6.0)	reference	reference		
Crawl						
Yes	873	95 (10.9)	1.35 (0.96-1.9)	1.08 (0.73-1.59)		
No	710	59 (8.3)	reference	reference		
Walk						
Yes	186	28 (9.0)	1.79 (1.15-2.78)	1.51 (0.95-2.39)		
No	1397	126 (15.1)	reference	reference		
Mobility						
Bicycle						
Yes	31	1 (3.2)	0.30 (0.04-2.25)	0.30 (0.04-2.32)		
No	1552	153 (9.9)	reference	reference		
Motorbike						
Yes	1549	152 (9.8)	1.74 (0.41-7.34)	1.73 (0.30-9.9)		
No	34	2 (5.9)	reference	reference		
Car						
Yes	99	12 (12.1)	1.3 (0.7-2.44)	1.34 (0.78-2.31)		
No	1484	142 (9.6)	reference	reference		
Walk						
Yes	108	10 (9.3)	0.94 (0.48-1.85)	0.97 (0.50-1.87)		
No	1475	144 (9.8)	reference	reference		
Public transportation						
Yes	40	5 (12.5)	1.34 (0.52-3.46)	1.44 (0.68-3.03)		
No	1543	149 (9.7)	reference	reference		
Number of times caretaker left the commune in the last 7 days						
0-2	872	44 (5.1)	reference	reference	reference	
3 or more	711	110 (15.5)	3.44 (2.39-4.96)	3.36 (2.02-5.58)	3.01 (1.78-5.10)	
Number of times infant left the commune in the last 7 days						
0	788	9 (1.1)	reference	reference	reference	
1 or more	795	145 (18.2)	19.31 (9.77-38.16)	18.9 (7.99-44.72)	18.3 (7.87-42.47)	

Day-care					
Day-care attendance					
Yes	1497	121 (8.1)	7.08 (4.41-11.36)	6.56 (3.92-10.98)	
No	86	33 (38.4)	reference	reference	
Pneumococcus carriage (n=1582)					
Pneumococcus					
Yes	353	48 (13.6)	1.67 (1.16-2.40)	1.58 (1.13-2.21)	1.25 (0.92-1.71)
No	1229	106 (8.6)	reference	reference	reference

\*Odds ratios adjusted by age group, considering clustering in each commune

\*\*Odds ratios adjusted by age group & day-care attendance, considering clustering in each commune

\*\*\*Odds ratio adjusted by day-care attendance, considering clustering in each commune

**Table S4:** A comparison of different logistic regression models to predict carriage, using DIC values

Covariates	DIC value
Number of Contacts	1682
PEI	1656
PEI + Infants' Age	1618
PEI+Infants' Age, with Commune as a random effect	1618
PEI + Infants' Age - Longer Duration Contacts Only, with Commune as a random effect	1619

**Table S5:** Values of the coefficients from the logistic regression (Equation 2) are given here.

Parameter	Mean Value	2.5%	97.5%
Fixed-effects model Intercept ( $\beta_0$ )	-3.33	-4.82	-2.05
Effect of PEI ( $\beta_1$ )	1.96	1.40	2.51
Commune ID: 21 ( $C_1$ )	0.13	-0.39	0.64
Commune ID: 17 ( $C_2$ )	0.21	-0.32	0.71
Commune ID: 16 ( $C_3$ )	0.06	-0.46	0.57


Commune ID: 23 ( $C_4$ )	0.01	-0.52	0.53
Commune ID: 18 ( $C_5$ )	-0.11	-0.68	0.43
Commune ID: 14 ( $C_6$ )	0.18	-0.32	0.70
Commune ID: 27 ( $C_7$ )	-0.12	-0.66	0.41
Commune ID: 10 ( $C_8$ )	-0.16	-0.71	0.37
Commune ID: 7 ( $C_9$ )	0.05	-0.50	0.58
Commune ID: 3 ( $C_{10}$ )	-0.43	-1.09	0.14
Commune ID: 25 ( $C_{11}$ )	-0.40	-1.03	0.17
Commune ID: 20 ( $C_{12}$ )	0.39	-0.13	0.90
Commune ID: 13 ( $C_{13}$ )	0.11	-0.43	0.61
Commune ID: 11 ( $C_{14}$ )	-0.02	-0.58	0.52
Commune ID: 5 ( $C_{15}$ )	-0.16	-0.73	0.35
Commune ID: 15 ( $C_{16}$ )	0.55	0.05	1.05
Commune ID: 24 ( $C_{17}$ )	0.32	-0.20	0.84
Commune ID: 12 ( $C_{18}$ )	-0.66	-1.34	-0.07
Commune ID: 1 ( $C_{19}$ )	-0.15	-0.71	0.38
Commune ID: 4 ( $C_{20}$ )	0.01	-0.51	0.50
Commune ID: 19 ( $C_{21}$ )	-0.37	-0.96	0.17
Commune: ID 22 ( $C_{22}$ )	0.09	-0.45	0.60
Commune ID: 2 ( $C_{23}$ )	-0.17	-0.73	0.36
Commune ID: 9 ( $C_{24}$ )	0.21	-0.33	0.74
Commune ID: 8 ( $C_{25}$ )	0.04	-0.50	0.55
Commune ID: 6 ( $C_{26}$ )	-0.17	-0.74	0.36

Commune ID: 26( $C_{27}$ )	-0.12	-0.72	0.45
Infants' Age ( $b_2$ )	0.04	0.01	0.08

**Table S6:** The VIF values quantifying multicollinearity among the covariates of the model are shown here. The values are well below 10 (which, as a 'rule of thumb', is when multicollinearity becomes influential).

Covariate	VIF
Mean PEI value	1.05
Infant Age	1.03
Commune	1.08

**Figure S4:** The Directed Acyclic Graph of the factors involved in the spread of pneumococcus to infants in Nha Trang is shown here



**Figure S5:** Distribution of 10000 MCMC samples of PEI for a randomly selected individual (Individual A), shown through a violin plot and boxplot nested within.

