Evolution of SARS-CoV-2 seroprevalence and clusters in school children from June 2020 to April 2021 reflect community transmission: prospective cohort study *Ciao Corona* Agne Ulyte\*, Thomas Radtke\*, Irene A. Abela, Sarah R. Haile, Priska Ammann, Christoph Berger, Alexandra Trkola, Jan Fehr, Milo A. Puhan#, Susi Kriemler#

Appendix 4 Detailed information of classes with potential clusters (3 or more newly seropositive children at T3) in schools with 2 or more classes with clusters

| School | School level | N of pupils:<br>T1+/T2+/T3+<br>negative<br>missing* | RT-PCR positive in T2-T3 |       | Quarantine/<br>isolation in T2-T3 |                | Information** (unrelated time points signify >2 weeks interval between all diagnosed or potential   | Possible i<br>individua |       |                | Transmission in class plausible | Comments  |
|--------|--------------|---|--------------------------|-------|-----------------------------------|----------------|---|-------------------------|-------|----------------|---------------------------------|---|
|        |              |   | Teacher                  | Child | Teacher                           | Child          | infections)   | Teacher                 | Child | House-<br>hold | ****                            |   |
| 1      | lower        | 1/0/4<br>10<br>3                                    | 0                        | 0     | 0                                 | 3              | 3 pupils in quarantine: n=1 due to PCR+<br>teacher not related to regular school, n=2<br>both due to PCR+ household member (HM)<br>of one of the 2 pupils, with whom both had<br>close contact  | 0                       | 0     | 1              | 1                               | Outbreak in grade 5 class<br>non-participating in this<br>study: 8 pupils PCR+, not<br>clear where Index came from,<br>family of 1 pupil and HM of<br>another pupil were tested<br>PCR+ |
|        | lower        | 1/0/3<br>5<br>11                                    | 1                        | 0     | 1                                 | 7              | 7 pupils in quarantine: n=5 due to PCR+ HM and n=2 for unknown reasons  | 0                       | 0     | 1              | 1                               |   |
|        | lower        | 0/0/4<br>8<br>8                                     | 0                        | 0     | 0                                 | 4              | 4 pupils in quarantine: n=1 due to PCR+ HM,<br>n=1 due to PCR+ teacher not related to<br>regular school, n=1 due to PCR+ family<br>friend, n=1 for unknown reasons  | 0                       | 0     | 1              | 0                               |   |
|        | lower        | 1/0/4<br>10<br>3                                    | 0                        | 1     | 1                                 | 4              | 1 pupil in isolation: PCR+ for unknown reasons. 3 pupils in quarantine: n=2 each due to PCR+ HM (unrelated time points), n=1 for unknown reason   | 0                       | 0     | 1              | 0                               |   |
| 2      | lower        | 0/0/4<br>8<br>11                                    | 0                        | 2     | 0                                 | 3              | 2 pupils in isolation: n=1 first in quarantine due to PCR+ HM then PCR+, n=1 PCR+ due to symptoms. 2 pupils in quarantine at unrelated time points each due to PCR+ HM  | 0                       | 0     | 1              | 1                               |   |
|        | middle       | 0/0/4<br>10<br>6                                    | 0                        | 0     | 0                                 | 1              | 1 pupil in quarantine due to PCR+ HM  | 0                       | 0     | 1              | ý                               |   |
|        | middle       | 0/0/3<br>3<br>16                                    | 0                        | 0     | 0                                 | 2              | 2 pupils in quarantine at unrelated time points each due to PCR+ HM   | 0                       | 0     | 1              | 0                               |   |
| 3      | lower        | 0/0/3<br>7<br>9                                     | 1                        | 1     | 1                                 | whole<br>class | Outbreak in a parallel class due to PCR+ child (probably infected by HM; index HM with severe symptoms but not tested). Then teacher and several children PCR+ in this and the parallel classes, the remaining children not tested; both classes (including this one) quarantined | 0                       | 0     | 1              | 1                               | Transmission of virus among pupils in the lower school level class probable, primary Index case most probably HM of a child.  |

|   | middle | 1/1/3<br>12<br>5 | 0 | 0 | 0               | 1               | 1 pupil in quarantine due to travel abroad after fall vacation   | 0 | 0 | 0 | ? |  |
|---|--------|------------------|---|---|-----------------|-----------------|--|---|---|---|---|--|
|   | middle | 1/1/3<br>7<br>8  | 1 | 3 | 1               | 3               | 1 teacher in isolation PCR+, 3 pupils in isolation, each PCR+ due to PCR+ HM (unrelated time points)   | ? | 0 | 1 | 1 |  |
| 4 | lower  | 1/0/3<br>2<br>15 | 1 | 3 | whole<br>school | whole<br>school | 3 pupils in isolation: n=1 PCR+ for unknown reason; n=1 due to PCR+ HM; n=1 PCR+ with symptoms   | 0 | ? | 1 | 1 | School outbreak with the whole school in quarantine                              |
|   | middle | 1/1/3<br>7<br>8  | 1 | 7 | _               |                 | 1 teacher PCR+, 3-6 days later PCR+ pupil<br>tested because of loss of taste. Testing of the<br>whole class 2 days later - 6 further pupils<br>PCR+, none with symptoms, 5 pupils PCR- | 1 | 0 | 0 | 1 |  |
|   | middle | 0/1/3<br>7<br>12 | 1 | 6 |                 |                 | 6 pupils in isolation: n=1 in quarantine due to PCR+ HM, then tested PCR+ in the outbreak testing together with 1 teacher and 5 further pupils PCR+                                    | ? | ? | 1 | 1 |  |
| 5 | lower  | 1/1/3<br>3<br>13 | 0 | 0 | 0               | 0               | No diagnosed or suspected SARS-CoV-2 infections known  | ? | ? | ? | ? |  |
|   | middle | 0/0/4<br>6<br>11 | 0 | 0 | 0               | 0               | No diagnosed or suspected SARS-CoV-2 infections known  | ? | ? | ? | ? |  |
| 6 | middle | 0/0/3<br>13<br>8 | 0 | 8 | 1               | whole<br>class  | First a few children PCR+, then in total 8 PCR+ shortly afterwards. Subsequently class in quarantine, teacher then tested PCR– but later tested seropositive                           | 0 | 1 | ? | 1 |  |
|   | middle | 1/1/3<br>3<br>15 | 0 | 3 | 0               | 3               | 3 pupils in quarantine at unrelated time points each due to PCR+ HM  | 0 | 0 | 1 | 0 |  |
| 7 | upper  | 0/1/3<br>10<br>3 | 0 | 2 | 0               | 2               | 2 pupils in isolation at unrelated time points for unknown reasons   | 0 | ? | ? | 0 | A class not participating in this study in quarantine during Christmas break due |
|   | upper  | 0/0/3<br>7<br>5  | 0 | 1 | 0               | 1               | 1 pupil in isolation for unknown reasons   | 0 | ? | ? | ? | to several PCR+ pupils   |
| 8 | lower  | 0/0/3<br>4<br>15 | 0 | 1 | 0               | 1               | 1 pupil in isolation: infected by a teacher not related to the regular school, whole class tested PCR-   | 0 | 0 | 0 | ? |  |
|   | middle | 1/0/3<br>12<br>8 | 0 | 1 | 0               | 4               | 1 pupil in isolation: infected by a teacher not related to the regular school. 3 pupils in quarantine at unrelated time points: n=1 due to PCR+ HM, n=2 of for unknown reasons         | 0 | 0 | 1 | 0 |  |
| 9 | middle | 2/0/3<br>8<br>11 | 0 | 1 | 0               | 6               | 1 pupil in isolation: PCR+ during quarantine due to a PCR+ HM. 6 pupils in quarantine all due to PCR+ HM   | 0 | 0 | 1 | 1 | Parallel class not participating in this study in quarantine                     |

|    | middle | 0/1/4<br>1<br>18 | 0 | 0 | 0 | 2 | 2 pupils in quarantine at unrelated time points due to PCR+ HM   | 0 | 0 | 1 | 0 | due to infected teacher and class assistant |
|----|--------|------------------|---|---|---|---|--|---|---|---|---|---|
| 10 | lower  | 1/0/5<br>9<br>9  | 0 | 0 | 0 | 5 | 1 pupil in isolation due to PCR+ HM.<br>4 pupils in quarantine: n=3 each due to PCR+<br>HM, n=1 due to PCR+ contact  | 0 | 0 | 1 | 1 |   |
|    | lower  | 0/0/3<br>12<br>8 | 0 | 0 | 0 | 3 | 3 pupils in quarantine: n=1 due to PCR+<br>teacher not related to the regular school,<br>n=2 both due to PCR+ HM of one of the 2<br>pupils, with whom both had close contact | 0 | 0 | 1 | 1 |   |

HM – household member.

<sup>\*</sup> T1+ denotes the number of seropositive pupils at T1 (June-July 2020); T2+ denotes the number of newly seropositive pupils at round T2 (October-November 2020); T3+ denotes the number of newly seropositive pupils at T3 (March-April 2021); negative denotes the number of seronegative pupils at round T3; missing denotes previously non-seropositive pupils who were not tested at T3. Total class size can be calculated by the sum of T1+, T2+, T3+, negative, and missing.

<sup>\*\*</sup> Dates reported denote month and day(s). September to December refer to the year 2020, January to April – to the year 2021.

<sup>\*\*\*</sup> Possible index cases of individual diagnosed or suspected (RT-PCR+ or quarantined) children in the class.

<sup>\*\*\*\*</sup> Transmission among pupils/teachers in class was considered possible (1) when the time between RT-PCR diagnoses or quarantine start dates was 2 weeks or below, and not probable (0) if longer. Transmission was defined as unclear (?) if no or only a single child was diagnosed or quarantined in the class according to the interview information.