




Figure S1. HVTN 100 clinical trial and immunogenicity data at month 12.5. A) Trial schema for the HVTN 100 clinical trial indicating the vaccination schedule and Env-specific B cell phenotyping and BCR repertoire analysis timepoints. Immune responses at month 12.5 to vaccine-matched antigens for the B cell analysis subset (n=14) in blue overlaid on the entire vaccinated cohort in yellow (n=185): Binding antibody responses to vaccine-matched B) gp120 and C) V1V2 proteins. D) Antibody-dependent cellular cytotoxicity (ADCC) activity of serum as measured by luciferase-based cytotoxicity assay. E) CD4 T cell responses (IL-2, IFN- γ , or CD40L) to vaccine-matched gp120 peptide pools. Colored circles indicate statistically “positive” response whereas open triangles indicate non-response.





Figure S2. Diversity of V_H , V_K and V_L s for the gp120⁻ and gp120⁺ repertoires as calculated by the A) Shannon or B) Simpson indices.

A

Heavy Chain

Kappa Chain

Lambda Chain



B

Heavy Chain

Kappa Chain

Lambda Chain





Figure S3. Proportions of V_H , V_K and V_L sequences encoded by specific V genes for the gp120- and gp120+ repertoires, by subject (one subject per row): A) subjects 1-7 and B) 8-14. The total number of sequences per repertoire is indicated in the center of the pie chart. Each V gene family (e.g., VH1, VH2...) is highlighted by a specific color that is identical across chains (e.g., all genes in the VH1, VL1 and VK1 families are shown in red); different V gene segments within a V gene family (e.g., VH1-2) are represented by different shades of the same color.

A Kappa Chain



B Lamda Chain




Figure S4. Rate of mutation among gp120+ and gp120- kappa and lambda sequences. Mean amino acid (a.a.) mutation rates per site in the respective framework (FR) or complementary determining (CDR) regions of the A) V_K and B) V_L sequences split out by V gene for the gp120+ or gp120- repertoires, followed by the difference in a.a. mutation rate between the gp120+ and gp120- sequences per V_K or V_L region. In the heatmap that shows the difference between mutation rates in the gp120+ and gp120- repertoires, color of heatmap indicates higher (red) or lower (purple) mutation rates in the gp120+ than in the gp120- repertoire.