

ONLINE SUPPLEMENT II: eTable3, eTable 4, and eTable 5

eTable 3. Associations between Perceived Stress Score (PSS) and Hospitalizations: B (SE) and OR (CI) from Generalized Marginal Models.

Covariates	Exposure <sup>a</sup>	<u>CHF Hosp.</u>		<u>Cardiac Hosp.</u>		<u>All Cause Hosp.</u>		<u>All Cause Hosp. or Death</u>	
		B (SE)	OR/CI	B (SE)	OR/CI	B (SE)	OR/CI	B (SE)	OR/CI
Unadjusted	Mean PSS <sup>c</sup>	0.04	1.05	<b>0.05*</b>	<b>1.05</b>	<b>0.06**</b>	<b>1.06</b>	<b>0.07**</b>	<b>1.07</b>
		(0.03)	0.99-1.10	<b>(0.02)</b>	<b>1.01-1.10</b>	<b>(0.02)</b>	<b>1.02-1.11</b>	<b>(0.02)</b>	<b>1.03-1.11</b>
Cov. Adjusted <sup>d</sup>	Mean PSS	0.04	1.04	<b>0.04*</b>	<b>1.05</b>	<b>0.06**</b>	<b>1.06</b>	<b>0.07*</b>	<b>1.07</b>
		(0.03)	0.99-1.10	<b>(0.02)</b>	<b>1.00-1.09</b>	<b>(0.02)</b>	<b>1.02-1.11</b>	<b>(0.02)</b>	<b>1.03-1.11</b>
Adjust. for Other Psych Scales <sup>e</sup>	Mean PSS	0.03	1.03	0.01	1.01	0.01	1.02	0.01	1.01
		(0.04)	0.94-1.12	(0.04)	0.94-1.09	(0.03)	0.95-1.09	(0.04)	.94-1.09
Unadjusted	Deviation PSS <sup>c</sup>	0.01	1.01	0.02	1.03	<b>0.03*</b>	<b>1.03</b>	<b>0.03*</b>	<b>1.03</b>
		(0.02)	0.98-1.04	(0.02)	0.99-1.06	<b>(0.01)</b>	<b>1.00-1.06</b>	<b>(0.01)</b>	<b>1.00-1.06</b>
Cov. Adjusted <sup>d</sup>	Deviation PSS	0.01	1.01	0.03	1.03	<b>0.03*</b>	<b>1.03</b>	<b>0.03*</b>	<b>1.03</b>
		(0.02)	0.97-1.04	(0.02)	0.99-1.06	<b>(0.01)</b>	<b>1.00-1.06</b>	<b>(0.01)</b>	<b>1.00-1.06</b>
Adjust. for Other Psych. Scales <sup>e</sup>	Deviation PSS	0.00	1.00	0.02	1.02	<b>0.03*</b>	<b>1.03</b>	0.03	1.03
		(0.02)	0.96-1.05	(0.02)	0.97-1.07	<b>(0.02)</b>	<b>1.00-1.07</b>	(0.02)	0.94-1.06

*Note.* PSS = Perceived Stress Scale score, HF Hosp. = heart failure-related hospitalizations, Cardiac Hosp. = cardiac-related hospitalizations, All-Cause Hosp. = hospitalizations due to any cause, All Cause Hosp. or Death = all cause hospitalization or death; Cov. Adjusted = analyses adjusted for medical covariates, Adjusted for Other Psych. Scales = analyses adjusted for medical covariates as well as Recent Anger, and STAI-State Anxiety.

<sup>b</sup>Outcomes are hospitalizations in the time prior to the next 6-month follow-up (baseline, 3-month, 9-month, 15-month, 21-month, 27 month, 33-month, and 39-month follow-ups) (see text for details). B (SE) values are parameter estimates from generalized estimating equations models.

<sup>a</sup>Exposure refers to the Mean PSS score and Deviation PSS score for their respective analyses.

<sup>c</sup>Mean PSS scores and Deviation PSS scores are entered concurrently. All models include time as a within-subject variable.

<sup>d</sup>Adjusted analyses include the following covariates: ejection fraction, NYHA class, hypertensive status, presence of type I or type II diabetes, implantable defibrillator status, age, gender, income, and race (parameter estimates for time or covariates not shown).

<sup>e</sup>Analyses including above covariates and STAI-State Anxiety and Recent Anger.

\* $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . These results are also bolded.

eTable 4. Associations between Anxiety (STAI) Score and Outcomes Variables: Parameter estimates B (SE) and OR (CI) from generalized marginal models.

Covariates	Exposure <sup>a</sup>	<u>CHF Hosp.</u>		<u>Cardiac Hosp.</u>		<u>All Cause Hosp.</u>		<u>All Cause Hosp. or Death</u>	
		B (SE)	OR/ CI	B (SE)	OR/ CI	B (SE)	OR/ CI	B (SE)	OR/ CI
Unadjusted	Mean STAI <sup>c</sup>	<b>0.05*</b> (0.02)	<b>1.05</b> 1.01-1.09	<b>0.04**</b> (0.02)	<b>1.05</b> 1.01-1.08	<b>0.06***</b> (0.02)	<b>1.06</b> 1.03-1.09	<b>0.06***</b> (0.02)	<b>1.07</b> 1.03-1.10
Cov. Adjusted <sup>d</sup>	Mean STAI	<b>0.05*</b> (0.02)	<b>1.05</b> 1.01-1.10	<b>0.05**</b> (0.02)	<b>1.05</b> 1.02-1.09	<b>0.06***</b> (0.02)	<b>1.06</b> 1.03-1.10	<b>0.07***</b> (0.02)	<b>1.07</b> 1.04-1.10
Adjust. for Other Psych Scales <sup>e</sup>	Mean STAI	<b>0.06*</b> (0.03)	<b>1.07</b> 1.00-1.14	0.06 (0.03)	1.06 1.00-1.12	<b>0.06*</b> (0.03)	<b>1.07</b> 1.01-1.12	<b>0.07*</b> (0.03)	<b>1.07</b> 1.02-1.13
Unadjusted	Deviation STAI <sup>c</sup>	0.00 (0.01)	1.00 0.97-1.02	0.01 (0.01)	1.01 0.99-1.04	0.01 (0.01)	1.01 0.99-1.04	0.01 (0.01)	1.01 .99-1.03
Cov. Adjusted <sup>d</sup>	Deviation STAI	0.00 (0.01)	1.00 0.97-1.03	0.01 (0.01)	1.01 0.99-1.04	0.01 (0.01)	1.01 0.99-1.03	0.01 (0.01)	1.01 .98-1.03
Adjust. for Other Psych. Scales <sup>e</sup>	Deviation STAI	-0.01 (0.02)	0.99 0.96-1.05	0.00 (0.01)	1.00 0.97-1.04	0.00 (0.01)	1.00 0.98-1.03	0.00 (0.01)	1.00 .98-1.03

Note. STAI = State Trait Anxiety Inventory-State Anxiety score, HF Hosp. = heart failure-related hospitalizations, Cardiac Hosp. = cardiac-related hospitalizations, All-Cause Hosp. = hospitalizations due to any cause, All Cause Hosp. or Death = all cause hospitalization or death;

Cov. Adjusted = analyses adjusted for medical covariates. Adjusted for Other Psych. Scales=analyses adjusted for medical covariates, as well as PSS, and Recent Anger.

<sup>a</sup>Exposure refers to the Mean STAI score and Deviation STAI score for their respective analyses. (See text for details).

<sup>b</sup>Outcomes are hospitalizations occurring after each assessment (baseline, 3-month, 9-month, 15-month, 21-month, 27-month, 33-month, and 39-month follow-ups). B (SE) are parameter estimates from generalized estimating equations models.

<sup>c</sup>Mean STAI scores and Deviation STAI scores are entered concurrently (see text). All models include time as a within-subject variable.

<sup>d</sup>Adjusted analyses for hospitalizations include the following covariates: ejection fraction, NYHA class, hypertensive status, presence of type I or type II diabetes, implantable defibrillator status, age, gender, income, and race (parameter estimates for time or covariates not shown).

<sup>e</sup>Analyses including above covariates and PSS, and Recent Anger.

\* $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . These results are also bolded

eTable 5. Associations between Anger Score and Outcomes Variables: Parameter estimates B (SE) and OR (CI) from generalized marginal models

Covariates	Exposure <sup>a</sup>	<u>CHF Hosp.</u>		<u>Cardiac Hosp.</u>		<u>All Cause Hosp.</u>		<u>All Cause Hosp. or Death</u>	
		B (SE)	OR/ CI	B (SE)	OR/ CI	B (SE)	OR/ CI	B (SE)	OR/ CI
Unadjusted	Mean Anger <sup>c</sup>	-0.02	0.98	0.01	1.01	0.02	1.02	0.02	1.02
		(0.03)	0.93-1.03	(0.02)	0.97-1.04	(0.02)	0.98-1.05	(0.02)	.99-1.05
Cov. Adjusted <sup>d</sup>	Mean Anger	-0.02	0.98	0.00	1.00	0.02	1.02	0.02	1.02
		(0.03)	0.92-1.04	(0.02)	0.97-1.04	(0.02)	0.98-1.05	(0.02)	.99-1.06
Adjust. for Other Psych Scales <sup>e</sup>	Mean Anger	<b>-0.06*</b>	<b>0.94</b>	-0.03	0.97	-0.02	0.98	-0.02	0.98
		<b>(0.03)</b>	<b>0.89-0.99</b>	(0.02)	0.93-1.01	(0.02)	0.94-1.02	(0.02)	.94-1.03
Unadjusted	Deviation Anger <sup>c</sup>	0.02	1.02	0.02	1.02	0.00	1.00	0.00	1.00
		(0.01)	1.00-1.05	(0.01)	0.99-1.04	(0.01)	0.98-1.02	(0.01)	.98-1.03
Cov. Adjusted <sup>d</sup>	Deviation Anger	0.02	1.02	0.02	1.02	0.00	1.00	0.01	1.01
		(0.01)	1.00-1.05	(0.01)	1.00-1.04	(0.01)	0.98-1.02	(0.01)	.98-1.03
Adjust. for Other Psych. Scales <sup>e</sup>	Deviation Anger	0.03	1.03	0.01	1.01	-0.01	0.99	0.00	1.00
		(0.01)	1.00-1.06	(0.01)	0.98-1.04	(0.01)	0.97-1.01	(0.01)	.98-1.02

*Note.* Anger= Recent Anger score, HF Hosp. = heart failure-related hospitalizations, Cardiac Hosp. = cardiac-related hospitalizations, All-Cause Hosp. = hospitalizations due to any cause, All Cause Hosp. or Death = all cause hospitalization or death; Cov. Adjusted = analyses adjusted for covariates, Adjusted for Other Psych. Scales = analyses adjusted for medical covariates as well as for PSS and STAI State Anxiety.

<sup>a</sup>Exposure refers to the Mean Anger and Deviation Anger score for their respective analyses. (See text for details).

<sup>b</sup>Outcomes are hospitalizations occurring after each assessment (baseline, 3-month, 9-month, 15-month, 21-month, 27-month, 33-month, and 39-month follow-ups. B (SE) values are parameter estimates from generalized estimating equation models.

<sup>c</sup>Mean Anger scores and Deviation Anger scores are entered concurrently (see text). All models include time as a within-subject variable.

<sup>d</sup>Adjusted analyses for hospitalizations include the following covariates: ejection fraction, NYHA class, hypertensive status, presence of type I or type II diabetes, implantable defibrillator status, age, gender, income, and race (parameter estimates for time or covariates not shown).

<sup>e</sup>Analyses including above covariates, PSS, and STAI-State Anxiety.

\* $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . These results are also bolded.