

Supplementary Materials

Substantive deviations from pre-registered protocol

The fruit and vegetable outcome variable for RQ2 was pre-registered as an ordinal, categorical change score (increased, no change, reduced intake) computed by comparing intakes at each timepoint (consuming a few portions per day vs all other) to pre-COVID-19 intakes retrospectively reported at baseline (consuming a few portions per day vs all other). However, upon formulating the outcome variable and running models, there was insufficient variance in the data, and GEE models were not converging. This was due to having no participants in the 'all other' ethnicity level for decreasing fruit and vegetable intake. In order to retain the change element of the fruit and vegetable outcome and maximise value for guiding health policy by identifying unhealthy dietary changes, the outcome variable for fruit and vegetable intake was changed to a binary, categorical change outcome (reduced intake vs all other) computed by comparing intakes at each timepoint (consuming a few portions per day vs all other) to pre-COVID-19 intakes retrospectively reported at baseline (consuming a few portions per day vs all other). The GEE models used the logit link function for a binomial distribution.

Given that dietary behaviours are likely to impact on other dietary behaviours, HFSS meals intake explanatory variable was included in the predictors for a change in HFSS snacks, and HFSS snacks was included in the predictors for a change in HFSS meals.

Bayes Factors Robustness checks

Robustness checks were performed using alternative priors. For smoking status, a prior mean difference below -4.0 and above 11.0 HFSS snack portions per month produced the same outcome of no evidence for an effect.

Testing linearity of logit in the binary sensitivity analysis models

Continuous explanatory variables were tested for linearity of logit, and converted to categorical variables where the assumption was violated.

For binary fruit and vegetables RQ2 and complete case analysis, physical activity violated the assumption.

- Physical activity: based on a lower limit cut-off of health recommendations of 150 minutes per week of moderate or 75 minutes of vigorous physical activity equating to 7.5 MET-hours per week [1,2], as the minimum recommended activity levels. (Table 3, Supplementary tables S2, S4 and S5).

For HFSS snacks sensitivity increase or decrease vs all other, HFSS meals violated the assumption.

- HFSS meals median intake: below median vs median and above at each timepoint. Supplementary tables S6 and S7.

Detailed description of study measures

The full wording of questions can be found at: <https://osf.io/bja7g/>

Time-invariant predictors are obtained from participants' baseline responses. Time-variant predictors are collected at baseline, 3- and 6-month follow-up surveys.

Time-invariant

Socio-demographic

- **Gender** (2 levels) *'Which of the following best describes how you think of yourself?'* **female** (female), **all others** (male, non-binary, prefer not to say)
- **Age.** continuous.
- **Ethnicity** (2 levels) **white** (1 white), **all others** (Mixed/Multiple ethnic groups, Asian/Asian British, Black/African/Caribbean/Black British, Chinese, Arab, Other ethnic group, prefer not to say)
- **Baseline occupation and working from home** (3 levels) **Unemployed**, **Employed from home**, **Employed not from home**.
 - All participants were asked, *'What is your current main occupation (during COVID-19)?'* Laid off during COVID-19, unemployed since before COVID-19, retired, homemaker, full-time parent or carer, unable to work due to disability, other, employed (full or part-time), self-employed (full or part-time), student, furloughed during COVID-19.
 - Participants who answered as employed (full or part-time), self-employed (full or part-time), student, furloughed during COVID-19 to the question *'What is your current main occupation (during COVID-19)?'* were asked, *'Can you do your work or study from home?'* Yes, I can do all the work or study from home, I can only do some work or study from home, or No, my work or study cannot be done at home.
 - **Unemployed** (Laid off during COVID-19, Unemployed since before COVID-19, Retired, Homemaker, Full-time parent or carer, Unable to work due to disability, Other), **Employed from home** (Employed AND Yes, I can do all the work or study from home, OR I can only do some work or study from home) **Employed not from home** (Employed AND No, my work or study cannot be done at home).
- **Composite socioeconomic score:** A sum score of education level and combined housing tenure and household income (Sum score 0-3):
 - **Education level.** *'What is the highest level of education that you have completed?'* **0**, below A-level (No formal qualification, GCSE/School certificate/O-level/CSE, Vocational qualifications (e.g. NVQ1+2)), or **1**, A-level or above (A-level/Higher school certificate or equivalent (e.g. NVQ3), Bachelor degree or equivalent (e.g. NVQ4), Masters/PhD/PGCE or equivalent, Other).
 - **Housing and Income.** *'What is your housing tenure?'* Owned outright, Mortgage Rented from local authority, Rented from private landlord, Belongs to housing association, Shared ownership (part owned, part rented), Other. *'What was your household annual income before COVID- 19?'* up to 13 499 GBP, 13 500-24 999 GBP, 25 000-49 999 GBP, ≥50 000 GBP, prefer not to say. **2** (Housing = Owned outright OR mortgage AND Income = 50 000+ GBP), **1** (Housing = All other AND Income = 50 000+ GBP), **1** (Housing = Owned outright OR mortgage AND Income = All other), **0** (Housing = All other AND Income = All other).

COVID-19 related (impact on well-being and lifestyle)

- **Living arrangements (3 levels): alone, with children (with or without adults), with adults only.**
 - **Alone.** Participants responding 'I live on my own' when asked *'How many persons other than yourself (including children) live with you now in the same flat or house?'* Participants answering, 'I live with 1 or more people' were asked: *'Do you live with any of these persons below?'* Being able to select multiple responses: With my partner, husband/wife, boyfriend/girlfriend, with children 0-5 years old, with children 6-15 years old, with persons aged 16-69 (family or friends), with persons aged 70+ (family or friends), persons who you believe may be vulnerable to COVID-19 for any reason, persons who are in poor health

- **With any children (with or without adults)** (if participants respond with 'children 0-5 years old' or 'with children 6-15 years old')
- **Adults only** (all other participants who did not say 'I live on my own' or did not select 'with children 0-5 years old' or 'with children 6-15 years old')

Time-variant

Socio-demographic

- **BMI.** Continuous. Participants were asked at baseline, 3 months and 6 months later, '*How much do you weigh? Please try to be as accurate as possible*'. Participants could answer in 1lb/0.4-5kg increments, or answer 'don't know' or 'prefer not to say'. Participants were also asked '*What is your height? Please try to be as accurate as possible*'. Participants could answer in 1 inch/2-3 cm increments, or answer 'don't know' or 'prefer not to say'. BMI was calculated as weight (kg)/height² (m), by converting height into metres. Prefer not to say or don't know responses were excluded.

COVID-19 related (impact on well-being and lifestyle)

- **Isolation** '*Which type of COVID-19 induced isolation are you experiencing?*' (2 levels) **total and some** (Total isolation/quarantine, Some isolation), **general and no isolation** (General isolation, No isolation)
- **Composite wellbeing, social and living measure:** At baseline: '*How would you rate the below aspects of your life in the since COVID-19? (i) Living, (ii) wellbeing, (iii) social, (iv) family*'. At follow-up: '*How would you rate the below aspects of your life in the past month?*' 1-5: 1 poor, 2, 3 average, 4, 5 excellent. Continuous mean score of living + wellbeing + social + family (1-5)

Behavioural

- **Fruit and vegetables intake.** Continuous, from baseline-since-COVID-19 to 6 months. This predictor was used in the (i) HFSS snacks and (ii) HFSS meals intake analysis only.
- **HFSS snacks and HFSS meals intake.** Continuous, from baseline-since-COVID-19 to 6 months. These predictors were used in the (iii) fruit and vegetables intake analysis only.
- **Physical activity. Continuous.** A continuous measure of MET-hours per week from baseline-since-COVID-19 to 6 months was computed using weekly strength training and assuming a session length of 45 minutes, and using weekly aerobic training frequency and aerobic session length.
 - Weekly strength training. '*Since COVID-19, on average, on how many days per week have you done strength training?*' 0 days per week, 1 day per week, 2 days per week, 3 days per week, 4 days per week or more.
 - Weekly aerobic training. '*Since covid-19, how long (in minutes) was your average session of moderate or vigorous aerobic physical activity?*' (15-480 minutes) and '*Since covid-19 on average, how many times per week have you done 15 minutes or more of moderate or vigorous aerobic physical activity?*' (0,1,2 to 13 times per week, or 14+ times per week).
- **Alcohol consumption. Binary.** Participants were asked '*How often did you have a drink containing alcohol in the past month?*', '*In the past month, how many units of alcohol did you drink on a typical day when you were drinking?*'. Scores were multiplied to compute a binary measure of weekly consumption, categorised into ≤ 14 and > 14 alcohol units.
- **Smoking.** At baseline, '*Which statement about tobacco use and cigarette smoking best describes you?*' At follow-ups, '*2 months ago/previously, we asked about your tobacco use. Your situation may have changed since then, or not. Which of the following best applies to you now?*' (2 levels) **Yes** (I smoke cigarettes (including hand-rolled) every day, I smoke cigarettes (including hand-rolled), but

not every day, I do not smoke cigarettes at all, but I do smoke tobacco of some kind (e.g. pipe, cigar or shisha), **No** (I have stopped smoking completely in the last year (at baseline)/in the past 2/3 months (follow-ups); I stopped smoking completely more than a year ago (baseline)/more than 2/3 months ago (follow-ups), I have never smoked any cigarettes/I have never been a regular smoker (i.e. smoked for a year or more).

Supplementary tables and figures

Figure S1. Changes in portion frequency consumption across the pandemic for (A) HFSS snacks intake, (B) HFSS meals intake and (C) fruit and vegetables intake.

Figure S2. Graphical illustrations of the time-varying associations of gender with meeting fruit and vegetable (FV) recommendations at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.

Figure S3. Graphical illustrations of the time-varying associations of (A) age and (B) alcohol consumption with increasing monthly HFSS snacks intake at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.

Figure S4. Graphical illustrations of the time-varying associations of eating behaviour score with a change in monthly (A) HFSS snacks and (B) meals intake at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.

Table S1. Weighted baseline participant characteristics for the total, included and excluded samples.

Table S2. Univariate unadjusted and adjusted models for each predictor variable and change in self-reported HFSS snacks and meals intake, and reduced FV intake.

Table S3. Unweighted mean changes in HFSS snacks and HFSS meals intake and changes in % meeting fruit and vegetable recommendations using complete cases.

Table S4. Univariate unadjusted and adjusted models for each predictor variable and change in self-reported HFSS snacks and HFSS meals intake, and reduced FV intake using complete cases.

Table S5. Fully adjusted GEE containing all predictor variables and the fully adjusted GEE model including significant explanatory variable*time interactions for a change in HFSS snacks and meals intake, and for reduced FV intake using complete cases.

Table S6. Univariate GEE models for a change in HFSS snacks intake using binary outcomes.

Table S7. Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant explanatory variable*time interactions from univariate models adjusted for time, for a change in HFSS snacks intake with binary outcomes using complete cases.

Table S8. Univariate unadjusted and adjusted models for composite eating behaviour score and change in self-reported HFSS snacks and HFSS meals intake, and reduced FV intake using complete cases.

Table S9. Fully adjusted models for a change in HFSS snacks and meals intake, and for reduced FV intake using complete cases, including the eating behaviour score explanatory variable.

Figure S1. Changes in portion frequency consumption across the pandemic for (A) HFSS snacks intake, (B) HFSS meals intake and (C) fruit and vegetables intake.




Figure S2. Graphical illustrations of the time-varying associations of gender with meeting fruit and vegetable recommendations at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.





Figure S3. Graphical illustrations of the time-varying associations of (A) age and (B) alcohol consumption with increasing monthly HFSS snacks intake at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.



Age categories: <25, 4.6%; 25-44, 23.1%; 45-64, 53.4%; >65, 19.0%.

Figure S4. Graphical illustrations of the time-varying associations of eating behaviour score with a change in monthly (A) HFSS snacks and (B) meals intake at baseline (May-June 2020), 3-months (August-September 2020) and 6-months follow-up (November-December 2020) from complete case analyses.




Table S1. Weighted baseline participant characteristics for the total, included and excluded samples.

	Total Sample Weighted (%)	Included sample Weighted (%)	Excluded sample Weighted (%)	p- value
N	2805	1532	1274	
Gender				
All other	48.3%	45.8%	51.3%	0.004
Female	51.7%	54.2%	48.7%	
Ethnicity				<0.001
All other	11.5%	8.0%	15.7%	
White	88.5%	92.0%	84.3%	
Mean BMI [SD] N= 2614	26.4 [5.2]	26.7 [5.2]	26.0 [5.2]	<0.001
Mean Age [SD]	48.3 [16.7]	52.9 [14.8]	42.8 [17.2]	<0.001
Occupation and work from home N= 2806				<0.001
Unemployed (retired persons and full-time parents/carers)	30.9%	35.7%	25.1%	
Employed working from home	40.4%	37.9%	43.4%	
Employed not working from home	28.7%	26.4%	31.5%	
Socioeconomic score				<0.001
Income <£50k, unowned Housing and no higher education	13.0%	10.3%	16.4%	
1 of: ≥£50K income, housing ownership/mortgage or higher education	40.5%	39.2%	42.0%	
2 of: ≥£50K income, housing ownership/mortgage or higher education	34.7%	37.6%	31.1%	
All of: ≥£50K income, housing ownership/mortgage and higher education	11.8%	12.9%	10.5%	
Living conditions				<0.001
Alone	18.8%	19.7%	17.8%	
With children (with or without adults)	17.3%	14.4%	20.7%	
With adults only	63.9%	65.9%	61.5%	
Isolation status N= 2761				<0.001
Total or some isolation	75.8%	79.0%	71.8%	
General or no isolation	24.2%	21.0%	28.2%	
Mean Quality of Life [SD] N=2704	3.2 [0.9]	3.3 [0.8]	3.1 [0.9]	<0.001
Mean HFSS Snacks portions per month N=2395	61.1 [49.9]	61.6 [50.5]	60.3 [48.8]	0.531
Mean HFSS Meals portions per month N=2409	7.4 [9.2]	6.6 [7.2]	8.9 [11.8]	<0.001
Mean Fruit and vegetables portions per month N=2438	39.4 [19.7]	40.5 [19.2]	37.7 [20.3]	0.001
Mean physical activity MET-Hours per week N=2587	17.6 [19.8]	18.3 [20.3]	16.7 [19.1]	0.040
Alcohol intake N=2541				0.044
≤14 weekly units	80.8%	82.1%	78.9%	
>14 weekly units	19.2%	17.9%	21.1%	

Smoking status N=2806				<0.001
Yes	26.5%	18.6%	35.9%	
No	73.5%	81.4%	64.1%	

SD: Standard Deviation.

Table S2. Univariate unadjusted and adjusted models for each predictor variable and change in self-reported HFSS snacks and meals intake, and reduced FV intake.

Predictor	Change in self-reported HFSS Snacks				Change in self-reported HFSS meals				Reduced FV intake			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p
Gender	12.220	<0.001	11.141	0.001	5.684	0.017	5.725	0.017	0.128	0.721	39.477	0.597
Time			192.667	<0.001			14.336	0.001			39.477	<0.001
Time*Gender			1.129	0.569			0.064	0.969			6.802	0.033
Ethnicity	0.544	0.461	0.697	0.404	<0.001	0.974	0.001	0.974	0.004	0.950	0.017	0.897
Time			41.321	<0.001			7.316	0.026			10.01	0.007
Time*Ethnicity			1.585	0.453			1.032	0.597			0.790	0.674
BMI	5.278	0.022	5.152	0.023	1.282	0.258	1.820	0.177	15.118	<0.001	17.188	<0.001
Time			0.486	0.784			3.841	0.147			8.034	<0.001
Time*BMI			6.882	0.032			4.648	0.098			2.411	0.300
Age	0.001	0.976	0.293	0.588	0.005	0.945	0.020	0.886	2.264	0.132	2.319	0.128
Time			71.567	<0.001			5.089	0.079			2.271	0.321
Time*Age			17.175	<0.001			1.815	0.404			0.200	0.905
Occupation and work from home	0.788	0.674	0.224	0.894	0.357	0.836	0.217	0.897	1.684	0.431	2.586	0.274
Time			223.915	<0.001			14.285	0.001			35.595	<0.001
Time*Occupation and work from home			7.140	0.129			2.355	0.671			2.642	0.619

Socioeconomic score	4.623	0.202	4.419	0.220	4.193	0.241	<0.001	1.000	9.647	0.022	8.712	0.033
Time			87.866	<0.001				10.964	0.004		20.027	<0.001
Time*Socioeconomic score			14.752	0.022				2.623	0.758		1.303	0.971
Living arrangements	1.084	0.582	0.962	0.618	5.822	0.054	5.991	0.050	3.505	0.173	2.838	0.242
Time			161.284	<0.001			21.977	<0.001			38.118	<0.001
Time*Living arrangements			6.349	0.175			4.819	0.306			4.057	0.398
Isolation status	81.509	<0.001	0.276	0.599	0.009	0.925	0.499	0.480	4.823	0.028	34.305	<0.001
Time			156.998	<0.001			22.932	<0.001			0.005	0.945
Time*Isolation status			1.528	0.466			0.346	0.841			1.942	0.379
Quality of life	39.870	<0.001	13.717	<0.001	0.005	0.946	0.193	0.661	9.897	0.002	12.070	0.001
Time			36.370	<0.001			4.75	0.093			1.453	0.484
Time*Quality of life			11.068	0.004			3.664	0.160			4.418	0.110
HFSS snacks intake					14.814	<0.001	16.668	<0.001	3.431	0.064	8.427	0.004
Time							0.329	0.848			25.778	<0.0001
Time*HFSS snacks intake							3.969	0.137			2.215	0.330
HFSS meals intake	44.221	<0.001	48.416	<0.001					17.063	<0.001	6.591	0.010
Time			77.207	<0.001							30.703	<0.001
Time*HFSS meals intake			18.409	<0.001							5.536	0.063

Fruit and vegetables intake	0.165	0.685	<0.001	0.993	0.748	0.387	0.297	0.586				
Time			66.725	<0.001			6.467	0.039				
Time*Fruit and vegetables intake			15.214	<0.001			2.886	0.236				
Physical activity	5.144	0.023	8.351	0.004	2.181	0.140	1.752	0.186	18.796	<0.001	15.154	<0.001
Time			157.374	<0.001			16.091	<0.001			41.646	<0.001
Time*Physical activity			8.200	0.017			2.896	0.235			0.097	0.953
Alcohol intake	0.001	0.970	1.790	0.181	0.138	0.710	0.114	0.735	0.242	0.622	0.079	0.779
Time			192.202	<0.001			15.872	<0.001			26.537	<0.001
Time*Alcohol intake			3.020	0.221			0.273	0.873			2.388	0.303
Smoking status	0.008	0.931	0.302	0.583	1.127	0.288	0.84	0.359	0.278	0.598	0.037	0.847
Time			117.28	<0.001			19.899	<0.001			29.553	<0.001
Time*Smoking status			2.352	0.309			5.043	0.080			1.934	0.380

Missingness depends on if participants responded to survey questions relating to the predictor variable. For the binary fruit and vegetables analysis, physical activity violated the linearity of logit assumption, and was converted to a binary variable of below minimum health recommendations (<7.5 MET-hours per week) vs meeting minimum health recommendations (≥ 7.5 MET-hours per week). $W\chi^2$: Wald Chi-square.

Table S3. Unweighted mean changes in HFSS snacks and HFSS meals intake and changes in the proportion meeting fruit and vegetable recommendations using complete cases.

	N	Change in HFSS Snacks		Change in HFSS Meals		% Meeting FV recommendations	
		Mean difference [95% CI]	p	Mean difference [95% CI]	p	Mean difference % [95% CI]	p
Pre-pandemic - May-Jun	1486	9.2 [7.2,11.2]	<0.001	-1.3 [0.9,1.7]	<0.001	0.00 [-0.02,0.02]	0.993
Pre-pandemic - Aug-Sep	1486	-4.9 [-6.8,-3.1]	<0.001	-1.3 [0.9,1.8]	<0.001	0.00 [-0.03,0.02]	0.993
Pre-pandemic - Nov-Dec	1486	0.3 [-1.5,2.0]	0.774	-0.7 [-1.2,-0.2]	0.001	0.02 [-0.01,0.05]	0.379
May-Jun - Aug-Sep	1486	-14.1 [-16.4,-11.9]	<0.001	-0.03 [-0.3,0.3]	0.824	0.00 [-0.03,0.02]	0.993
Aug-Sep - Nov-Dec	1486	5.2 [3.5,6.9]	<0.001	0.6 [0.2,1.0]	0.001	0.02 [-0.01,0.05]	0.298
May-Jun - Nov-Dec	1486	-8.9 [-11.4,-6.4]	<0.001	0.6 [0.2,0.9]	0.001	0.02 [-0.01,0.05]	0.379

CI: Confidence Interval.

Table S4. Univariate unadjusted and adjusted models for each predictor variable and change in self-reported HFSS snacks and HFSS meals intake, and reduced FV intake using complete cases.

Predictor	Change in self-reported HFSS Snacks intake				Change in self-reported HFSS meals intake				Reduced FV intake			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p	W χ^2	p
Gender	10.469	0.001	9.864	0.002	5.390	0.020	5.459	0.019	0.003	0.953	0.059	0.809
Time			187.119	<0.001			11.749	0.003			29.464	<0.001
Time*Gender			2.392	0.302			0.485	0.784			6.469	0.039
Ethnicity	0.975	0.323	1.147	0.284	2.930	0.087	3.211	0.073	0.001	0.976	0.073	0.787
Time			42.283	<0.001			8.162	0.017			8.293	0.016
Time*Ethnicity			1.125	0.570			1.332	0.514			1.293	0.524
BMI	4.141	0.042	4.475	0.034	1.443	0.230	1.839	0.175	19.17	<0.001	21.595	<0.001
Time			0.584	0.747			3.604	0.165			9.540	0.008
Time*BMI			6.827	0.033			4.717	0.095			4.259	0.119
Age	0.234	0.628	0.754	0.385	0.219	0.640	0.411	0.522	0.596	0.440	0.605	0.437
Time			68.023	<0.001			3.157	0.206			2.334	0.311
Time*Age			16.053	<0.001			1.533	0.465			0.007	0.997
Occupation and work from home	0.156	0.925	0.080	0.961	0.973	0.615	0.682	0.711	1.337	0.512	1.631	0.442
Time			218.606	<0.001			9.808	0.007			24.722	<0.001
Time*Occupation and work from home			9.701	0.046			3.857	0.426			1.426	0.840

Socioeconomic score	2.476	0.480	**	**	2.216	0.529	1.977	0.577	6.075	0.108	**	**
Time			73.356	<0.001			9.516	0.009			**	**
Time*Socioeconomic score			7.292	0.063			5.578	0.472			**	**
Living arrangements	1.066	0.587	1.058	0.589	5.101	0.078	5.339	0.069	5.408	0.067	4.758	0.093
Time			154.476	<0.001			12.034	0.002			26.309	<0.001
Time*Living arrangements			8.838	0.065			2.132	0.711			4.47	0.346
Isolation status		<0.001										
	94.128	1	0.131	0.718	0.058	0.810	0.590	0.442	2.326	0.127	0.313	0.576
Time			144.098	<0.001			16.252	<0.001			23.867	<0.001
Time*Isolation status			1.526	0.466			0.154	0.926			1.987	0.370
Quality of life		<0.001										
	37.700	1	11.477	0.001	0.030	0.862	0.131	0.717	6.844	0.009	8.791	0.003
Time			34.275	<0.001			4.228	0.121			1.183	0.553
Time*Quality of life			10.814	0.004			3.745	0.154			2.718	0.257
HFSS snacks intake					16.672	<0.001	18.789	<0.001	3.414	0.065	7.774	0.005
Time							0.766	0.682			16.548	<0.001
Time*HFSS snacks intake							1.857	0.395			1.419	0.492
HFSS meals intake		<0.001										
	27.342	1	35.700	<0.001					14.570	<0.001	7.007	0.008
Time			65.614	<0.001							19.123	<0.001
Time*HFSS meals intake			18.077	<0.001							6.700	0.035

Impact of COVID-19 pandemic on diet behaviour among UK adults: a longitudinal analysis of data from the HEBECO study

Fruit and vegetables intake	0.054	0.816	0.222	0.638	0.510	0.475	0.294	0.588				
Time			61.166	<0.001			4.772	0.092				
Time*Fruit and vegetables intake			13.087	0.001			2.355	0.308				
Physical activity (MET-hours per week)	5.039	0.025	9.338	0.002	2.598	0.107	2.310	0.129	19.970	<0.001	15.338	<0.001
Time			148.862	<0.001			7.461	0.024			28.332	<0.001
Time*Physical activity			8.271	0.016			0.323	0.851			1.390	0.499
Alcohol consumption	0.159	0.690	0.688	0.407	0.008	0.929	0.005	0.945	0.001	0.973	0.002	0.966
Time			182.269	<0.001			15.732	<0.001			16.573	<0.001
Time*Alcohol consumption			4.165	0.125			0.559	0.756			1.790	0.409
Smoking status	0.614	0.433	1.267	0.260	2.307	0.129	1.867	0.172	1.515	0.218	0.744	0.388
Time			113.324	<0.001			9.506	0.009			21.952	<0.001
Time*Smoking status			4.617	0.099			1.165	0.559			3.022	0.221

****Model did not compute due to numerical errors. For the binary fruit and vegetables analysis, physical activity violated the linearity of logit assumption, and was converted to a binary variable of below minimum health recommendations (<7.5 MET-hours per week) vs meeting minimum health recommendations (≥7.5 MET-hours per week). Missingness with each model depends on if participants responded to survey questions relating to the predictor variable. W χ^2 : Wald Chi-square.**

Table S5. Fully adjusted GEE containing all predictor variables and the fully adjusted GEE model including significant predictor*time interactions for a change in HFSS snacks and meals intake, and for reduced FV intake using complete cases.

Change in self-reported HFSS Snacks QIC = 4214122.36			Change in self-reported HFSS Meals QIC = 184695.497		Reduced FV intake QIC = 2282.216	
All predictors (N=1451)	p-value	B [95% CI]	p-value	B [95% CI]	p-value	OR [95% CI]
Gender	<0.001		0.010		0.737	
All other		Reference		Reference		Reference
Female		6.210 [3.127,9.294]		0.948 [0.227,1.668]		0.949 [0.697,1.290]
Ethnicity	0.110		0.064		0.753	
All other		Reference		Reference		Reference
White		-6.051 [-13.483,1.380]		-1.123 [-2.312,0.065]		0.889 [0.426,1.853]
BMI	0.166	0.233 [-0.097,0.564]	0.084	-0.056 [-0.119,0.008]	0.006	1.035 [1.010,1.061]
Age	0.062	0.126 [-0.006,0.259]	0.496	0.010 [-0.018,0.038]	0.625	0.997 [0.984,1.010]
Occupation and work from home	0.597		0.692		0.655	
Unemployed		Reference		Reference		Reference
Employed working from home		1.972 [-1.894,5.839]		0.077 [-0.982,0.828]		1.204 [0.811,1.788]
Employed not working from home		1.626 [-3.080,6.332]		-0.412 [-1.485,0.661]		1.145 [0.727,1.805]
Socioeconomic score	0.258		0.384		0.410	
Income <£50k, unowned Housing and no higher education		Reference		Reference		Reference
1 of: ≥£50K income, housing ownership/mortgage or higher education		0.684 [-10.392,11.761]		1.739 [-1.103,4.580]		1.238 [0.473,3.240]
2 of: ≥£50K income, housing ownership/mortgage or higher education		4.401 [-6.592,15.393]		2.076 [-0.695,4.846]		1.420 [0.550,3.665]
All of: ≥£50K income, housing ownership/mortgage and higher education		3.692 [-7.167,14.551]		2.134 [-0.629,4.897]		1.078 [0.408,2.852]
Living conditions	0.473		0.039		0.938	
Alone		Reference		Reference		Reference
With children (with or without adults)		2.181 [-3.082,7.444]		-0.476 [-1.901,0.948]		0.990 [0.586,1.672]

With adults only		2.439 [-1.471,6.348]		0.658 [-0.375,1.691]		0.943 [0.640,1.390]
Isolation status	0.507		0.373		0.709	
Total or some isolation		Reference		Reference		Reference
General or no isolation		-0.711[-2.813,1.391]		0.188 [-0.226,0.602]		1.053 [0.803,1.380]
Quality of Life	<0.001	-2.785 [-4.318,-1.252]	0.971	0.005 [-0.288,0.299]	0.098	0.867 [0.731,1.027]
HFSS Snacks			<0.001	0.018 [0.010,0.027]	0.324	1.002 [0.998,1.005]
HFSS Meals	<0.001	0.731 [0.453,1.008]			0.142	1.013 [0.996,1.031]
Fruit and vegetables	0.401	0.034 [-0.045,0.112]	0.306	-0.008 [-0.025,0.008]		
Physical activity (MET-Hours per week)	0.102	-0.036 [-0.080,0.007]	0.074	-0.009 [-0.018,0.001]	0.004	0.671 [0.511,0.881]*
Alcohol intake	0.141		0.441		0.960	
≤14 weekly units		Reference		Reference		Reference
>14 weekly units		1.659 [-0.548,3.867]		0.186 [-0.287,0.658]		1.008 [0.740,1.373]
Smoking status	0.238		0.283		0.544	
Yes		Reference		Reference		Reference
No		2.600 [-1.720,6.920]		-0.461 [-1.304,0.381]		0.884 [0.592,1.318]

All predictors + significant full model interactions		Change in self-reported HFSS Snacks QIC = 4187090.57		Reduced FV intake QIC = 2280.087	
	p-value				p-value
Time*Age	0.016				
Time*HFSS meals intake	0.001				
Time*Gender				0.042	

*For the fully adjusted GEE model including explanatory variable*time interactions, Type III tests for the explanatory variable*time interactions are shown only. QIC is a relative, 'lower is better' measure of goodness of fit. For the binary fruit and vegetables analysis, physical activity violated the linearity of logit assumption, and was converted to a binary variable of below minimum health recommendations (<7.5 MET-hours per week) vs meeting minimum health recommendations (≥7.5 MET-hours per week). *OR for 'meeting minimum health recommendations' with 'below minimum health recommendations' as reference. W χ^2 : Wald Chi-square, B: Beta parameter, SE: Standard Error, CI: Confidence Interval, OR: Odds Ratio.*

Table S6. Univariate GEE models for a change in HFSS snacks intake using binary outcomes.

Change in self-reported HFSS Snacks								
Predictor	Increase vs all other				Decrease vs all other			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	<i>W</i> χ^2	p-value	<i>W</i> χ^2	p-value	<i>W</i> χ^2	p-value	<i>W</i> χ^2	p-value
Gender	11.269	<0.001	10.735	0.001	8.883	0.003	9.027	0.003
Time			53.031	<0.001			190.436	<0.001
Time*Gender			3.305	0.192			2.311	0.315
Ethnicity	0.004	0.949	0.003	0.956	0.054	0.816	0.072	0.788
Time			6.970	0.031			31.307	<0.001
Time*Ethnicity			1.272	0.529			0.839	0.657
BMI	4.782	0.029	4.851	0.028	0.348	0.555	0.640	0.424
Time			0.360	0.835			6.840	0.033
Time*BMI			1.332	0.514			0.082	0.960
Age	1.107	0.293	0.662	0.416	2.363	0.124	2.572	0.109
Time			30.120	<0.001			25.181	<0.001
Time*Age			15.183	0.001			1.816	0.403
Occupation and work from home	1.139	0.566	0.822	0.663	0.675	0.713	0.827	0.661
Time			56.224	<0.001			207.576	<0.001
Time*Occupation and work from home			8.393	0.078			5.067	0.280

Socioeconomic score	3.288	0.349	2.850	0.415	8.228	0.042	8.367	0.039
Time			23.238	<0.001			92.627	<0.001
Time*Socioeconomic score			8.147	0.228			1.341	0.969
Living arrangements	1.864	0.394	2.180	0.336	6.675	0.036	7.481	0.024
Time			47.547	<0.001			150.283	<0.001
Time*Living arrangements			7.303	0.121			6.351	0.174
Isolation status	34.294	<0.001	1.266	0.261	74.726	<0.001	0.104	0.747
Time			31.881	<0.001			141.312	<0.001
Time*Isolation status			3.523	0.172			0.396	0.821
Quality of life	15.749	<0.001	6.046	0.014	13.323	<0.001	2.132	0.144
Time			8.893	0.012			11.559	0.003
Time*Quality of life			2.466	0.291			0.098	0.952
HFSS meals intake	19.176	<0.001	23.181	<0.001	16.208	<0.001	24.190	<0.001
Time			73.444	<0.001			240.892	<0.001
Time*HFSS meals intake			2.692	0.260			2.901	0.234
Fruit and vegetables	0.195	0.658	0.079	0.779	0.511	0.475	0.742	0.389
Time			12.075	0.002			34.272	<0.001
Time*Fruit and vegetables			0.576	0.750			1.061	0.588

Physical activity	1.577	0.209	2.153	0.142	0.019	0.891	0.500	0.480
Time			34.026	<0.001			113.971	<0.001
Time*Physical activity			1.658	0.437			0.011	0.994
Alcohol consumption	0.298	0.585	1.182	0.277	0.182	0.670	3.340	0.068
Time			63.287	<0.001			156.775	<0.001
Time*Alcohol consumption			8.884	0.012			5.309	0.070
Smoking status	0.261	0.609	0.096	0.756	0.049	0.825	0.085	0.770
Time			33.454	<0.001			112.573	<0.001
Time*Smoking status			2.868	0.238			3.099	0.212

Due to violating the linearity of logit assumption, HFSS meals intake was converted to a binary variable of 'below median' and 'median and above' intakes. Missingness with each model depends on if participants responded to survey questions relating to the predictor variable. $W\chi^2$: Wald Chi-square.

Table S7. Full GEE model containing all predictor variables and the full GEE model containing all predictor variables and significant explanatory variable*time interactions from univariate models adjusted for time, for a change in HFSS snacks intake using binary outcomes using complete cases.

Change in self-reported HFSS snacks intake						
N=1451	Increase vs all other			Decrease vs all other		
	Type III Test			Type III Test		
	W χ^2	p-value	QIC	W χ^2	p-value	QIC
All predictors			5588.258			5278.921
Gender	15.120	<0.001		12.365	<0.001	
Ethnicity	0.427	0.514		1.419	0.234	
BMI	4.057	0.044		0.028	0.867	
Age	0.014	0.905		10.347	0.001	
Occupation and work from home	0.753	0.686		2.768	0.251	
Socioeconomic score	4.697	0.195		6.537	0.088	
Living conditions	1.049	0.592		5.08	0.079	
Isolation status	1.648	0.199		0.704	0.401	
Quality of Life	7.634	0.006		7.022	0.008	
HFSS meals intake	19.344	<0.001		24.292	<0.001	
Fruit and vegetables	0.003	0.958		0.837	0.360	
Physical activity (MET-Hours per week)	0.073	0.788		0.066	0.798	
Alcohol consumption	3.161	0.075		5.442	0.020	
Smoking status	0.001	0.970		0.017	0.896	
All predictors and significant interactions: N=1451			5574.692			
Time*Age	14.137	0.001				
Time*Alcohol consumption	9.825	0.007				

Models also included Time as a covariate. Type III tests are shown only. Due to violating the linearity of logit assumption, HFSS meals intake was converted to a binary variable of 'below median' and 'median and above' intakes. There were no material changes in significance of main effects with the addition of time interactions, except for baseline BMI for an increase in HFSS snacks vs all other, which became non-significant. QIC is a relative, 'lower is better' measure of goodness of fit. W χ^2 : Wald Chi-square.

Table S8. Univariate unadjusted and adjusted models for composite eating behaviour score and change in self-reported HFSS snacks and HFSS meals intake, and reduced FV intake using complete cases.

Predictor	Change in self-reported HFSS Snacks intake				Change in self-reported HFSS meals intake				Reduced FV intake			
	Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction		Unadjusted models		Adjusted models w/interaction	
	W X-S	p	W X-S	p	W X-S	p	W X-S	p	W X-S	p	W X-S	p
Composite eating behaviour score N=1193	25.474	<0.001	18.645	<0.001	0.204	0.652	<0.001	0.984	6.099	0.014	6.248	0.012
Time			10.456	0.005			14.720	0.001			5.914	0.052
Time*Composite eating behaviour score			72.869	<0.001			9.258	0.010			0.246	0.884

W χ^2 : Wald Chi-square.

Table S9. Fully adjusted models for a change in HFSS snacks and meals intake, and for reduced FV intake using complete cases, including eating behaviour score.

	Change in self-reported HFSS Snacks		Change in self-reported HFSS Meals		Reduced FV intake	
	QIC = 3183339.18		QIC = 135661.533		QIC = 1802.887	
All predictors (N=1166)	W χ^2	p-value	W χ^2	p-value	W χ^2	p-value
Eating Behaviour score	13.558	<0.001	0.118	0.731	0.326	0.568
All predictors and significant interactions (N=1166)	Change in self-reported HFSS Snacks		Change in self-reported HFSS Meals			
	QIC = 3127023.43		QIC = 135368.316			
Time*Eating Behaviour score	50.783	<0.001	7.351	0.025		

Models also included Time as a covariate. Type III tests are shown only for Eating behaviour score in the model. QIC is a relative, 'lower is better' measure of goodness of fit. W χ^2 : Wald Chi-square.

References:

- [1] Lee I-M. Physical Activity and Weight Gain Prevention. JAMA 2010;303:1173. <https://doi.org/10.1001/jama.2010.312>.
- [2] Kaminsky LA, Montoye AHK. Physical Activity and Health: What Is the Best Dose? J Am Heart Assoc n.d.;3:e001430. <https://doi.org/10.1161/JAHA.114.001430>.