

# Supplementary Materials

## Supplementary Materials 1





### Pre-analysis data cleaning

**Participants (N = 237) were excluded based on meeting one or more of the following pre-registered quality control checks:**

1. Completed the study quicker than would be expected given a reasonable reading rate (8 minutes, 5th percentile of soft-launch data)
2. Failed to answer attention questions appropriately (e.g., place slider in X position)
3. Provided discrepant data during screening and the main survey regarding the COVID-19 vaccine they had previously received (i.e., AstraZeneca or Pfizer)
4. Identified as having experienced side effects to previous COVID-19 vaccines, then did not select any side effects
5. Spent more than 5 minutes on the 2-minute timed manipulation page without proceeding
6. Identified as having previously received a specific COVID-19 vaccine, then stated they had never heard of that vaccine

# Supplementary Materials 2

## Wording used for the Positive and Negative Attribute Frame

Negative Attribute Frame (Standard EU Wording)	Positive Attribute Frame	Positive Attribute Frame (Accompanying Infographic)
<p><b>Very common</b> (may affect more than 1 in 10 people)</p>	<p><b>Very common</b> (9 in 10 or fewer people may <b><i>not be affected</i></b>)</p>	 <p><b>Very Common:</b> 9 in 10 or fewer <b>not affected</b> (light to dark blue circles)</p>
<p><b>Common</b> (may affect up to 1 in 10 people)</p>	<p><b>Common</b> (90 in 100 or more people may <b><i>not be affected</i></b>)</p>	 <p><b>Common:</b> 90 in 100 (blue circles) or more <b>not affected</b></p>
<p><b>Uncommon</b> (may affect up to 1 in 100 people)</p>	<p><b>Uncommon</b> (990 in 1,000 or more people may <b><i>not be affected</i></b>)</p>	 <p><b>Uncommon:</b> 990 in 1,000 (blue circles) or more <b>not affected</b></p>
<p><b>Rare</b> (may affect up to 1 in 1,000 people)</p>	<p><b>Rare</b> (9,990 in 10,000 or more people may <b><i>not be affected</i></b>)</p>	 <p><b>Rare:</b> 9,990 in 10,000 (blue circles) or more <b>not affected</b></p>
<p><b>Very Rare</b> (may affect up to 1 in 10,000 people)</p>	<p><b>Very Rare</b> (99,990 in 100,000 or more people may <b><i>not be affected</i></b>)</p>	 <p><b>Very Rare:</b> 99,990 in 100,000 (blue circles) or more <b>not affected</b></p>

## **Supplementary Materials 3**

### **Patient Information Leaflets (PILs) employed in the present experiment**

The following pages contain the PILs used to present side effect information.

High fidelity versions of these documents were embedded in Qualtrics.

# Negative PIL (AstraZeneca Vaccine)

## Package leaflet: Information for the user

### Vaxzevria suspension for injection COVID-19 Vaccine (ChAdOx1-S [recombinant])

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

**Read all of this leaflet carefully before the vaccine is given because it contains important information for you.**

#### What Vaxzevria is and what it is used for

Vaxzevria is used for preventing COVID-19 caused by the SARS-CoV-2 virus.

Vaxzevria is given to adults aged 18 years and older.

The vaccine causes the immune system (the body's natural defences) to produce antibodies and specialised white blood cells that work against the virus, so giving protection against COVID-19.

None of the ingredients in this vaccine can cause COVID-19

#### Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

#### The following side effects may occur with Vaxzevria:

##### Very Common (may affect more than 1 in 10 people)

- tenderness, pain, warmth, itching, or bruising where the injection is given
- feeling tired (fatigue) or generally feeling unwell
- chills or feeling feverish
- headache
- feeling sick (nausea)
- joint pain or muscle ache

##### Common (may affect up to 1 in 10 people)

- swelling or redness where the injection is given
- fever ( $\geq 38^{\circ}\text{C}$ )
- being sick (vomiting) or diarrhoea
- low level of blood platelets
- pain in legs or arms
- flu-like symptoms, such as high temperature, sore throat, runny nose, cough and chills
- physical weakness or lack of energy

##### Uncommon (may affect up to 1 in 100 people)

- sleepiness or feeling dizzy
- abdominal pain or decreased appetite
- enlarged lymph nodes
- excessive sweating, itchy skin, rash or hives
- sleepiness or deep unresponsiveness and inactivity

##### Very Rare (may affect up to 1 in 10,000 people)

- blood clots often in unusual locations (e.g. brain, bowel, liver, spleen) in combination with low level of blood platelets
- serious nerve inflammation, which may cause paralysis and difficulty breathing (Guillain-Barré syndrome [GBS])

##### Not known (cannot be estimated from the available data)

- severe allergic reaction (anaphylaxis)
- hypersensitivity
- rapid swelling under the skin in areas such as the face, lips, mouth and throat (which may cause difficulty in swallowing or breathing)
- capillary leak syndrome (a condition causing fluid leakage from small blood vessels)

# Positive PIL (AstraZeneca Vaccine)

Package leaflet: Information for the user  
Vaxzevria suspension for injection  
COVID-19 Vaccine (ChAdOx1-S [recombinant])

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

Read all of this leaflet carefully before the vaccine is given because it contains important information for you.

## What Vaxzevria is and what it is used for

Vaxzevria is used for preventing COVID-19 caused by the SARS-CoV-2 virus.

Vaxzevria is given to adults aged 18 years and older.

The vaccine causes the immune system (the body's natural defences) to produce antibodies and specialised white blood cells that work against the virus, so giving protection against COVID-19. None of the ingredients in this vaccine can cause COVID-19

## Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

### The following side effects may occur with Vaxzevria:

#### Very Common (9 in 10 or fewer people may *not be affected*)

- tenderness, pain, warmth, itching, or bruising where the injection is given
- feeling tired (fatigue) or generally feeling unwell
- chills or feeling feverish
- headache
- feeling sick (nausea)
- joint pain or muscle ache

#### Common (90 in 100 or more people may *not be affected*)

- swelling or redness where the injection is given
- fever ( $\geq 38^{\circ}\text{C}$ )
- being sick (vomiting) or diarrhoea
- low level of blood platelets
- pain in legs or arms
- flu-like symptoms, such as high temperature, sore throat, runny nose, cough and chills
- physical weakness or lack of energy

#### Uncommon (990 in 1,000 or more people may *not be affected*)

- sleepiness or feeling dizzy
- abdominal pain or decreased appetite
- enlarged lymph nodes
- excessive sweating, itchy skin, rash or hives
- sleepiness or deep unresponsiveness and inactivity

#### Very Rare (99,990 in 100,000 or more people may *not be affected*)

- blood clots often in unusual locations (e.g. brain, bowel, liver, spleen) in combination with low level of blood platelets
- serious nerve inflammation, which may cause paralysis and difficulty breathing (Guillain-Barré syndrome [GBS])

#### Not known (cannot be estimated from the available data)


- severe allergic reaction (anaphylaxis)
- hypersensitivity
- rapid swelling under the skin in areas such as the face, lips, mouth and throat (which may cause difficulty in swallowing or breathing)
- capillary leak syndrome (a condition causing fluid leakage from small blood vessels)

In the images below:  
● Blue circles represent people **not affected**  
● Yellow circles represent people who **are affected**


Very Common: 9 in 10 or fewer **not affected** (light to dark blue circles)




Common: 90 in 100 (blue circles) or more **not affected**



Uncommon: 990 in 1,000 (blue circles) or more **not affected**



Very Rare: 99,990 in 100,000 (blue circles) or more **not affected**



# Negative PIL (Pfizer Vaccine)

## Package leaflet: Information for the user

### Comirnaty concentrate for dispersion for injection COVID-19 mRNA Vaccine (nucleoside modified)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

**Read all of this leaflet carefully before you receive this vaccine because it contains important information for you.**

#### What Comirnaty is and what it is used for

Comirnaty is a vaccine used for preventing COVID-19 caused by SARS-CoV-2 virus.

Comirnaty is given to adults and adolescents from 12 years of age and older.

The vaccine causes the immune system (the body's natural defences) to produce antibodies and blood cells that work against the virus, so giving protection against COVID-19.

As Comirnaty does not contain the virus to produce immunity, it cannot give you COVID-19.

#### Possible side effects

Like all vaccines, Comirnaty can cause side effects, although not everybody gets them.

#### The following side effects may occur with Comirnaty:

##### Very common side effects (may affect more than 1 in 10 people)

- injection site: pain, swelling
- tiredness
- headache
- muscle pain
- chills
- joint pain
- diarrhoea
- fever

Some of these side effects were slightly more frequent in adolescents 12 to 15 years than in adults.

##### Common side effects (may affect up to 1 in 10 people)

- injection site redness
- nausea
- vomiting

##### Uncommon side effects (may affect up to 1 in 100 people)

- enlarged lymph nodes
- feeling unwell
- arm pain
- insomnia
- injection site itching
- allergic reactions such as rash or itching

##### Rare side effects (may affect up to 1 in 1,000 people)

- temporary one sided facial drooping
- allergic reactions such as hives or swelling of the face

##### Not known (cannot be estimated from the available data)

- severe allergic reaction
- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain
- extensive swelling of the vaccinated limb
- swelling of the face (swelling of the face may occur in patients who have had facial dermatological fillers)

# Positive PIL (Pfizer Vaccine)

**Package leaflet: Information for the user**  
**Comirnaty concentrate for dispersion for injection**  
 COVID-19 mRNA Vaccine (nucleoside modified)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

**Read all of this leaflet carefully before you receive this vaccine because it contains important information for you.**

**What Comirnaty is and what it is used for**

Comirnaty is a vaccine used for preventing COVID-19 caused by SARS-CoV-2 virus.

Comirnaty is given to adults and adolescents from 12 years of age and older.

The vaccine causes the immune system (the body's natural defences) to produce antibodies and blood cells that work against the virus, so giving protection against COVID-19.

As Comirnaty does not contain the virus to produce immunity, it cannot give you COVID-19.

**Possible side effects**

Like all vaccines, Comirnaty can cause side effects, although not everybody gets them.

**The following side effects may occur with Comirnaty:**

**Very common side effects** (9 in 10 or fewer people may *not be affected*)

- injection site: pain, swelling
- tiredness
- headache
- muscle pain
- chills
- joint pain
- diarrhoea
- fever

**Very Common:** 9 in 10 or fewer **not affected** (light to dark blue circles)




Some of these side effects were slightly more frequent in adolescents 12 to 15 years than in adults.

**Common side effects** (90 in 100 or more people may *not be affected*)

- injection site redness
- nausea
- vomiting


**Common:** 90 in 100 (blue circles) or more **not affected**



**Uncommon side effects** (990 in 1,000 or more people may *not be affected*)

- enlarged lymph nodes
- feeling unwell
- arm pain
- insomnia
- injection site itching
- allergic reactions such as rash or itching


**Uncommon:** 990 in 1,000 (blue circles) or more **not affected**



**Rare side effects** (9,990 in 10,000 or more people may *not be affected*)

- temporary one sided facial drooping
- allergic reactions such as hives or swelling of the face

**Rare:** 9,990 in 10,000 (blue circles) or more **not affected**



**Not known** (cannot be estimated from the available data)

- severe allergic reaction
- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain
- extensive swelling of the vaccinated limb
- swelling of the face (swelling of the face may occur in patients who have had facial dermatological fillers)

# Negative PIL (Moderna Vaccine)

## Package leaflet: Information for the user

### Spikevax dispersion for injection COVID-19 mRNA Vaccine (nucleoside modified)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

**Read all of this leaflet carefully before you receive this vaccine because it contains important information for you.**

#### What Spikevax is and what it is used for

Spikevax is a vaccine used to prevent COVID-19 caused by SARS-CoV-2. It is given to adults and children aged 12 years and older. The active substance in Spikevax is mRNA encoding the SARS-CoV-2 Spike protein. The mRNA is embedded in SM-102 lipid nanoparticles.

As Spikevax does not contain the virus, it cannot give you COVID-19.

#### How the vaccine works

Spikevax stimulates the body's natural defences (immune system). The vaccine works by causing the body to produce protection (antibodies) against the virus that causes COVID-19.

#### Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

#### The following side effects may occur with Spikevax:

##### Very common (may affect more than 1 in 10 people)

- swelling/tenderness of the underarm glands
- headache
- nausea
- vomiting
- muscle ache, joint aches, and stiffness
- pain or swelling at the injection site
- feeling very tired
- chills
- fever

##### Common (may affect up to 1 in 10 people)

- rash
- rash, redness, or hives at the injection site (some of which may occur some time after the injection)
- injection)

##### Uncommon (may affect up to 1 in 100 people)

- itchiness at the injection site

##### Rare (may affect up to 1 in 1,000 people)

- temporary one-sided facial drooping (Bell's palsy)
- swelling of the face (Swelling of the face may occur in patients who have had facial cosmetic injections.)
- dizziness
- decreased sense of touch or sensation

##### Frequency unknown (cannot be estimated from the available data)

- severe allergic reactions with breathing difficulties (anaphylaxis)
- reaction of increased sensitivity or intolerance by the immune system (hypersensitivity)
- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain.

# Positive PIL (Moderna Vaccine)

## Package leaflet: Information for the user

### Spikevax dispersion for injection COVID-19 mRNA Vaccine (nucleoside modified)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

Read all of this leaflet carefully before you receive this vaccine because it contains important information for you.

#### What Spikevax is and what it is used for

Spikevax is a vaccine used to prevent COVID-19 caused by SARS-CoV-2. It is given to adults and children aged 12 years and older. The active substance in Spikevax is mRNA encoding the SARS-CoV-2 Spike protein. The mRNA is embedded in SM-102 lipid nanoparticles.

As Spikevax does not contain the virus, it cannot give you COVID-19.

#### How the vaccine works

Spikevax stimulates the body's natural defences (immune system). The vaccine works by causing the body to produce protection (antibodies) against the virus that causes COVID-19.

#### Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

#### The following side effects may occur with Spikevax:

##### Very common (9 in 10 or fewer people may *not be affected*)

- swelling/tenderness of the underarm glands
- headache
- nausea
- vomiting
- muscle ache, joint aches, and stiffness
- pain or swelling at the injection site
- feeling very tired
- chills
- fever


##### Very Common: 9 in 10 or fewer *not affected* (light to dark blue circles)



##### Common (90 in 100 or more people may *not be affected*)

- rash
- rash, redness, or hives at the injection site (some of which may occur some time after the injection)


##### Common: 90 in 100 (blue circles) or more *not affected*



##### Uncommon (990 in 1,000 or more people may *not be affected*)

- itchiness at the injection site


##### Uncommon: 990 in 1,000 (blue circles) or more *not affected*



##### Rare (9,990 in 10,000 or more people may *not be affected*)

- temporary one-sided facial drooping (Bell's palsy)
- swelling of the face (Swelling of the face may occur in patients who have had facial cosmetic injections.)
- dizziness
- decreased sense of touch or sensation

##### Rare: 9,990 in 10,000 (blue circles) or more *not affected*



##### Frequency unknown (cannot be estimated from the available data)

- severe allergic reactions with breathing difficulties (anaphylaxis)
- reaction of increased sensitivity or intolerance by the immune system (hypersensitivity)
- inflammation of the heart muscle (myocarditis) or inflammation of the lining outside the heart (pericarditis) which can result in breathlessness, palpitations or chest pain.

## Supplementary Materials 4

### Demographic Information (Full Sample)

#### Descriptive Statistics: Gender

Question wording:

“Which of the following most accurately describes you?”

Options:

- woman
- man
- non-binary/genderqueer/agender/gender fluid (non-binary)
- not sure
- prefer not to say
- other - let me type

Gender	N
woman	706
man	511
non-binary	3
not sure	1
prefer not to say	1

#### Descriptive Statistics: Education

Question wording:

“Please select your highest level of education:”

Options:

- GNVQ / GSVQ / GCSE/ SCE standard (GCSE)
- NVQ1, NVQ2 (NVQ1/2)
- NVQ3/ SCE Higher Grade/ Advanced GNVQ/ GCE A/AS or similar (A-Levels)
- NVQ4 / HNC / HND / Bachelor’s degree or similar (Bachelors)
- NVQ5 or post-graduate diploma (Post-Graduate)
- Primary/Secondary school (no qualifications) (Primary/Secondary)
- Other (please specify)

Education	N
Bachelors	413
A-Levels	285
GCSE	255
Post-Graduate	121
NVQ1/2	70
Primary / Secondary	53
Other	25

## Descriptive Statistics: Employment

Question wording:

“Which of the following best describes you?”

Options:

- Employed full-time (full-time)
- Employed part-time (part-time)
- Full-time parenting / caring responsibilities (parent / carer)
- Retired
- Self employed (self-employed)
- Student
- Unemployed and not looking for a job / Long-term sick or disabled (unemployed not looking)
- Unemployed but looking for a job (unemployed looking)
- Other (not stated)

Employment	N
full-time	407
retired	321
part-time	153
self-employed	98
unemployed (not looking)	92
parent / carer	70
unemployed (looking)	38
student	24
other	19

## Descriptive Statistics: Ethnicity

Question wording:

“Please select the option that best describes your ethnicity”

Options:

- Asian or Asian British - Bangladeshi
- Asian or Asian British - Indian (A/AB\_I)
- Asian or Asian British - Pakistani (A/AB\_P)
- Asian or Asian British - Any other Asian background (A/AB\_Other)
- Black or Black British - African (B/BB\_A)
- Black or Black British - Caribbean (B/BB\_C)
- Black or Black British - Any other Black background
- Mixed - White and Black Caribbean (M/W&BC)
- Mixed - White and Black African
- Mixed - White and Asian (M/W&A)
- Mixed - Any other Mixed background
- Other ethnic groups - Chinese (OEG\_C)
- Other ethnic groups - Any other ethnic group (ONS)
- White - British (W/B)
- White - Irish (W/I)
- White - Any other White background (W/O)

---

Ethnicity	N
W/B	1064
W/O	58
A/AB_I	19
W/I	16
B/BB_A	11
M/W&BC	10
A/AB_P	9
OEG_C	7
A/AB_Other	5
B/BB_C	5
M/Other	5
A/AB_B	4
M/W&A	4
OEG	3
M/W&BA	1
ONS	1

---

## Supplementary Materials 5

### Demographic Information (by Experimental Condition)

#### Gender\*Condition

	man	non-binary	not sure	prefer not to say	woman
-ve familiar	79	1	0	0	123
-ve same	81	0	0	0	122
-ve unfamiliar	87	1	0	1	116
+ve familiar	93	1	1	0	106
+ve same	90	0	0	0	117
+ve unfamiliar	81	0	0	0	122

#### Education\*Condition

	GCSE	NVQ1/2	A-Levels	Bachelors	Post-Graduate	Other	Primary / Secondary
-ve familiar	48	13	46	61	21	7	7
-ve same	44	10	50	66	14	4	15
-ve unfamiliar	44	11	47	71	27	2	3
+ve familiar	39	10	44	76	17	5	10
+ve same	36	16	53	65	21	5	11
+ve unfamiliar	44	10	45	74	21	2	7

#### Employment\*Condition

	full-time	part-time	parent / carer	other	retired	self-employed	student	unemployed (not looking)	unemployed (looking)
-ve familiar	61	36	10	1	47	17	6	16	9
-ve same	63	28	13	2	52	19	4	16	6
-ve unfamiliar	73	29	10	2	46	21	5	15	4
+ve familiar	70	21	10	7	58	14	2	14	5
+ve same	68	16	10	3	66	16	3	17	8
+ve unfamiliar	72	23	17	4	52	11	4	14	6

### Ethnicity\*Condition

Categories are:

- 1 = "A/AB\_Other"
- 2 = "A/AB\_B"
- 3 = "A/AB\_I"
- 4 = "A/AB\_P"
- 5 = "B/BB\_A"
- 6 = "B/BB\_C"
- 7 = "M/Other"
- 8 = "M/W&A"
- 9 = "M/W&BA"
- 10 = "M/W&BC"
- 11 = "ONS"
- 12 = "OEG"
- 13 = "OEG\_C"
- 14 = "W/O"
- 15 = "W/B"
- 16 = "W/I"

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
-ve familiar	0	1	1	1	1	0	1	0	1	2	0	0	0	8	183	4
-ve same	2	2	3	1	3	1	0	2	0	0	0	0	2	10	174	3
-ve unfamiliar	1	0	4	3	2	0	2	0	0	2	0	0	3	14	170	4
+ve familiar	0	1	3	1	1	2	1	0	0	4	0	0	1	9	177	1
+ve same	0	0	5	0	3	2	1	2	0	1	1	2	0	8	181	1
+ve unfamiliar	2	0	3	3	1	0	0	0	0	1	0	1	1	9	179	3

### Months Since Last COVID Vaccine\*Condition

	0	1	2	3	4	5	6	7	8	9	10	11	13
-ve familiar	0	4	13	31	34	68	41	7	4	0	0	0	1
-ve same	0	6	14	24	30	65	53	3	8	0	0	0	0
-ve unfamiliar	0	6	11	25	25	66	50	10	6	3	2	1	0
+ve familiar	1	8	7	22	29	56	62	15	1	0	0	0	0
+ve same	0	1	14	26	25	65	66	9	1	0	0	0	0
+ve unfamiliar	0	2	10	22	36	71	46	10	5	1	0	0	0

### Side Effects from Previous COVID Vaccines\*Condition

	first dose	second dose	both doses	no side effects
-ve familiar	61	28	21	93
-ve same	61	26	17	99
-ve unfamiliar	62	24	26	93
+ve familiar	57	20	22	102
+ve same	65	23	23	96
+ve unfamiliar	66	20	21	96

## Supplementary Materials 6

Full statistical model for the primary outcome (including interactions between the factors (Framing/Familiarity) and the planned covariate (Baseline Booster Intention))

Variables are:

- intent\_fv\_t2 (primary outcome: booster intention (intent) for the framed vaccine (fv) post-manipulation (t2))
- intent\_fv\_t1 (covariate: booster intention (intent) for the framed vaccine (fv) at baseline (t1))

Full model accounting for interactions with the covariate

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	48462.8455	1	173.6635	0.0000
intent_fv_t1	543030.3892	1	1945.9146	0.0000
Framing	25.7614	1	0.0923	0.7613
Familiarity	4466.3893	2	8.0025	0.0004
intent_fv_t1:Framing	76.6206	1	0.2746	0.6004
intent_fv_t1:Familiarity	5867.6007	2	10.5131	0.0000
Framing:Familiarity	6002.1555	2	10.7542	0.0000
intent_fv_t1:Framing:Familiarity	4268.1448	2	7.6473	0.0005
Residuals	337664.7592	1210	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5321	0.6166
Framing	0.0000	0.0001
Familiarity	0.0044	0.0131
intent_fv_t1:Framing	0.0001	0.0002
intent_fv_t1:Familiarity	0.0057	0.0171
Framing:Familiarity	0.0059	0.0175
intent_fv_t1:Framing:Familiarity	0.0042	0.0125

**Full model with pre-registered contrasts for Vaccine Familiarity**

- cont1 = Same Vaccine vs. Other Vaccines [Familiar and Unfamiliar combined]
- cont2 = Familiar Vaccine vs. Unfamiliar Vaccine

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	48462.8455	1	173.6635	0.0000
intent_fv_t1	543030.3892	1	1945.9146	0.0000
Framing	25.7614	1	0.0923	0.7613
cont1	3134.4070	1	11.2319	0.0008
cont2	1335.9564	1	4.7873	0.0289
intent_fv_t1:Framing	76.6206	1	0.2746	0.6004
intent_fv_t1:cont1	4420.8055	1	15.8417	0.0001
Framing:cont1	1413.4722	1	5.0651	0.0246
intent_fv_t1:cont2	1553.5052	1	5.5669	0.0185
Framing:cont2	4593.6316	1	16.4610	0.0001
intent_fv_t1:Framing:cont1	1225.7388	1	4.3924	0.0363
intent_fv_t1:Framing:cont2	3122.1111	1	11.1879	0.0008
Residuals	337664.7592	1210	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5321	0.6166
Framing	0.0000	0.0001
cont1	0.0031	0.0092
cont2	0.0013	0.0039
intent_fv_t1:Framing	0.0001	0.0002
intent_fv_t1:cont1	0.0043	0.0129
Framing:cont1	0.0014	0.0042
intent_fv_t1:cont2	0.0015	0.0046
Framing:cont2	0.0045	0.0134
intent_fv_t1:Framing:cont1	0.0012	0.0036
intent_fv_t1:Framing:cont2	0.0031	0.0092

## Supplementary Materials 7

### Pre-registered ANCOVA (without interactions between factors and covariate)

The following presents the results from the pre-registered ANCOVA that did not account for interactions between the covariate (baseline booster intention) and the Framing/Familiarity factors.

#### Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	81587.3803	1	285.3356	0.0000
intent_fv_t1	607341.4178	1	2124.0559	0.0000
Framing	667.2802	1	2.3337	0.1269
Familiarity	820.2857	2	1.4344	0.2387
Framing:Familiarity	1987.1180	2	3.4748	0.0313
Residuals	347410.7352	1215	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5951	0.6361
Framing	0.0007	0.0019
Familiarity	0.0008	0.0024
Framing:Familiarity	0.0019	0.0057

#### Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	81587.3803	1	285.3356	0.0000
intent_fv_t1	607341.4178	1	2124.0559	0.0000
Framing	667.2802	1	2.3337	0.1269
cont1	801.2642	1	2.8023	0.0944
cont2	16.7860	1	0.0587	0.8086
Framing:cont1	236.0587	1	0.8256	0.3637
Framing:cont2	1747.3457	1	6.1110	0.0136
Residuals	347410.7352	1215	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5951	0.6361
Framing	0.0007	0.0019
cont1	0.0008	0.0023
cont2	0.0000	0.0000
Framing:cont1	0.0002	0.0007
Framing:cont2	0.0017	0.0050

### Framing\*Familiarity Means

Framing	Familiarity	emmean	SE	df	lower.CL	upper.CL
Negative Frame	Unfamiliar	77.0667	1.1868	1215	74.7383	79.3951
Positive Frame	Unfamiliar	82.0994	1.1939	1215	79.7572	84.4417
Negative Frame	Familiar	79.7128	1.1885	1215	77.3810	82.0445
Positive Frame	Familiar	78.8776	1.1952	1215	76.5327	81.2226
Negative Frame	Same	81.1027	1.1989	1215	78.7505	83.4549
Positive Frame	Same	81.3391	1.1946	1215	78.9955	83.6827

## Supplementary Materials 8

The effect of positive framing on vaccine switches

Full model including previous vaccine type (“prevax\_type”)

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	48152.2239	1	173.6703	0.0000
intent_fv_t1	466284.3106	1	1681.7448	0.0000
Framing	68.7237	1	0.2479	0.6187
Familiarity	4162.4448	2	7.5063	0.0006
prevax_type	1763.3901	1	6.3600	0.0118
intent_fv_t1:Framing	143.0262	1	0.5159	0.4728
intent_fv_t1:Familiarity	5531.5062	2	9.9752	0.0001
Framing:Familiarity	7062.0571	2	12.7353	0.0000
intent_fv_t1:prevax_type	945.5318	1	3.4102	0.0650
Framing:prevax_type	7.3229	1	0.0264	0.8709
Familiarity:prevax_type	207.0593	2	0.3734	0.6885
intent_fv_t1:Framing:Familiarity	5204.8395	2	9.3861	0.0001
intent_fv_t1:Framing:prevax_type	0.3171	1	0.0011	0.9730
intent_fv_t1:Familiarity:prevax_type	317.3683	2	0.5723	0.5644
Framing:Familiarity:prevax_type	1611.3401	2	2.9058	0.0551
intent_fv_t1:Framing:Familiarity:prevax_type	930.3705	2	1.6778	0.1872
Residuals	332160.1508	1198	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.4569	0.5840
Framing	0.0001	0.0002
Familiarity	0.0041	0.0124
prevax_type	0.0017	0.0053
intent_fv_t1:Framing	0.0001	0.0004
intent_fv_t1:Familiarity	0.0054	0.0164
Framing:Familiarity	0.0069	0.0208
intent_fv_t1:prevax_type	0.0009	0.0028
Framing:prevax_type	0.0000	0.0000
Familiarity:prevax_type	0.0002	0.0006
intent_fv_t1:Framing:Familiarity	0.0051	0.0154
intent_fv_t1:Framing:prevax_type	0.0000	0.0000
intent_fv_t1:Familiarity:prevax_type	0.0003	0.0010
Framing:Familiarity:prevax_type	0.0016	0.0048
intent_fv_t1:Framing:Familiarity:prevax_type	0.0009	0.0028

## Vaccine Switch: AstraZeneca to Pfizer and Moderna

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	34492.1950	1	105.8593	0.0000
intent_fv_t1	139372.9781	1	427.7470	0.0000
Framing	11.5190	1	0.0354	0.8510
Familiarity	833.9037	1	2.5593	0.1104
intent_fv_t1:Framing	7.8985	1	0.0242	0.8764
intent_fv_t1:Familiarity	1172.9565	1	3.5999	0.0585
Framing:Familiarity	4799.7488	1	14.7308	0.0001
intent_fv_t1:Framing:Familiarity	3707.3265	1	11.3781	0.0008
Residuals	130983.8324	402	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5019	0.5155
Framing	0.0000	0.0001
Familiarity	0.0030	0.0063
intent_fv_t1:Framing	0.0000	0.0001
intent_fv_t1:Familiarity	0.0042	0.0089
Framing:Familiarity	0.0173	0.0353
intent_fv_t1:Framing:Familiarity	0.0134	0.0275

## Vaccine Switch: Pfizer to Moderna

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	26187.1494	1	59.7228	0.0000
intent_fv_t1	111498.8315	1	254.2860	0.0000
Framing	1744.2603	1	3.9780	0.0475
intent_fv_t1:Framing	791.0549	1	1.8041	0.1808
Residuals	86818.6664	198	NA	NA

	eta.sq	eta.sq.part
intent_fv_t1	0.5565	0.5622
Framing	0.0087	0.0197
intent_fv_t1:Framing	0.0039	0.0090

## Supplementary Materials 9

### Secondary Predictor: Booster Side Effect Severity

- `sesev_fv_t2` (perceived side effect severity associated with the framed vaccine post-manipulation)
- `sesev_fv_t1` (perceived side effect severity associated with the framed vaccine at baseline)

### Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	57189.5779	1	183.2956	0.0000
<code>sesev_fv_t1</code>	263557.1698	1	844.7145	0.0000
Framing	1222.8893	1	3.9194	0.0480
Familiarity	40.5057	2	0.0649	0.9372
<code>sesev_fv_t1:Framing</code>	1411.1856	1	4.5229	0.0336
<code>sesev_fv_t1:Familiarity</code>	654.8262	2	1.0494	0.3505
Framing:Familiarity	157.2217	2	0.2520	0.7773
<code>sesev_fv_t1:Framing:Familiarity</code>	314.8565	2	0.5046	0.6039
Residuals	377528.9467	1210	NA	NA

	eta.sq	eta.sq.part
<code>sesev_fv_t1</code>	0.3935	0.4111
Framing	0.0018	0.0032
Familiarity	0.0001	0.0001
<code>sesev_fv_t1:Framing</code>	0.0021	0.0037
<code>sesev_fv_t1:Familiarity</code>	0.0010	0.0017
Framing:Familiarity	0.0002	0.0004
<code>sesev_fv_t1:Framing:Familiarity</code>	0.0005	0.0008

Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	57189.5779	1	183.2956	0.0000
sesev_fv_t1	263557.1698	1	844.7145	0.0000
Framing	1222.8893	1	3.9194	0.0480
cont1	40.0779	1	0.1285	0.7201
cont2	0.5846	1	0.0019	0.9655
sesev_fv_t1:Framing	1411.1856	1	4.5229	0.0336
sesev_fv_t1:cont1	609.9385	1	1.9549	0.1623
Framing:cont1	53.6660	1	0.1720	0.6784
sesev_fv_t1:cont2	51.2097	1	0.1641	0.6855
Framing:cont2	100.9352	1	0.3235	0.5696
sesev_fv_t1:Framing:cont1	156.2693	1	0.5009	0.4793
sesev_fv_t1:Framing:cont2	152.7560	1	0.4896	0.4842
Residuals	377528.9467	1210	NA	NA

	eta.sq	eta.sq.part
sesev_fv_t1	0.3935	0.4111
Framing	0.0018	0.0032
cont1	0.0001	0.0001
cont2	0.0000	0.0000
sesev_fv_t1:Framing	0.0021	0.0037
sesev_fv_t1:cont1	0.0009	0.0016
Framing:cont1	0.0001	0.0001
sesev_fv_t1:cont2	0.0001	0.0001
Framing:cont2	0.0002	0.0003
sesev_fv_t1:Framing:cont1	0.0002	0.0004
sesev_fv_t1:Framing:cont2	0.0002	0.0004

# Supplementary Materials 10

## Secondary Predictor: Perceived Booster Risk

- serisk\_fv\_t2 (perceived risk associated with the framed vaccine post-manipulation)
- sesev\_fv\_t1 (perceived risk associated with the framed vaccine at baseline)

## Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	81175.1304	1	193.8057	0.0000
serisk_fv_t1	215659.8866	1	514.8883	0.0000
Framing	436.2962	1	1.0417	0.3076
Familiarity	894.8706	2	1.0683	0.3439
serisk_fv_t1:Framing	21.8767	1	0.0522	0.8193
serisk_fv_t1:Familiarity	4039.4547	2	4.8221	0.0082
Framing:Familiarity	74.8987	2	0.0894	0.9145
serisk_fv_t1:Framing:Familiarity	919.5879	2	1.0978	0.3340
Residuals	506805.9582	1210	NA	NA

	eta.sq	eta.sq.part
serisk_fv_t1	0.2885	0.2985
Framing	0.0006	0.0009
Familiarity	0.0012	0.0018
serisk_fv_t1:Framing	0.0000	0.0000
serisk_fv_t1:Familiarity	0.0054	0.0079
Framing:Familiarity	0.0001	0.0001
serisk_fv_t1:Framing:Familiarity	0.0012	0.0018

Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	81175.1304	1	193.8057	0.0000
serisk_fv_t1	215659.8866	1	514.8883	0.0000
Framing	436.2962	1	1.0417	0.3076
cont1	109.4773	1	0.2614	0.6093
cont2	754.8287	1	1.8022	0.1797
serisk_fv_t1:Framing	21.8767	1	0.0522	0.8193
serisk_fv_t1:cont1	1791.4699	1	4.2771	0.0388
Framing:cont1	11.7394	1	0.0280	0.8671
serisk_fv_t1:cont2	2526.0142	1	6.0309	0.0142
Framing:cont2	65.7193	1	0.1569	0.6921
serisk_fv_t1:Framing:cont1	708.3207	1	1.6911	0.1937
serisk_fv_t1:Framing:cont2	267.7285	1	0.6392	0.4242
Residuals	506805.9582	1210	NA	NA

	eta.sq	eta.sq.part
serisk_fv_t1	0.2885	0.2985
Framing	0.0006	0.0009
cont1	0.0001	0.0002
cont2	0.0010	0.0015
serisk_fv_t1:Framing	0.0000	0.0000
serisk_fv_t1:cont1	0.0024	0.0035
Framing:cont1	0.0000	0.0000
serisk_fv_t1:cont2	0.0034	0.0050
Framing:cont2	0.0001	0.0001
serisk_fv_t1:Framing:cont1	0.0009	0.0014
serisk_fv_t1:Framing:cont2	0.0004	0.0005

# Supplementary Materials 11

## Correlations and PCA for Framed Vaccine Side Effect Perceptions

### (Satisfaction, Happiness and Anxiety)

**Correlations between variables** Variables are: switchsat\_fv\_t2 (satisfaction with the framed vaccine post-manipulation) switchhap\_fv\_t2 (happiness with the framed vaccine post-manipulation) switchanx\_rev\_fv\_t2 (anxiety with the framed vaccine post-manipulation - variable reversed (+ve association))

```
##
## Pearson's product-moment correlation
##
## data:  sat_dat$switchsat_fv_t2 and sat_dat$switchhap_fv_t2
## t = 78.489, df = 1220, p-value < 2.2e-16
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
##  0.9038498 0.9224377
## sample estimates:
##      cor
## 0.9136199

##
## Pearson's product-moment correlation
##
## data:  sat_dat$switchsat_fv_t2 and sat_dat$switchanx_rev_fv_t2
## t = 16.839, df = 1220, p-value < 2.2e-16
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
##  0.3876229 0.4786818
## sample estimates:
##      cor
## 0.4342611

##
## Pearson's product-moment correlation
##
## data:  sat_dat$switchhap_fv_t2 and sat_dat$switchanx_rev_fv_t2
## t = 17.55, df = 1220, p-value < 2.2e-16
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
##  0.4030299 0.4926355
## sample estimates:
##      cor
## 0.4489607
```

**PCA component loadings for post-manipulation measures**

	PC1	PC2	PC3
switchsat_fv_t2	0.6135009	-0.3383353	-0.7135446
switchhap_fv_t2	0.6338972	-0.3278683	0.7004832
switchanx_rev_fv_t2	0.4709469	0.8820610	-0.0133223

**PCA component loadings for pre-manipulation measures**

	PC1	PC2	PC3
switchsat_fv_t1	0.6469202	-0.2877156	0.7061969
switchhap_fv_t1	0.6539002	-0.2671390	-0.7078498
switchanx_rev_fv_t1	0.3923122	0.9197045	0.0153197

**Secondary Predictor: Satisfaction and Happiness combined**

- av\_sathap\_t2 (average satisfaction and happiness associated with the framed vaccine post-manipulation)
- av\_sathap\_t1 (average satisfaction and happiness associated with the framed vaccine at baseline)

**Main Effects Model**

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	68464.8110	1	177.3429	0.0000
av_sathap_t1	629741.7591	1	1631.2060	0.0000
Framing	2184.7699	1	5.6592	0.0175
Familiarity	1042.8219	2	1.3506	0.2595
av_sathap_t1:Framing	849.7667	1	2.2011	0.1382
av_sathap_t1:Familiarity	1420.4558	2	1.8397	0.1593
Framing:Familiarity	2803.5778	2	3.6310	0.0268
av_sathap_t1:Framing:Familiarity	2651.9911	2	3.4347	0.0325
Residuals	467131.3843	1210	NA	NA

	eta.sq	eta.sq.part
av_sathap_t1	0.5173	0.5741
Framing	0.0018	0.0047
Familiarity	0.0009	0.0022
av_sathap_t1:Framing	0.0007	0.0018
av_sathap_t1:Familiarity	0.0012	0.0030
Framing:Familiarity	0.0023	0.0060
av_sathap_t1:Framing:Familiarity	0.0022	0.0056

**Model with pre-registered contrasts for Vaccine Familiarity**

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	68464.8110	1	177.3429	0.0000
av_sathap_t1	629741.7591	1	1631.2060	0.0000
Framing	2184.7699	1	5.6592	0.0175
cont1	89.8365	1	0.2327	0.6296
cont2	949.0046	1	2.4582	0.1172
av_sathap_t1:Framing	849.7667	1	2.2011	0.1382
av_sathap_t1:cont1	7.8678	1	0.0204	0.8865
Framing:cont1	1323.4860	1	3.4282	0.0643
av_sathap_t1:cont2	1405.2058	1	3.6399	0.0566
Framing:cont2	1461.2307	1	3.7850	0.0519
av_sathap_t1:Framing:cont1	1588.5089	1	4.1147	0.0427
av_sathap_t1:Framing:cont2	987.7901	1	2.5587	0.1100
Residuals	467131.3843	1210	NA	NA

	eta.sq	eta.sq.part
av_sathap_t1	0.5173	0.5741
Framing	0.0018	0.0047
cont1	0.0001	0.0002
cont2	0.0008	0.0020
av_sathap_t1:Framing	0.0007	0.0018
av_sathap_t1:cont1	0.0000	0.0000
Framing:cont1	0.0011	0.0028
av_sathap_t1:cont2	0.0012	0.0030
Framing:cont2	0.0012	0.0031
av_sathap_t1:Framing:cont1	0.0013	0.0034
av_sathap_t1:Framing:cont2	0.0008	0.0021

**Secondary Predictor: Anxiety**

**Main Effects Model**

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	80477.3132	1	140.8544	0.0000
switchanx_fv_t1	584744.8174	1	1023.4421	0.0000
Framing	1449.0382	1	2.5362	0.1115
Familiarity	1743.2143	2	1.5255	0.2179
switchanx_fv_t1:Framing	3160.2594	1	5.5312	0.0188
switchanx_fv_t1:Familiarity	524.9404	2	0.4594	0.6318
Framing:Familiarity	318.5277	2	0.2787	0.7568
switchanx_fv_t1:Framing:Familiarity	2257.7873	2	1.9758	0.1391
Residuals	691334.8598	1210	NA	NA

	eta.sq	eta.sq.part
switchanx_fv_t1	0.4358	0.4582

	eta.sq	eta.sq.part
Framing	0.0011	0.0021
Familiarity	0.0013	0.0025
switchanx_fv_t1:Framing	0.0024	0.0046
switchanx_fv_t1:Familiarity	0.0004	0.0008
Framing:Familiarity	0.0002	0.0005
switchanx_fv_t1:Framing:Familiarity	0.0017	0.0033

### Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	80477.3132	1	140.8544	0.0000
switchanx_fv_t1	584744.8174	1	1023.4421	0.0000
Framing	1449.0382	1	2.5362	0.1115
cont1	1041.1486	1	1.8223	0.1773
cont2	788.8024	1	1.3806	0.2402
switchanx_fv_t1:Framing	3160.2594	1	5.5312	0.0188
switchanx_fv_t1:cont1	4.6664	1	0.0082	0.9280
Framing:cont1	174.0626	1	0.3047	0.5811
switchanx_fv_t1:cont2	518.5455	1	0.9076	0.3409
Framing:cont2	160.4686	1	0.2809	0.5962
switchanx_fv_t1:Framing:cont1	2086.2670	1	3.6515	0.0563
switchanx_fv_t1:Framing:cont2	152.6742	1	0.2672	0.6053
Residuals	691334.8598	1210	NA	NA

	eta.sq	eta.sq.part
switchanx_fv_t1	0.4358	0.4582
Framing	0.0011	0.0021
cont1	0.0008	0.0015
cont2	0.0006	0.0011
switchanx_fv_t1:Framing	0.0024	0.0046
switchanx_fv_t1:cont1	0.0000	0.0000
Framing:cont1	0.0001	0.0003
switchanx_fv_t1:cont2	0.0004	0.0007
Framing:cont2	0.0001	0.0002
switchanx_fv_t1:Framing:cont1	0.0016	0.0030
switchanx_fv_t1:Framing:cont2	0.0001	0.0002

## Supplementary Materials 12

### Positive Framing Influences Judgement Regarding Side Effect Prevalence

#### Side effects presented from each PIL

The following presents the 14 side effects associated with each vaccine type (i.e., AstraZeneca, Pfizer, Moderna). For each of the side effects, participants were required to: 1) categorise into risk brackets (very common, common, uncommon, rare, very rare) and, 2) rate based on their perception of risk (out of 100,000 people how many would experience the side effect). As side effects are identical to those included in the manufacturer PILs, risk categories differ across vaccine types. As such, metrics derived for analysis of risk categorisation involved a single value associated with the percentage of correct assignments across side effects. This was performed for the 11 side effects that have a discrete risk bracket. Within each group of 14 side effects, 3 were categorised as of ‘unknown risk’ based on RCT data. Frequency estimates of these items were analysed separately in a sub analysis to determine whether positive framing results in generally reduced risk perceptions when no concrete information regarding risk is provided.

### AstraZeneca Side Effects

<i>Category</i>	<i>Side Effect (N = 11)</i>
Very Common	Injection site pain
Very Common	Tiredness (fatigue)
Very Common	Muscle ache/pain
Very Common	Chills
Common	Injection site redness
Common	Vomiting
Uncommon	Diarrhea
Uncommon	Enlarged lymph nodes (swelling of underarm glands)
Uncommon	Dizziness
Very Rare	Blood clots often in unusual locations (e.g. brain, bowel, liver, spleen) in combination with low level of blood platelets
Very Rare	Serious nerve inflammation, which may cause paralysis and difficulty breathing (Guillain-Barré syndrome)

### AstraZeneca Unknown Side Effects

<i>Category</i>	<i>Side Effect (N = 3)</i>
Not known	Severe allergic reaction (anaphylaxis)
Not known	hypersensitivity
Not known	capillary leak syndrome (a condition causing fluid leakage from small blood vessels)

### Pfizer Side Effects

Category	Side Effect (N = 11)
Very Common	Injection site pain
Very Common	Tiredness (fatigue)
Very Common	Muscle ache/pain
Very Common	Chills
Common	Injection site redness
Common	Vomiting
Uncommon	Diarrhea
Uncommon	Enlarged lymph nodes (swelling of underarm glands)
Uncommon	Insomnia
Rare	Temporary one-sided facial drooping
Rare	Swelling of the face

### Pfizer Unknown Side Effects

Category	Side Effect (N = 3)
Not known	Severe allergic reaction (anaphylaxis)
Not known	Inflammation of the heart (myocarditis or pericarditis)
Not known	Extensive swelling of the vaccinated limb

Moderna Side Effects	
Category	Side Effect (N = 11)
Very Common	Injection site pain
Very Common	Tiredness (fatigue)
Very Common	Muscle ache/pain
Very Common	Chills
Common	Injection site redness
Common	Rash
Common	Injection site hives
Uncommon	injection site itching
Rare	Temporary one-sided facial drooping
Rare	Swelling of the face
Rare	Decreased sense of touch or sensation

Moderna Unknown Side Effects	
Category	Side Effect (N = 3)
Not known	Severe allergic reaction (anaphylaxis)
Not known	Hypersensitivity
Not known	Inflammation of the heart (myocarditis or pericarditis)

## Supplementary Materials 13

### Secondary Predictor: Verbal Prevalence Categories

Overall\_CategoryPercent = percentage of correctly identified categories

#### Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	1258379.599	1	3900.3990	0.0000
Framing	1786.274	1	5.5366	0.0188
Familiarity	1923.618	2	2.9812	0.0511
Framing:Familiarity	1047.111	2	1.6228	0.1978
Residuals	392316.166	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0045	0.0045
Familiarity	0.0048	0.0049
Framing:Familiarity	0.0026	0.0027

#### Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	1258379.5992	1	3900.3990	0.0000
Framing	1786.2735	1	5.5366	0.0188
cont1	1590.3548	1	4.9294	0.0266
cont2	329.1191	1	1.0201	0.3127
Framing:cont1	1003.2909	1	3.1097	0.0781
Framing:cont2	42.6316	1	0.1321	0.7163
Residuals	392316.1662	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0045	0.0045
cont1	0.0040	0.0040
cont2	0.0008	0.0008
Framing:cont1	0.0025	0.0026
Framing:cont2	0.0001	0.0001

## Supplementary Materials 14

### Secondary Predictor: Frequency Estimates

Overall\_FrequencyPercent = percentage of correctly estimated frequency brackets

#### Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	602115.9981	1	1777.6846	0.0000
Framing	2185.5526	1	6.4526	0.0112
Familiarity	6577.8779	2	9.7102	0.0001
Framing:Familiarity	340.6265	2	0.5028	0.6049
Residuals	411868.9225	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0052	0.0053
Familiarity	0.0156	0.0157
Framing:Familiarity	0.0008	0.0008

#### Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	602115.9981	1	1777.6846	0.0000
Framing	2185.5526	1	6.4526	0.0112
cont1	3225.6804	1	9.5235	0.0021
cont2	3333.4302	1	9.8416	0.0017
Framing:cont1	1.8231	1	0.0054	0.9415
Framing:cont2	338.6588	1	0.9999	0.3175
Residuals	411868.9225	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0052	0.0053
cont1	0.0077	0.0078
cont2	0.0079	0.0080
Framing:cont1	0.0000	0.0000
Framing:cont2	0.0008	0.0008

## Supplementary Materials 15

Frequency estimates for side effect of ‘unknown prevalence’

Main Effects Model

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	4533390495	1	29.5304	0.0000
Framing	7457785	1	0.0486	0.8256
Familiarity	537592293	2	1.7509	0.1740
Framing:Familiarity	119664905	2	0.3897	0.6773
Residuals	186675244185	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0000	0.0000
Familiarity	0.0029	0.0029
Framing:Familiarity	0.0006	0.0006

Model with pre-registered contrasts for Vaccine Familiarity

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	4.533390e+09	1	29.5304	0.0000
Framing	7.457785e+06	1	0.0486	0.8256
cont1	2.443971e+08	1	1.5920	0.2073
cont2	2.947225e+08	1	1.9198	0.1661
Framing:cont1	5.626220e+04	1	0.0004	0.9847
Framing:cont2	1.195929e+08	1	0.7790	0.3776
Residuals	1.866752e+11	1216	NA	NA

	eta.sq	eta.sq.part
Framing	0.0000	0.0000
cont1	0.0013	0.0013
cont2	0.0016	0.0016
Framing:cont1	0.0000	0.0000
Framing:cont2	0.0006	0.0006


Mean estimates \* Framing

	emmean	SE	df	lower.CL	upper.CL
Negative Frame	2004.294	501.2572	1216	1020.8692	2987.719
Positive Frame	1848.045	501.2893	1216	864.5571	2831.533


# Supplementary Materials 16

Post-hoc Familiarity \* Framing \* Baseline interactions among those with scores above and below the mean of baseline booster intention

## Side Effect Severity



### Booster Acceptance: Satisfaction and Happiness



### Booster Acceptance: Anxiety

