

Supplemental Figures:

Supplemental Figure 1: Dose response curves in response to WT and patient-encoded variants.

(A) EC50 comparison between IFN- λ 2 and IFN- λ 2Var from dose response curve described in Figure 2A.

(B) Dose-response curve of downstream ISG induction following stimulation with either IFN- λ 3 or IFN- λ 3Var1. Normalized relative luciferase units (RLU) shown.

(C) EC50 comparison between IFN- λ 3 and IFN- λ 3Var1. EC50 not quantifiable for IFN- λ 3Var1.

(D) Dose-response curve of downstream ISG induction following stimulation with either IFN- λ 3 or IFN- λ 3Var2. Normalized relative luciferase units (RLU) shown.

(E) EC50 comparison between IFN- λ 3 and IFN- λ 3Var2.

(F) Induction of ISGs following stimulation of healthy control tHIOs with either WT IFN- λ 2 (yellow dots), patient-encoded IFN- λ 2Var (peach dots), or no treatment (black dots). HP1 = Healthy parent 1; HP2 = healthy parent 2; SI = small intestinal enteroids of unrelated healthy control

(G) EC50 comparison between INF- λ 2 and INF- λ 2Var in dimerization assay with IFNLR1 and IL10RB

(H) EC50 comparison between INF- λ 3 and INF- λ 3Var1 in dimerization assay with IFNLR1 and IL10RB

(B-E, G, H) Data depicts three independent experiments, each data point is an independent experiment. RLU were normalized to blank wells (no protein). (C,E,G,H) Statistical significance determined by unpaired t-test between the groups.

Supplemental Figure 1

