

Supplementary Materials

Table of Contents

Supplementary Table 1.....	page 2
Supplementary Table 2.....	page 3
Supplementary Table 3.....	page 4

Supplementary Table 1. Participant demographics by number of follow-up years

		Maximum years of follow-up					Study Cohort
	Missing	Baseline only	1 year	2 years	3 years	4 or more years	
		n=96	n=44	n=40	n=40	n=89	n=213
Female, no (%)	0	25 (26)	5 (11)	9(23)	8 (20)	19 (21)	41 (19)
Age (yrs), mean (SD)	0	56.4 (13)	59.9 (12)	53.4 (14)	58.8 (12)	56.2 (13)	56.9 (13)
Black, no (%)	0	28 (29)	12 (27)	8 (20)	6 (15)	20 (22)	46 (22)
Smoking, no (%)	5	26 (27)	10 (23)	9 (23)	4 (10)	11 (13)	34 (16)
Education, no (%)	25						
Some Highschool		7 (8)	3 (7)	1 (3)	1 (3)	3 (4)	8 (4)
Completed Highschool		69 (69)	27 (68)	21 (60)	26 (68)	53 (62)	127 (64)
Completed College		20 (23)	10 (27)	13 (37)	11 (29)	29 (34)	63 (32)
BMI, mean (SD)	0	31.1 (8)	28.5 (6.4)	29.5 (5.7)	31.8 (7.3)	30.3 (6)	30.1 (6.3)
SBP, mmHg mean (SD)	0	138 (24)	132 (18)	129 (22)	130 (17)	128 (19)	130 (19)
DM, no (%)	0	48 (50)	18 (37)	19 (48)	17 (43)	37 (45)	91 (43)
CAD, no (%)	0	37 (38)	12 (24)	10 (25)	10 (25)	17 (20)	49 (23)
eGFRcysc, mean(SD)	0	40.1 (23)	41.9 (18)	42.3 (21)	45.7 (18)	54.4 (21)	47.9 (21)
eGFRcr-cysc, mean (SD)	0	40.2 (26)	40.8 (18)	40.9 (22)	45.4 (19)	52.2 (23)	46.8 (22)
Hemoglobin (gm/dL), mean (SD)	20	12.6 (1.9)	13 (2)	12.8 (2)	12.9 (2)	13.3 (2)	13.1 (2)
Phosphorus (mg/dL), mean (SD)	0	4 (0.9)	3.8 (0.7)	3.9 (0.7)	3.7 (0.6)	3.4 (0.6)	3.6 (0.7)
Albuminuria (mg/dL) median (IQR)	5	217 (23, 1265)	106 (22, 929)	324 (68, 1011)	114 (13, 696)	45 (15, 689)	105 (10.8, 594.3)
Frail, no (%)	57	11 (13)	6 (17)	3 (10)	2 (6)	3 (4)	14 (8)
Gait speed (m/s), mean (SD)	0	0.94 (0.21)	0.93 (0.20)	0.97 (0.21)	0.97 (0.25)	0.98 (0.18)	0.97 (0.2)

HS=high school, BMI=body mass index, SBP=systolic blood pressure, DM=diabetes mellitus, CAD=coronary artery disease, GFRcysc=Cystatin C based eGFR, GFRcr-cysc=Cystatin C based eGFR with creatinine, PTH=parathyroid hormone. Frailty was defined using slight modifications of criteria originally established by Fried et al.^{1,2}

Supplementary Table 2. Associations with annual changes in gait speed using eGFR with combined cystatin and creatinine

		Model 1	Model 2	Sensitivity Analysis
	Mean (SD)	Adjusted annual change (95% CI)	Adjusted annual change (95% CI)	Adjusted annual change (95% CI)
eGFRcr-cysc				
≥60 (n=52)	1.02 (0.22)	Reference	Reference	Reference
40-59 (n=75)	0.96 (0.20)	-0.016 (-0.039, 0.008)	-0.016 (-0.039, 0.007)	-0.012 (-0.033, 0.009)
30-39 (n=38)	0.89 (0.17)	-0.023 (-0.051, 0.005)	-0.032 (-0.063, -0.002)	-0.016 (-0.054, -0.022)
<30 (n=48)	0.98 (0.20)	-0.044 (-0.077, -0.011)	-0.044 (-0.074, -0.014)	-0.034 (-0.065, -0.003)
Per 30 ml/min per 1.73m²				
eGFRcr-cysc		-0.016 (-0.031, -0.002)	-0.019 (-0.032, -0.006)	-0.016 (-0.029, -0.003)
Age (per 1 year)		0.001 (-0.000, 0.002)	0.001 (-0.000, 0.002)	0.001 (-0.000, 0.002)
Sex (female)		0.004 (-0.033, 0.042)	0.003 (-0.032, 0.039)	0.009 (-0.025, 0.044)
Height (per 1 inch)		-0.000 (-0.004, -0.003)	-0.001 (-0.004, 0.003)	-0.002 (-0.005, 0.002)
Weight (per 10 lbs)		-0.002 (-0.004, 0.000)	-0.001 (-0.003, 0.000)	-0.002 (-0.003, 0.000)
African American race		-0.011 (-0.029, 0.008)	-0.015 (-0.033, 0.003)	-0.012 (-0.028, 0.004)
Diabetes			-0.024 (-0.043, -0.005)	-0.023 (-0.041, -0.005)
Hemoglobin (per 1gm/dL)				0.008 (0.003, 0.013)
P for continuous GFRcysc		0.026	0.004	0.016
P value for age		0.219	0.075	0.051
P value for sex		0.824	0.866	0.597
P value for height		0.859	0.684	0.354
P value for weight		0.081	0.101	0.088
P value for African American race		0.263	0.109	0.135
P value diabetes			0.012	0.012
P value for hemoglobin				0.002

Model 1: Age, sex, race (African American vs. other), height, weight, education, smoking, baseline gait speed

Model 2: + DM, Any CVD (CAD, PVD, stroke)

(Sensitivity Analysis): + Bicarbonate, hemoglobin, phosphate. Note hemoglobin is the main effect and interaction term with time

GFRcr-cysc=Cystatin C based eGFR with creatinine, DM=diabetes mellites, CVD=cardiovascular disease, CAD=coronary artery disease, PVD=peripheral vascular disease

Supplementary Table 3. Associations with fastest tertile of decline in gait speed over 3 years of follow-up using eGFR with combined cystatin and creatinine

	Model 1		Model 2	
	IRR (95% CI)	P-value	IRR (95% CI)	P-value
eGFRcr-cysc (per 30 ml/min per 1.73m² lower)	1.39 (1.06, 1.83)	0.018*	1.44 (1.07, 1.93)	0.016*
Age (per 1 yr older)	1.01 (0.99, 1.03)	0.357	1.00 (0.98, 1.03)	0.721
Height (in)	1.04 (0.96, 1.12)	0.388	1.04 (0.95, 1.13)	0.385
Weight (lbs)	1.00 (1.00, 1.01)	0.417	1.00 (1.00, 1.01)	0.302
Female gender	0.76 (0.33, 1.76)	0.518	0.80 (0.31, 2.11)	0.659
African American	1.37 (0.97, 1.94)	0.077	1.50 (1.03, 2.19)	0.034*
Education (Greater than HS vs HS or less)	0.85 (0.52, 1.38)	0.501	0.89 (0.59, 1.32)	0.552
Current smoking	1.81 (1.01, 3.25)	0.047*	1.61 (0.93, 2.78)	0.088
Diabetes			1.16 (0.80, 1.69)	0.427
Any cardiovascular disease			1.46 (1.05, 2.04)	0.025*
Log CRP (per 1 log mg/L)			1.04 (0.90, 1.19)	0.595

Fastest tertile of gait speed decline defined as a median 17.5% per year decrease in gait speed [IQR 12, 28.3]

GFRcr-cysc=Cystatin C based eGFR with creatinine, HS=high school, CRP=C-reactive protein; ** $p<0.01$, * $p<0.05$