

SUPPLEMENTSupplemental Table 1: PCR assay primers and probes by *Plasmodium* species

<i>Plasmodium</i> species	Factor	Final molarity	Sequence (5' to 3')
<i>malariae</i>	Forward Primer	300 nM	AGTTAAGGGAGTGAAGACGATCAGA
	Reverse Primer	300 nM	CAACCCAAAGACTTTGATTTCTCATAA
	Probe	200 nM	FAM/ATGAGTGTTTCTTTTAGATAGC/MGBNFQ
<i>ovale</i> spp.	Forward Primer	400 nM	CCRACTAGGTTTTGGATGAAVVRTTTTT
	Reverse Primer	400 nM	AACCCAAAGACTTTGATTTCTCATAA
	Probe	200 nM	VIC/CRAAAGGAATTYTCTTATT/MGBNFQ
<i>vivax</i>	Forward Primer	400 nM	ACGCTTCTAGCTTAATCCACATAACT
	Reverse Primer	400 nM	ATTTACTCAAAGTAACAAGGACTTCCAAGC
	Probe	200 nM	FAM/TTCGTATCG/ZEN/ACTTTGTGCGCATTTTGC/3IABkFQ
<i>falciparum</i>	Forward Primer	300 nM	ATTGCTTTTGAGAGGTTTTGTTACTTT
	Reverse Primer	300 nM	GCTGTAGTATTCAAACACAATGAACTCAA
	Probe	200 nM	FAM/CATAACAGACGGGTAGTCAT
<i>ovale curtisi</i>	Forward Primer	300 nM	TTTTGAAGAATACATTAGGATACAATTAATG
	Reverse Primer	300 nM	CATCGTTCCTCTAAGAAGCTTTACAAT
	Probe	200 nM	HEX/CCTTTTCCC/ZEN/TATTCTACTTAATTCGCAATTCATG
<i>ovale wallikeri</i>	Forward Primer	300 nM	TTTTGAAGAATATATTAG-GATACATTATAG
	Reverse Primer	300 nM	CATCGTTCCTCTAAGAAGCTTTACAAT
	Probe	200 nM	FAM/CCTTTTCCCTTTTCTACTTAATTCGCTATTCATG/TAMRA

Supplemental Table 2: PCR assay control reactions

<i>Plasmodium</i> species	Parasites / μ L	N Controls*	# Positive [†]	% Positive	Mean Ct	Standard deviation Ct
<i>malariae</i>	Std. 1 – 10,000 p/ μ L	20	20	100	23.7	0.64
	Std. 2 – 1000 p/ μ L	20	20	100	27.4	0.69
	Std. 3 – 100 p/ μ L	20	20	100	31.1	0.96
	Std. 4 – 10 p/ μ L	20	20	100	34.5	1.44
	Std. 5 – 1 p/ μ L	20	6	30	37.4	1.47
	Neg. – 0 p/ μ L	20	0	0	--	--
<i>ovale</i> spp.	Std. 1 – 10,000 p/ μ L	19	19	100	27.7	2.93
	Std. 2 – 1000 p/ μ L	20	19	95	31.0	2.79
	Std. 3 – 100 p/ μ L	19	19	100	34.3	2.39
	Std. 4 – 10 p/ μ L	20	14	70	37.0	1.64
	Std. 5 – 1 p/ μ L	20	10	50	40.7	1.31
	Neg. – 0 p/ μ L	20	0	0	--	--
<i>vivax</i>	Std. 1 – 10,000 p/ μ L	20	20	100	25.7	0.54
	Std. 2 – 1000 p/ μ L	20	20	100	28.5	0.86
	Std. 3 – 100 p/ μ L	20	18	90	32.5	0.86
	Std. 4 – 10 p/ μ L	20	20	100	36.1	0.59
	Std. 5 – 1 p/ μ L	20	15	75	39.0	1.07
	Neg. – 0 p/ μ L	20	0	0	--	--
<i>falciparum</i>	Std. 1 – 10,000 p/ μ L	20	20	100	20.4	0.46
	Std. 2 – 1000 p/ μ L	20	20	100	24.4	0.29
	Std. 3 – 100 p/ μ L	20	20	100	27.7	0.35
	Std. 4 – 10 p/ μ L	20	19	95	31.7	0.26
	Std. 5 – 1 p/ μ L	20	10	50	35.5	0.84
	Neg. – 0 p/ μ L	20	0	0	--	--

Note: Negative controls used water for each *Plasmodium* species.

*Controls were run in duplicate per PCR plate.

[†]PCR assays in which negative controls amplified incorrectly (e.g., due to contamination) are re-run until controls remain negative; Results from original assays are then discarded as invalid and the re-run assay results are reflected in final analyses.

Supplemental Table 3: Comparison of characteristics between students in the full SMPS cohort vs. sample of students in the sub-cohort for non-falciparum analysis.

Student characteristics	Full cohort	Sub-cohort	p-value*
n	17,131	3,456	
Age (years), median (IQR)	11 (9, 13)	11 (9, 13)	0.884
Male, n (%)	8457 (49)	1761 (51)	0.093
Malaria RDT+, n (%)	3328 (20)	686 (20)	0.589
Elevation (m), median (IQR)	1230 (1058, 1458)	1230 (1058, 1467)	0.925
Epidem. risk strata, n (%)			0.966
High	8806 (51)	1768 (51)	
Moderate	2180 (13)	448 (13)	
Low	2952 (17)	602 (17)	
Very low	3193 (19)	638 (18)	
Region, n (%)			1
Arusha	2757 (16)	552 (16)	
Iringa	1583 (9)	320 (9)	
Kagera	3078 (18)	619 (18)	
Mara	2228 (13)	452 (13)	
Mtwara	1511 (9)	307 (9)	
Rukwa	1506 (9)	301 (9)	
Tabora	2045 (12)	413 (12)	
Tanga	2423 (14)	492 (14)	

IQR: Interquartile range; rapid diagnostic test

*Continuous variables are compared using a Kruskal-Wallis test. Categorical variables are compared using Chi-square test.

Supplement Table 4: Prevalences and 95% confidence intervals of species-specific malaria infections

	Prevalence*		
	n	%	95% CI
Total student population	3456	100.00	---
Total infections†			
Po	814	23.6	22.2 – 25.0
Pm	136	3.9	3.3 – 4.6
Pv	11	0.3	0.2 – 0.6
Pf	755	21.8	20.5 – 23.3
Single-species infections			
Po	519	15.0	13.9 – 16.3
Pm	24	0.7	0.5 – 1.0
Pv	4	0.1	0.1 – 0.3
Pf	429	12.4	11.4 – 13.6
Mixed-species infections‡			
<i>P. falciparum</i> co-infection			
Po+Pf	224	6.5	5.7 – 7.4
Pm+Pf	44	1.3	1.0 – 1.7
Pv+Pf	2	0.1	0.02 – 0.2
Po+Pm+Pf	55	1.6	1.2 – 2.1
Po+Pv+Pf	1	0.03	0.01 – 0.2
Pm+Pv+Pf	0	0	0
Po+Pm+Pv+Pf	0	0	0
<i>Non-falciparum</i> only			
Po+Pm	12	0.3	0.2 – 0.6
Po+Pv	3	0.1	0.03 – 0.3
Pm+Pv	1	0.03	0.01 – 0.2
Po+Pm+Pv	0	0	0

CI: Confidence interval; Pf: *Plasmodium falciparum*; Po: *Plasmodium ovale* spp.; Pm: *Plasmodium malariae*; Pv: *Plasmodium vivax*

*Prevalences represent point prevalences. Percents are calculated out of the total study population.

†Total infections include both single and mixed-species infections.

‡Categories are mutually exclusive.

Supplemental Table 5. Sensitivity analysis: *P. falciparum* and *P. malariae* prevalence estimated under a real-time PCR Ct of 45 (N=750)*,†

<i>Plasmodium</i> species	Epidemiologic risk strata			Total N=750	Crude prevalence		Weighted prevalence [§]	
	Low [‡] n=250	Moderate n=250	High n=250		%	95% CI	%	95% CI
<i>P. falciparum</i>								
PCR positive	5 (2.0%)	47 (18.8%)	105 (42.0%)	157 (20.9%)	20.9%	17.8%-24.5%	24.6%	20.6%-29.2%
Ct <40	5 (2.0%)	45 (18.0%)	103 (41.2%)	153 (20.4%)				
Ct 40 to <45	0 (0.0%)	2 (0.8%)	2 (0.8%)	4 (0.5%)				
PCR Negative	245 (98.0%)	203 (81.2%)	145 (58.0%)	593 (79.1%)				
<i>P. malariae</i>								
PCR positive	2 (0.8%)	6 (2.4%)	12 (4.8%)	20 (2.7%)	2.7%	1.6%-4.1%	3.1%	1.8%-4.9%
Ct <40	1 (0.4%)	6 (2.4%)	11 (4.4%)	18 (2.4%)				
Ct 40 to <45	1 (0.4%)	0 (0.0%)	1 (0.4%)	2 (0.3%)				
PCR Negative	248 (99.2%)	244 (97.6%)	238 (95.2%)	730 (97.3%)				

*Real-time PCR assay targeting the 18S ribosomal subunit; a Ct of 45 was employed for sensitivity analysis

†Sensitivity analysis population is a stratified random sample of the non-falciparum analysis population, where sampling was stratified by epidemiologic malaria risk strata to maintain malaria transmission heterogeneity within the sample.

‡Low epidemiological risk classification comprises 'very low' and 'low' strata.

§Weighted prevalence estimates are weighted to the distribution of students in the non-falciparum analysis population by epidemiologic malaria risk strata to account for oversampling of the moderate risk strata in the sensitivity analysis population.

Supplemental Table 6. Student infection characteristics by PCR+ vs. PCR– non-falciparum infection

Student characteristics, n (%) ^{*,†}	<i>P. ovale</i> spp.			<i>P. malariae</i>			<i>P. vivax</i>		
	PCR +		p-value	PCR +		p-value	PCR +		p-value
	Any Infection	No infection		Any Infection	No infection		Any Infection	No infection	
N	814	2138		136	2138		11	2138	
Age (years)									
Median (IQR)	11 (9, 13)	11 (9, 13)	0.128	12 (9, 13)	11 (9, 13)	0.038	11 (9.5, 12)	11 (9, 13)	0.95
Male, n (%)	402 (49.4)	1075 (50.3)	0.694	83 (61.0)	1075 (50.3)	0.019	8 (72.7)	1075 (50.3)	0.226
Fever [‡] , n (%)	23 (3.6)	27 (1.5)	0.003	12 (10.9)	27 (1.5)	<0.001	0 (0.0)	27 (1.5)	1
Epidemiological risk strata [§] , n (%)			<0.001			<0.001			0.001
High	565 (69.4)	770 (36.0)		125 (91.9)	770 (36.0)		10 (90.9)	770 (36.0)	
Moderate	63 (7.7)	327 (15.3)		10 (7.4)	327 (15.3)		1 (9.1)	327 (15.3)	
Low	160 (19.7)	433 (20.3)		1 (0.7)	433 (20.3)		0 (0.0)	433 (20.3)	
Very low	26 (3.2)	608 (28.4)		0 (0.0)	608 (28.4)		0 (0.0)	608 (28.4)	

PCR: polymerase chain reaction; Po: *Plasmodium ovale* spp.; Pm: *Plasmodium malariae*; Pv: *Plasmodium vivax*; RDT: rapid diagnostic test

*Continuous variables are compared using a Kruskal-Wallis test. Categorical variables are compared using chi-square; fisher's exact test was applied where cell counts were <5.

[†]PCR negative represents students confirmed by PCR to have no malaria infection of any species.

[‡]Fever was defined as temperature at time of survey $\geq 38^{\circ}\text{C}$. Temperature is missing for n=618 (17.9%) subjects; percentages are calculated out of non-missing data.

[§]Epidemiological risk strata are defined as follows, based on *P. falciparum* prevalences in children from the 2014-15 Tanzanian SMPS: "very low" if prevalence <5%, "low" 5 to <10%, "moderate" 10 to <50%, and "high" $\geq 50\%$.