

Supplemental Material

Imputation of B-lines

In patients with missing B-line data in ≤ 2 out of 4 zones (n=7), we imputed B-line data from anatomically adjacent zones: Zones 1 and 2, zones 3 and 4, zones 5 and 6, zones 7 and 8.

	Zones 1 & 2	Zones 3 & 4	Zones 5 & 6	Zones 7 & 8
Correlation between adjacent zones (rho, P)	0.32 (p<0.001)	0.14 (p=0.13)	0.28 (p=0.002)	0.13 (p=0.154)
Imputed zones (n)	n=1	n=1	n=2	n=3

Supplementary Table S1. Baseline characteristics of total population and according to lung ultrasound

	Total	Negative LUS	Positive LUS	p-value
	N=123	N=107	N=16	
Age (years)	75 ± 9	74 ± 9	75 ± 9	0.74
Male	57 (46%)	50 (47%)	7 (44%)	0.82
Smoking	52 (42%)	43 (40%)	9 (56%)	0.23
Body mass index (kg/cm2)	23.8 ± 6.4	23.8 ± 6.4	23.7 ± 6.3	0.95
FEV ₁				
FEV ₁ > 50%	29 (25%)	24 (24%)	5 (31%)	0.85
FEV ₁ 30 - 50%	46 (40%)	40 (41%)	6 (38%)	
FEV ₁ < 30%	39 (34%)	34 (35%)	5 (31%)	
Diabetes	16 (13%)	13 (12%)	3 (19%)	0.46
Hypertension	44 (36%)	36 (34%)	8 (50%)	0.20
Acute myocardial infarction	33 (27%)	29 (27%)	4 (25%)	0.86
Atrial fibrillation	20 (16%)	18 (17%)	2 (13%)	0.66
Diuretics pre-admission	52 (42%)	44 (41%)	8 (50%)	0.50
Diuretics prior to LUS	61 (50%)	55 (51%)	6 (38%)	0.30
Respiration frequency (breaths/minute)	27 ± 7	27 ± 7	27 ± 7	0.96
Peripheral O ₂ saturation (%)	89 ± 7	89 ± 7	87 ± 7	0.21
Systolic blood pressue (mmHg)	139 ± 25	138 ± 24	144 ± 28	0.33
Pulse (beats/minute)	95 ± 18	95 ± 19	93 ± 13	0.67
C-reactive protein (ng/L)	20 (7-60)	21 (6-60)	16 (10-71)	0.84
NT-proBNP (ng/L)	455 (125-1247)	444 (120-1029)	855 (130-2949)	0.29
High sensitive Troponin T (ng/L)	30 (18-52)	31 (17-52)	28 (20-45)	0.66
Creatinine (umol/L)	69 (58-92)	69 (57-98)	71 (59-89)	0.72
Hemoglobin (g/dL)	13.6 ± 1.9	13.6 ± 1.9	13.7 ± 1.5	0.74
Arterial O ₂ (kPa)	8.8 ± 3.4	8.9 ± 3.6	8.3 ± 1.5	0.54
Chest X-ray - congestion	36 (30%)	32 (30%)	4 (27%)	0.76
Chest X-ray - infiltrate	29 (24%)	21 (20%)	8 (53%)	0.005
Length of stay (days)	5 (3-7)	5 (3-8)	4 (3-6)	0.41

FEV₁=forced expiratory volume in the 1st second, (n=114), LUS=lung ultrasound, NTproBNP=N-terminal pro-brain natriuretic peptide

Supplementary Table 2. Predictors of the total number of B-lines on LUS. Analyzed by negative binomial regression with total B-lines number as dependent variable¹

Independent variable	IRR (95% CI)	p-value
Age (per 5 years)	1.14 (1.07-1.21)	<0.001
Male	1.02 (0.80-1.29)	0.88
BMI (per unit)	0.98 (0.69-1.10)	0.047
Smoking	0.87 (-0.38- 0.10)	0.25
Emphysema	1.12 (0.87-1.43)	0.39
FEV ₁ 30-50%	0.97 (0.71-1.33)	0.86
FEV ₁ <30%	0.78 (0.56-1.08)	0.14
Concurrent acute HF	1.40 (1.11-1.76)	0.004
NTproBNP (per ln increase)	1.05 (0.97-1.14)	0.20
CRP (per ln increase)	1.11 (1.03-1.20)	0.007
Creatinine (per umol/L)	1.15 (0.85-1.55)	0.37
Arterial pressure O ₂ (kPa)	0.96 (0.93- 0.99)	0.005
Diuretics given prior to LUS	1.37 (1.09-1.72)	0.007
Diuretics newly initiated	1.44 (1.04-1.99)	0.029
Non-invasive ventilation	0.88 (0.68-1.14)	0.34
Chest X-ray - congestion	1.27 (0.99-1.64)	0.061
Chest X-ray - infiltrate	1.40, (1.08-1.83)	0.012

BMI=body mass index, CRP=C-reactive protein, FEV₁=forced expiratory volume in the 1st second, (n=114), HF=heart failure, IRR=incidence-rate ratios, LUS=lung ultrasound, NTproBNP=N-terminal pro-brain natriuretic peptide

¹one unit increase in the independent variable leads to % change in sum of B-lines in 8 zones.