

Supplemental to:

Association of Biological Age with Tumor Microenvironment in Patients with Esophageal Adenocarcinoma

C.J. Ravensbergen^{1†}, Y. van Holstein^{2†}, S.C. Hagenaars¹, A.S.L.P. Crobach³, S. Trompet², J.E.A. Portielje⁴, N.A. de Glas⁴, D. van Heemst², F. van den Bos², R.A.E.M. Tollenaar¹, W.E. Mesker^{1±}, S.P. Mooijaart^{2*} & M. Slingerland^{4*}

¹Department of Surgery, Section Surgical Oncology, Leiden University Medical Center, Leiden, the Netherlands

²Department of Internal Medicine, Section of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands


³Department of Pathology, Leiden University Medical Center, Leiden, the Netherlands

⁴Department of Medical Oncology, Leiden University Medical Center, Leiden, the Netherlands

[†]Contributed equally


^{*}Shared senior author

[±]Correspondence to: Dr. W.E. Mesker, Department of Surgery, Leiden University Medical Center, Albinusdreef 2, P.O. Box 9600, 2300RC Leiden, the Netherlands. Mail: w.e.mesker@lumc.nl




Supplementary Figure S1. Flowchart of cohort selection. Out of the 76 patients with esophageal adenocarcinoma selected for this study, 72 were eligible for histologic analysis and subsequently included.


A




B



Supplementary Figure S2. Histologic intratumoral stroma percentage and age. (A) Scatter correlation plot of histologic intratumoral stroma percentage and chronological age. (B) Stacked bar plot of the proportions of stroma-high and stroma-low tumors, as scored by the standardized histologic tumor-stroma ratio, in strata of biological age.



Supplementary Figure S3. Scatter correlation plot of histologic tumor immune cell infiltrate (TICI) and chronological age.



Supplementary Figure S4. Spearman correlation coefficient matrix of chronological age, histologic parameters, and blood cell counts. *X*, not significant.

Supplementary Table S1. Statistics for a multiple linear regression model containing strata of biological age, chronological age, male sex, clinical T and N stage as predictor variables and histologic intratumoral stroma percentage as dependent variable. The fit older adult group was selected as a reference.

Variables	Coefficient (B)	SE	95% CI	T statistic	P-value
Constant	64.02	31.30	1.500 to 126.5	2.045	0.045*
Younger adult	-4.981	8.551	-22.06 to 12.10	-0.583	0.562
Frail older adult	-14.66	6.233	-27.11 to -2.213	-2.352	0.022*
Chronological age	-0.251	0.356	-0.961 to 0.460	-0.704	0.484
Male sex	0.473	5.960	-11.43 to 12.38	0.079	0.937
cT stage	0.998	3.295	-5.583 to 7.579	0.303	0.763
cN stage	-0.215	2.803	-5.813 to 5.383	-0.077	0.939

$R^2_{adj} = 0.051$, $P = 0.151$

Abbreviations: SE, standard error; CI, confidence interval.

*Statistically significant at $P < 0.05$, two-tailed

Supplementary Table S2. Statistics for a multiple linear regression model containing strata of biological age, chronological age, male sex, clinical T and N stage as predictor variables and blood thrombocyte count at the time of diagnosis as dependent variable. The fit older adult group was selected as a reference.

Variables	Coefficient (B)	SE	95% CI	T statistic	P-value
Constant	248.0	128.0	-7.732 to 503.6	1.937	0.057
Younger adult	15.77	35.01	-54.16 to 85.71	0.451	0.654
Frail older adult	-44.94	26.06	-97.00 to 7.114	-1.725	0.089
Chronological age	0.821	1.455	-2.085 to 3.728	0.565	0.574
Male sex	-32.76	24.50	-81.71 to 16.18	-1.337	0.186
cT stage	-4.651	13.47	-31.55 to 22.25	-0.345	0.731
cN stage	15.99	11.56	-7.110 to 39.10	1.383	0.172

$R^2_{adj} = 0.010$, $P = 0.363$

Abbreviations: SE, standard error; CI, confidence interval.

*Statistically significant at $P < 0.05$, two-tailed

Supplementary Table S3. Statistics for a multiple linear regression model containing strata of biological age, chronological age, male sex, clinical T and N stage as predictor variables and blood leukocyte count at the time of diagnosis as dependent variable. The fit older adult group was selected as a reference.

Variables	Coefficient (B)	SE	95% CI	T statistic	P-value
Constant	12.08	3.746	4.599 to 19.57	3.225	0.002*
Younger adult	-0.865	1.025	-2.912 to 1.182	-0.884	0.402
Frail older adult	-0.838	0.763	-2.361 to 0.686	-1.098	0.276
Chronological age	-0.034	0.043	-0.119 to 0.051	-0.797	0.428
Male sex	-0.376	0.717	-1.809 to 1.057	-0.524	0.602
cT stage	-0.325	0.394	-1.113 to 0.462	-0.825	0.412
cN stage	0.549	0.339	-0.127 to 1.226	1.623	0.110

$R^2_{adj} = -0.017$, $P = 0.571$

Abbreviations: SE, standard error; CI, confidence interval.

*Statistically significant at $P < 0.05$, two-tailed