

Supplementary Information

Table S1: Characteristics of included sites, 2012-2019 influenza seasons

Coordinating site	Included patient specimens	Number of contributing seasons	Country income level ^a	Specimen distribution (+/- flu)	Distribution: number of comorbidities
Canada (Halifax)	1,351	2	High income		
France (Lyon)	211	2	High income		
Czech Republic (Prague)	461	4	High income		
Spain (Valencia)	21,304	7	High income		
Poland (Warsaw)	22	1	High income		
Kazakhstan (Almaty)	259	1	Upper middle income		
China (Beijing)	4,656	3	Upper middle income		
Lebanon (Beirut)	522	1	Upper middle income		
Romania (Bucharest)	1,822	3	Upper middle income		
Argentina (Buenos Aires)	902	2	Upper middle income		
Colombia (Cartagena)	42	1	Upper middle income		
Brazil (Curitiba)	378	1	Upper middle income		
Turkey (Ankara)	2,373	4	Upper middle income		
South Africa (Johannesburg)	4,705	3	Upper middle income ^b		
Peru (Lima)	530	3	Upper middle income		
Mexico (Mexico City)	1,971	4	Upper middle income		
Russian Federation (Moscow)	8,549	6	Upper middle income		
Serbia (Novi Sad)	1,386	2	Upper middle income		
China (Shanghai)	2,764	3	Upper middle income		
Russian Federation (St. Petersburg)	14,184	6	Upper middle income		
Côte d'Ivoire (Abidjan)	983	2	Lower middle income		
Tunisia (Ariana)	77	1	Lower middle income		
Kenya (Nairobi)	1,002	2	Lower middle income		
India (Srinagar)	2,667	4	Lower middle income		
All sites	73,121				
^a According to World Bank GNI per capita, calendar year 2019					
^b Romania designated upper middle income in 2 out of 3 years of contribution to network					

Figure S1: Age distribution of patients enrolled by site, and whether they were influenza-laboratory confirmed, GIHSN, 2012-2019




Table S2: Collation of dataset for analysis

We removed 86 specimens serving as internal controls for the coordinating sites and 1531 specimens missing patient age. We then removed 54,954 specimens for which questionnaires were partially or fully completed but which did not meet GIHSN criteria for inclusion (i.e., patients were recently discharged, were institutionalized, or were not residents of the hospital catchment area). Finally, we removed 279 specimens with inconsistent subtype or lineage data (e.g., a “yes” for A/H1N1pdm09 and a “yes” for unable to be subtyped) and 93 influenza-positive specimens with coinfections of more than one influenza strain.

The data used in the regression analyses of the influenza-positive data excludes 88 influenza-positive specimens for which ICU admission information is unavailable (ICU model) and 91 influenza-positive specimens for which it is unknown whether the patient died in the hospital (in-hospital death model). The data used in the regression analysis of influenza-negative specimens excludes 352 influenza-negative specimens for which ICU admission information is unavailable (ICU model), 321 influenza-negative specimens for which it is unknown whether the patient died in the hospital (in-hospital death model), and 257 influenza-negative specimens for which comorbidity information is not recorded (both models).

Figure S2: GIHSN data collection protocol




Table S3: Demographic characteristics of all patients hospitalized with severe acute respiratory illness by country income level, GIHSN 2012-2019

		Country Income Level			
	Total	High Income	Upper Middle Income	Lower Middle Income	<i>p-value</i>
	N = 73121	N = 23349	N = 45043	N = 4729	
	n (%)	n (%)	n (%)	n (%)	
Age group					
<5 years	29450 (40.3)	4460 (19.1)	23149 (51.4)	1841 (38.9)	<0.001
5-64 years	25191 (34.5)	5579 (23.9)	17845 (39.6)	1767 (37.4)	
≥65 years	18480 (25.3)	13310 (57.0)	4049 (9.0)	1121 (23.7)	
Sex					
female	34161 (46.7)	10833 (46.4)	21075 (46.8)	2253 (47.6)	0.26
male	38960 (53.3)	12516 (53.6)	23968 (53.2)	2476 (52.4)	
Number of comorbidities					
zero	35125 (48.0)	6497 (27.8)	27284 (60.6)	1344 (28.4)	<0.001
one	21711 (29.7)	6645 (28.5)	12993 (28.8)	2073 (43.9)	
two or more	16028 (21.9)	10207 (43.7)	4509 (10.0)	1312 (27.7)	
missing	257 (0.4)	0 (0)	257 (0.6)	0 (0)	
Comorbidities					
cardiovascular disease	14597 (20.0)	8723 (37.4)	4657 (10.3)	1217 (25.7)	<0.001
chronic obstructive pulmonary disease	9449 (12.9)	6467 (27.7)	2217 (4.9)	765 (16.2)	
asthma	3552 (4.9)	1978 (8.5)	1464 (3.3)	110 (2.3)	
diabetes	7488 (10.2)	5563 (23.8)	1434 (3.2)	491 (10.4)	
immunosuppression (rheumatologic/autoimmune, neoplasm)	20288 (27.8)	6143 (26.3)	11779 (26.2)	2366 (50.0)	
renal disease	3871 (5.3)	2781 (11.9)	895 (2.0)	195 (4.1)	
cirrhosis	1236 (1.7)	782 (3.4)	427 (1.0)	27 (0.6)	
neuromuscular disorder	2327 (3.2)	1239 (5.3)	946 (2.1)	142 (3.0)	
missing	257 (0.4)	0 (0)	257 (0.6)	0 (0)	
Smoking status of patient or household					
currently smokes	15872 (21.7)	3509 (15.0)	11644 (25.9)	719 (15.2)	<0.001
has smoked previously	13801 (18.9)	6396 (27.4)	6681 (14.8)	724 (15.3)	
has never smoked	42358 (57.9)	12634 (54.1)	26461 (58.7)	3263 (69.0)	
missing	1090 (1.5)	810 (3.5)	257 (0.6)	23 (0.5)	

Figure S3: Distribution of all and influenza positive-only specimens by country income bracket

Distribution of All Patient Specimens by Country Income Bracket

Distribution of Influenza-Positive Patient Specimens by Country Income Bracket




Table S4: Comparing the odds ratios of infection with A/H1N1pdm09 or A/H3N2 in patients with laboratory-confirmed influenza using Tukey contrasts

Dataset	Outcome	Linear Hypothesis	Estimate	<i>p</i>
All age groups	ICU admission	A/H3N2 – A/H1N1pdm09 = 0	-1.00	< 0.001
All age groups	In-hospital death	A/H3N2 – A/H1N1pdm09 = 0	-0.86	< 0.001
≥65 years only	ICU admission	A/H3N2 – A/H1N1pdm09 = 0	-0.76	< 0.001
≥65 years only	In-hospital death	A/H3N2 – A/H1N1pdm09 = 0	-0.56	< 0.05

Figure S4: Risk factors for (A) influenza-related intensive care unit admission, (B) influenza-related in-hospital death, (C) non-influenza intensive care unit admission, and (D) non-influenza in-hospital death, among patients from HIC hospitalized with severe acute respiratory illness

