

Patient	Disease	Clinical	Time since sclerosis	Pathology group	Degree of infiltrate	Necrotic keratinocytes
U1	Unaffected	none	none	normal	unaffected	none
U2	Unaffected	none	none	normal	unaffected	none
U3	Unaffected	none	none	normal	unaffected	none
U4	Unaffected	none	none	normal	unaffected	none
U5	Unaffected	none	none	normal	unaffected	none
U6	Unaffected	none	none	normal	unaffected	none
U7	Unaffected	none	none	normal	unaffected	none
U8	Unaffected	none	none	normal	unaffected	none
U9	Unaffected	none	none	normal	unaffected	none
A1	Affected	stable	late	Scl + epi	mild	mild
A2	Affected	progressive	early	Scl + epi	moderate	mild
A3	Affected	softening	late	Scl + epi	mild	mild
A4	Affected	softening	early	Scl + peri	moderate	none
A5	Affected	progressive	late	Scl + epi	mild	moderate
A6	Affected	stable	late	sclerotic	mild	none
A7	Affected	progressive	late	Scl + epi	mild	moderate
A8	Affected	stable	early	sclerotic	mild	none
A9	Affected	stable	late	Scl + peri	mild	none
A10	Affected	stable	early	sclerotic	mild	none
A11	Affected	progressive	early	sclerotic	mild	none
A12	Affected	stable	late	Scl + epi	mild	moderate
A13	Affected	stable	early	Scl + epi	moderate	mild
A14	Affected	stable	early	Scl + epi	mild	moderate
A15	Affected	stable	late	Scl + epi	moderate	none
A16	Affected	progressive	early	Scl + epi	moderate	moderate
A17	Affected	progressive	early	Scl + epi	mild	mild

**Supplementary Table S3: Clinical and histopathologic correlative characteristics of patients.** U (unaffected); A (affected); Clinical: as determined by patient and physician assessment; early (less than 1 year since sclerosis onset); late (greater than 1 year); sclerotic (sclerotic changes without significant inflammation); Scl + peri (sclerotic changes and perivascular inflammation without significant epidermal changes); Scl + epi (sclerotic changes with necrotic keratinocytes, melanophages, and/or interface dermatitis); normal (normal histopathology); for the degree of infiltrate, mild (scattered immune cells either in a perivascular or interface pattern); moderate (a moderate inflammatory cell infiltrate in the dermis and/or a combination of a perivascular and interface inflammatory cell infiltrate); for necrotic keratinocytes, none (no necrotic keratinocytes identified), mild (at least 1 identified in the pathology section), or moderate (multiple identified in the pathology section).