

Appendix C: Risk of bias assessment for each of the studies

Abbreviations: m: male; f: female; No information on status: no assessment of the participants' physical health, mental health or fall history

Table C.1 Risk of bias assessment for the studies comparing higher and lower risk older adults

| Bias domain | Total bias | 1 – Confounding factors | 2 – Participant selection | 3 – Group allocation based on fall risk assessment | 4 – Intended experiment | 5 – Motion analysis outcome parameters | 6 – Missing data | 7 – Result reporting |
|------------------|------------|--|--|---|---|--|---|---|
| Stairs | | | | | | | | |
| Ackermans (2019) | Low | No information on gender. | No information on gender. | Low Retrospective fall history | Low Subjects were allowed to take breaks to avoid fatigue. | Low | Low | Low |
| Ackermans (2021) | Moderate | No information on gender balance within groups. | Moderate Gender is unbalanced: 28 m, 53 f | Low Prospective fall history | Low | Low | Low | Low |
| Brodowski (2021) | Serious | Serious Participants in the hospitalized group were significantly older than in the healthy group (p < 0.001, mean age difference 7 years). | Moderate Gender is unbalanced: 43 m, 70 f 3 of the 60 hospitalized subjects were excluded because they could not perform the timed-up-and-go test. | Moderate Health status (hospitalized versus healthy) | Low | Low | Moderate 4 of the 60 healthy participants were excluded because they performed the task very rapidly or very slowly. 4 of the 56 hospitalized subjects could not perform the 2 nd trial. | Moderate Significance level was not reported, but sample size was sufficient for the reported effect size. |
| Carli (2014) | Moderate | No information on gender balance within groups. | Moderate Gender is unbalanced: 4 m, 30 f. | Moderate Physical level | No information on fatigue prevention strategy. | Low | Low | Low |

| | | | | | | | | |
|----------------------|----------|---|--|--|--|--|--|---|
| Oh-Park (2012) | Moderate | Low | Moderate Gender is unbalanced: 147 m, 224 f. | Moderate Functional decline over 1 year | Low Subjects were allowed a brief rest. | Low | Low | Moderate Significance level was not reported, but sample size was sufficient for the reported effect size. |
| Oh-Park (2011) | Moderate | No information on gender balance within groups | Moderate Gender is unbalanced: 204 m & 309 f. | Low Retrospective fall history | Low Subjects were allowed a brief rest. | Low | Low | Low |
| Zietz (2011) | Serious | Serious Higher risk participants were significantly older ($p < 0.05$, mean age difference 7 years). | Moderate Gender is unbalanced: 3 m, 19 f | Moderate Physical and mental level | Moderate 4 out of the 7 high fall risk older adults (and no other participant) used the handrail. No information on fatigue prevention strategy. | Serious No information on how the center of mass is calculated. | Low | Low |
| Perturbations | | | | | | | | |
| Gerards (2021) | Low | Low | Low | Low Retrospective fall history | No information on randomization. No information on fatigue prevention strategy. | Low | Low One subject was excluded because of technical failure | Low |
| Pijnappels (2005) | Moderate | Low | No information on status. | Moderate Physical level | No information on fatigue prevention strategy. No information on randomization. | Low | Moderate Older adults (but no younger adults) sometimes performed “a lowering strategy at mid-swing”, and these trials were | Low |

| | | | | | | | | |
|------------------|----------|---|--|-----------------------------------|--|-----|---|-----|
| | | | | | | | removed from the analysis. | |
| Obstacles | | | | | | | | |
| Brach (2011) | Moderate | Low | Moderate Gender is unbalanced: 20 m, 87 f Exclusion of subjects with low walking speed (36 out of the 120 recruited subjects). | Low Retrospective fall history | No information on randomization. No information on fatigue prevention strategy. | Low | Moderate 13 out of 84 subjects did not complete the follow-up. | Low |
| Chou (2003) | Moderate | Moderate Gender balance is different across groups Y: 7 m, 2 f O: 1 m, 5 f | Low | Low | Low | Low | Low | Low |
| Hansson (2021) | Moderate | Low | Moderate Gender is unbalanced: 10 m, 91 f | Low | Low | Low | Low | |
| Guadagnin (2020) | Moderate | Low | Moderate Exclusion of male subjects. | Low Retrospective fall history | No information on fatigue prevention strategy. No information on randomization. | Low | Low | Low |
| Pan (2016) | Moderate | No information on gender. | No information on gender. | Moderate Physical level | Low Subjects were allowed to rest if needed. | Low | Low | Low |
| Pieruccini- | Moderate | Low | Moderate Exclusion of male subjects. | Low Retrospective fall history | No information on fatigue prevention strategy. | Low | Low | Low |

| | | | | | | | | |
|---------------|----------|-----|---|-----------------------------------|--|-----|-----|-----|
| Faria (2019) | | | | | No information on randomization. | | | |
| Uemura (2011) | Moderate | Low | Moderate Gender is unbalanced: 26 m, 50 f. | Low Retrospective fall history | No information on fatigue prevention strategy. | Low | Low | Low |

Table C.2 Risk of bias assessment for the studies comparing younger and older adults

| Bias domain | Total bias | 1 – Confounding factors | 2 – Participant selection | 4 – Intended experiment | 5 – Motion analysis outcome parameters | 6 – Missing data | 7 – Result reporting |
|------------------|------------|---|---|--|--|---|----------------------|
| Stairs | | | | | | | |
| Begg (2000) | Moderate | Low | Moderate Exclusion of male subjects. | No information on fatigue prevention strategy. | Low | Low | Low |
| Bosse (2012) | Low | Low | No information on status. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Chiu (2015) | Low | Low | No information on status. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Christina (2002) | Moderate | Moderate Gender balance is different across groups Y: 5 m, 7 f O: 8 m, 4 f | Low | No information on fatigue prevention strategy. | Low | Low 1 participant slipped and was removed from the analysis. | Low |

| | | | | | | | |
|------------------|----------|---|---|--|-----|---|-----|
| Crosbie (2003) | Low | Low | Low | Low | Low | Low | Low |
| Dewolf (2021) | Moderate | Moderate Gender balance is different across groups Y: 4 m, 4 f O: 9 m, 1 f | Moderate Gender is unbalanced: 13 m, 5 f | Low | Low | Low | Low |
| Dixona (2018) | Low | No information on gender. | Low | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Francksen (2022) | Low | No information on gender. | No information on status or gender. | Low | Low | Low 2 out of 27 younger adults and 1 out of 33 older adults were excluded due to technical issues. | Low |
| Francksen (2020) | Low | No information on gender. | No information on status or gender. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Foster (2019) | Low | Low | No information on status. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Hamel (2005) | Moderate | Low | Moderate Exclusion of male subjects. | No information on randomization. No information on fatigue prevention strategy. | Low | Low Only trials of a pre-specified speed were included in the analysis. | Low |

| | | | | | | | |
|---------------|----------|---------------------------|--|--|---|---|-----|
| Hsue (2014) | Moderate | Low | Moderate Exclusion of subjects with fall history. | Low | Low | Low | Low |
| Hsue (2009) | Moderate | Low | Moderate Exclusion of male subjects. | Low | Low | Low | Low |
| Kim (2009) | Moderate | Low | Moderate Exclusion of subjects with low physical test scores. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Larsen (2008) | Moderate | No information on gender. | Moderate Exclusion of subjects who do not participate in physical activities at least once per week. No information on gender. | No information on randomization. No information on fatigue prevention strategy. | Low | Low 1 out of 19 older subjects was excluded because they could not walk up the stairs without the handrails. | Low |
| Mian (2007) | Moderate | Low | Moderate Exclusion of female subjects. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Novak (2011) | Moderate | Low | Moderate Exclusion of subjects with fall history. Gender is unbalanced 21 m, 36 f. | No information on fatigue prevention strategy. | Low | Low 1 subject per group was excluded due to technical failure. | Low |
| Novak (2016) | Moderate | Low | No information on status. | Low Trials were randomized to prevent fatigue affecting the results. | Moderate Markers placed on the sacrum were used as a proxy for the | Low | Low |

| | | | | | | | |
|----------------------|----------|---|---|--|-----------------|---|-----|
| | | | | | center of mass. | | |
| Reeves (2007) | Moderate | Moderate Gender balance is different across groups Y: 10 m, 7 f O: 5 m, 10 f | No information on status. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Perturbations | | | | | | | |
| Afschrift (2019) | Low | No information on gender. | No information on gender. | Low | Low | Low | Low |
| Bosqué e (2021) | Low | No information on gender. | No information on status or gender. | Low | Low | Low | Low |
| Debelle (2021) | Moderate | Moderate Gender balance is different across groups Y: 8 m, 9 f O: 3 m, 14 f | Moderate Gender is unbalanced: 11 m, 23 f No information on status. | Low | Low | Low Trials for which the subjects required more than 15 recovery steps were not analyzed (9 younger adults trials and 9 older adults trials) | Low |
| Jeon (2022a) | Moderate | Moderate Gender balance is different across groups Y: 4 m, 6 f O: 7 m, 3 f | Low | Low | Low | Moderate 1 out of 10 young subjects was excluded because they had no arm reaction. | Low |

| | | | | | | | |
|-----------------|----------|---------------------------|--|---|-----|---|-----|
| Jeon (2022b) | Moderate | Low | No information on status | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Kazanski (2020) | Moderate | Low | Low | Low There was an approximately 2 minute rest between trials. | Low | Moderate 2 of the 17 older adults (and none of the younger adults) used the handrail, and were removed from the analysis. | Low |
| Laudani (2021) | Moderate | Low | Moderate Gender is unbalanced: 4 m, 16 f . Exclusion of highly active individuals. | Low | Low | Low | Low |
| Liu (2009) | Moderate | No information on gender. | No information on status or gender. | No information on randomization. No information on fatigue prevention strategy. | Low | Moderate “Participants who successfully recovered with two feet (i.e., stance foot and swing foot) on one force-plate were excluded from the current study due to the difficulty in performing inverse dynamics analyses.” | Low |
| Martelli (2017) | Moderate | Low | Moderate Exclusion of subjects with fall history. | Low Subjects were allowed to rest if needed. No information on randomization. | Low | Low 2 out of 24 trials (both in elderly subjects) were removed due to technical failure. | Low |

| | | | | | | | |
|-----------------|----------|---|---|--|--|---|-----|
| McCrum (2016) | Moderate | Low | Moderate Exclusion of male subjects. No information on status | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| McIntosh (2016) | Moderate | Low | Moderate Gender is unbalanced: 6 m, 15 f | Low | Low | Low | Low |
| Nachmani (2020) | Serious | No information on gender. | No information on gender. | Low Subjects were allowed to rest if needed. | Moderate The center of mass was calculated as the midpoint between both ASIS markers. | Serious 13 of the 35 older adults (and none of the younger adults) failed to complete the task and were removed from the analysis. | Low |
| Qiao (2018) | Moderate | Low | Moderate Exclusion of subjects with fall history or BMI > 30. | Low | Low | Low | Low |
| Ren (2022) | Moderate | No information on gender. | Moderate Exclusion of subjects with fall history. No information on gender. | Low | Low | Low | Low |
| Roeles (2018) | Moderate | Moderate Gender balance is different across groups Y: 6 m, 3 f O: 2 m, 7 f | Low | No information on fatigue prevention strategy. | Low | Low | Low |
| Rum (2020) | Moderate | Low | Moderate Gender is unbalanced: 4 m, 16 f | Low | Low | Low | Low |

| | | | | | | | |
|------------------|----------|--|---|--|-----|---|---|
| Shulman (2018) | Low | Low | Low | No information on fatigue prevention strategy. | Low | Low | Low |
| Shulman (2019) | Low | Low | No information on status. | No information on fatigue prevention strategy. | Low | Low 4 of the 560 trials (all in older adults) were removed from the analysis because participants contacted the floor with the forefoot prior to heel contact. | Low |
| Tropea (2015) | Serious | Moderate Gender balance is different across groups Y: 4 m, 2 f O: 2 m, 4 f | Low Exclusion of participants with low physical health (as assessed by a physical therapist) | No information on fatigue prevention strategy. No information on randomization. | Low | Low | Serious Significance level was not reported, and sample size was too low for the reported effect size. |
| Yoo (2021) | Moderate | Moderate Gender balance is different across groups Y: 7 m, 7 f O: 4 m, 10 f | Moderate Gender is unbalanced: 11 m, 17 f | Low | Low | Low | Low |
| Obstacles | | | | | | | |

| | | | | | | | |
|------------------|----------|-----|--|--|-----|--|-----|
| Caetano (2016) | Moderate | Low | Moderate Gender is unbalanced: 25 m, 46 f. | No information on fatigue prevention strategy. | Low | Moderate 22% of the older adults (and none of the younger adults) made at least 1 mistake, and trials with a mistake were removed from the analysis. | Low |
| Chen (1991) | Low | Low | Low | No information on fatigue prevention strategy. | Low | Low | Low |
| Chen (1994) | Moderate | Low | Low | No information on fatigue prevention strategy. | Low | Moderate Trials in which participants failed to step over an obstacle were removed from the analysis. There is no information on how often this occurred. | Low |
| Chien (2018) | Moderate | Low | Moderate Exclusion of subjects with fall history | Low Subjects rested one minute between trials. | Low | Low | Low |
| Draganich (2004) | Low | Low | No information on status. | Low | Low | Low | Low |
| Eyal (2020) | Moderate | Low | Moderate Exclusion of subjects with fall history. | Low | Low | Low | Low |
| Hahn (2004) | Low | Low | Low | No information on randomization. No information on fatigue prevention strategy. | Low | Low 19 out of 6630 individual data points were removed because they were | Low |

| | | | | | | | |
|-----------------|----------|---------------------------|--|--|-----|----------------------------------|-----|
| | | | | | | “outside reasonable variability” | |
| Huang (2008) | Low | No information on gender. | No information on status or gender. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| Kim (2013) | Low | Low | Low | No information on fatigue prevention strategy. | Low | Low | Low |
| Kulkarni (2021) | Moderate | Low | Moderate Gender is unbalanced: 7 m, 24 f No information on status. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Low |
| LoJacono (2018) | Moderate | No information on gender. | No information on gender. | Low | Low | Low | Low |
| Lowrey (2007) | Low | Low | Low | Low | Low | Low | Low |
| Lu (2006) | Low | No information on gender. | No information on status or gender. | No information on fatigue prevention strategy. No information on randomization. | Low | Low | Low |
| Luo (2022) | Serious | Low | Serious Exclusion of: <ul style="list-style-type: none"> • subjects with fall history. • female subjects. | Low | Low | Low | Low |
| Maidana (2018) | Low | Low | Low | Low | Low | Low | Low |
| McFadyen (2002) | Moderate | Low | Moderate Exclusion of female subjects. | Low | Low | Low | Low |

| | | | | | | | |
|----------------------|----------|---------------------------|---|--|-----|---|---|
| Mckenzie (2004) | Moderate | Low | Moderate Gender is unbalanced: 12m, 20 f | No information on fatigue prevention strategy. | Low | Moderate 6 out of 17 older adults and 8 out of 15 younger adults were not analyzed due to technical failure. | Low |
| Park (2012) | Moderate | Low | Moderate Gender is unbalanced: 6 m, 12 f | Low | Low | Low | Low |
| Tomar (2019) | Serious | Low | Moderate Exclusion of female subjects. | No information on randomization. No information on fatigue prevention strategy. | Low | Low | Serious Significance level was not reported, and sample size was too low for the reported effect size. |
| Uchiyama (2012) | Low | No information on gender. | Low No information on gender. | Low | Low | Low | Low |
| Wang (2010) | Low | Low | Low | No information on fatigue prevention strategy. No information on randomization. | Low | Low | Low |
| Weerdesteijn (2005a) | Moderate | Low | Moderate Gender is unbalanced: 26 m, 98 f Exclusion of subjects without fall history. | No information on fatigue prevention strategy. | Low | Low 1% of trials were removed from the analysis because the subject performed an | Low |

| | | | | | | | |
|----------------------|---------|-----|---|-----|-----|------------------------------------|-----|
| | | | | | | “intermediate avoidance strategy”. | |
| Weerde steyn (2005b) | Serious | Low | Serious Exclusion of male subjects. Subjects were recruited from elderly sports groups. | Low | Low | Low | Low |