


Supplemental Figure 1: Mortality protected among 55-74 years age groups of IL1RN haplotype CTA-1/2 carriers.



SUPPLEMENTAL FIGURE 1: IL1RN CTA 1/2 Protective Haplotype is Associated with Decreased Mortality Among Men.
2A. Comparison of mortality rate among men between IL1RN CTA-1/2 vs. TTG-0/1/2 haplotype by decade.
2B. Mortality as percentage is shown for *IL1RN* CTA and TTG-0/1/2 carriers among men. Total number of patients are shown in parentheses for each age group.

Supplemental Table 1: Demography, inflammation biomarkers levels in COVID-19 patients.

	Number of observations	Plasma Levels Mean (SD)
Demography		
Age	2589	61.2 (18.7)
BMI	2532	30.43 (7.71)
Sex (% Male)	1380 (53.3%)	
Inflammation Markers		
LAB - IL-1Ra	465	2897.28 (2449.21)
LAB - IL1 β	642	7.02 (10.05)
LAB - IL2	645	6.92 (19.89)
LAB - IL6	1229	40.76 (161.30)
IL1 β - MAX	642	7.43 (13.64)
IL2 - MAX	645	7.31 (34.37)
IL6 - MAX	1174	107.10 (421.30)
LAB - CRP	2031	109.50 (86.96)
CRP - MAX	1984	152.00 (109.60)
LAB - PROCALCITONIN	1928	1.68 (17.50)
LAB - FERRITIN	2022	1232.0 (2583.0)
LAB – D-dimer	2046	1290.0 (3634.0)
D-dimer - MAX	2039	3277.0 (7282.0)
LAB - C3 COMPLEMENT	176	117.3 (39.82)
LAB - C4 COMPLEMENT	172	30.42 (14.91)
Mortality	397 (15.3%)	

IL-1Ra – interleukin-1 receptor antagonist, CRP – C - reactive protein, IL1 β – interleukin-1 beta, IL2 – interleukin-2, IL6 –interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. Biomarkers data are presented as mean $\pm$ standard deviation (SD).

Supplemental Table 2: Association of plasma inflammation cytokine biomarkers and mortality - with the *IL1RN* rs315952 CC genotype

	<i>IL1RN</i> rs315952 (N= 2589)		
	CC (N=237)	CT/TT (N=2352)	
Demography	Mean (SD)	Mean (SD)	FDR- p value
Age	61.09 (17.39)	61.21 (18.78)	0.9190
BMI	30.13 (7.75)	30.88 (21.15)	0.8955
Male %	90 (51.4%)	1290 (53.4%)	0.8955
Inflammation Markers			
IL-1Ra	2650 (2160)	2940 (2500)	0.4893
LAB - IL1	5.40 (0.86)	7.19 (10.54)	0.0007
LAB - IL2	4.84 (2.22)	7.14 (20.88)	0.0520
LAB - IL6	34.92 (77.32)	41.15 (167.57)	0.5168
IL1 - MAX	5.42 (0.87)	7.64 (14.32)	0.0007
IL2 - MAX	5.15 (4.88)	7.54 (36.08)	0.2581
IL6 - MAX	94.43 (224.73)	109.47 (438.63)	0.5500
LAB-CRP	119.88 (90.64)	108.65 (86.58)	0.2581
CRP - MAX	157.87 (110.06)	149.62 (110.13)	0.4839
LAB - PROCALCITONIN	3.96 (42.07)	1.41 (12.27)	0.4893
LAB - FERRITIN	1599.22 (3561.04)	1186.84 (2437.36)	0.2581
LAB – D-dimer	1932.25 (4818.88)	1233.02 (3495.84)	0.1820
D-dimer - MAX	4048.79 (8573.90)	3207.75 (7133.68)	0.3283
Mortality	40 (16.9%)	357 (15.2%)	1.11 (0.74, 1.63); 0.61

Summary of demographics and inflammation biomarkers by genotype. CRP – denotes – C - reactive protein, IL1 β – interleukin-1 beta, IL-2 – interleukin-2, IL6 – interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. Biomarkers are presented as mean $\pm$ standard deviation (SD). Adjusted p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For mortality comparison between genotypes - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for age, gender and BMI.

Supplemental Table 3: Association of plasma inflammation cytokine and mortality with the *IL1RN* rs9005 AA genotype

	<i>IL1RN</i> rs9005 (N= 2589)		
	AA (N=296)	AG/GG (N=2293)	
Demography	Mean (SD)	Mean (SD)	FDR- p value
Age	57.96 (17.95)	61.62 (18.71)	0.0030
BMI	30.75 (7.20)	30.40 (7.77)	0.4420
Male %	167 (56.4%)	1213 (52.9%)	0.4245
Inflammation Markers			
IL-1Ra	3810 (3640)	2800 (2270)	0.1207
LAB - IL1 β	5.80 (2.96)	7.19 (10.64)	0.0607
LAB - IL2	5.06 (2.10)	7.17 (21.17)	0.0715
LAB - IL6	52.75 (280.97)	39.05 (139.55)	0.7245
IL1 β - MAX	5.84 (2.67)	7.65 (14.49)	0.0520
IL2 - MAX	4.61 (2.21)	7.68 (36.60)	0.1062
IL6 - MAX	119.13 (390.96)	106.65 (426.85)	0.7963
LAB - CRP	108.61 (80.24)	109.81 (87.82)	0.8340
CRP - MAX	139.79 (102.60)	151.68 (110.97)	0.1690
LAB - PROCALCITONIN	0.56 (1.29)	1.79 (18.46)	0.0520
LAB - FERRITIN	1049.35 (1634.20)	1246.97 (2656.75)	0.1690
LAB - D-dimer	1177.93 (3766.41)	1310.93 (3623.26)	0.7245
D-dimer - MAX	2521.78 (5413.44)	3377.50 (7470.43)	0.0858
Mortality	Death = 35 (11.7%)	Death = 362 (15.8%)	0.87 (0.58, 1.27); 0.48

Summary of demographics and inflammation biomarkers by genotype. CRP – denotes – C - reactive protein, IL1 β – interleukin-1 beta, IL-2 – interleukin-2, IL6 – interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. Biomarkers are presented as mean $\pm$ standard deviation (SD). Adjusted p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For mortality comparison between genotypes - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for age, gender and BMI.

Supplemental Table 4: Comparison inflammation biomarkers between ≤80 and >80 years age group

	ALL (N=2589)		
	≤80 years (N=2145)	>80 Years (N=443)	
Demography	Mean (SD)	Mean (SD)	FDR – p value
Age	55.95 (15.96)	86.63 (4.33)	<0.001
BMI	31.11 (7.46)	27.16 (8.04)	<0.001
Gender (male %)	1185 (55.2%)	195 (44.0%)	<0.001
Inflammation Markers			
LAB - IL-1Ra	2751.45 (2313.42)	3706.57 (2988.51)	0.0124
LAB - IL1β	6.94 (9.23)	7.38 (13.29)	0.9630
LAB - IL2	6.80 (20.82)	7.49 (14.60)	0.9630
LAB - IL6	40.44 (157.38)	41.05 (176.82)	0.9630
IL1β - MAX	7.37 (13.69)	7.70 (13.42)	0.9630
IL2 - MAX	7.17 (36.89)	7.99 (17.86)	0.9630
IL6 - MAX	115.28 (457.68)	77.58 (222.28)	0.1450
LAB - CRP	111.96 (88.51)	100.24 (79.84)	0.0315
CRP - MAX	154.01 (112.57)	135.33 (98.08)	0.0124
LAB - PROCALCITONIN	1.86 (19.40)	0.80 (3.47)	0.1030
LAB – D-dimer	1156.80 (3377.45)	1876.81 (4530.08)	0.0124
D-dimer - MAX	3286.07 (7462.52)	3268.96 (6450.67)	0.9630
rs419598 (CC%)	144 (6.7%)	31 (7.2%)	0.9099
Mortality	251 (11.7%)	146 (33.0%)	3.92 (3.04, 5.04); < 0.001

Summary of demographics and inflammation biomarkers by genotype. CRP – denotes – C - reactive protein, IL1β – interleukin-1 beta, IL-2 – interleukin-2, IL6 – interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. Biomarkers are presented as mean ± standard deviation (SD). Adjusted p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For mortality comparison between age groups - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for gender and BMI.

Supplemental Table 5: Co-morbidities and mortality among ≤80 and >80 years age group

	≤80 years (N=2145)	>80 Years (N=443)	OR (95% CI); p value
Age (years)	56.0 (16.0)	86.6 (4.3)	0.0001
BMI	31.1 (7.46)	27.16 (8.04)	0.0001
Gender (Male %)	1185 (55.2%)	195 (44.1%)	0.0001
Metabolic Syndrome	248 (11.6%)	38 (8.6%)	1.28 (0.86 , 1.86) ; 0.212
CV	1299 (60.6%)	398 (89.8%)	7.65 (5.51 , 10.88) ; < 0.001
Pulm	345 (16.1%)	74 (16.7%)	1.23 (0.92 , 1.64) ; 0.161
Diabetes	724 (33.8%)	145 (32.7%)	1.21 (0.96 , 1.52) ; 0.103
Coronary Artery Disease	252 (11.7%)	139 (31.4%)	4.15 (3.21 , 5.37) ; < 0.001
Heart Failure	187 (8.7%)	108 (24.4%)	4.32 (3.25 , 5.75) ; < 0.001
Hyperlipdemia	828 (38.6%)	285 (64.3%)	3.49 (2.79 , 4.39) ; < 0.001
Hypertension	1075 (50.1%)	361 (81.5%)	5.59 (4.29 , 7.37) ; < 0.001
Peripheral Vascular Disease	131 (6.10%)	47 (10.6%)	2.02 (1.4 , 2.87) ; < 0.001
Asthma	259 (12.1%)	39 (8.8%)	0.82 (0.56 , 1.17) ; 0.279
COPD	130 (6.1%)	55 (12.4%)	2.5 (1.75 , 3.54) ; < 0.001
Dialysis	108 (5.0%)	13 (2.9%)	0.54 (0.28 , 0.96) ; 0.05
Cancer	196 (9.1%)	100 (22.6%)	2.93 (2.21 , 3.88) ; < 0.001
CKD	285 (13.3%)	68 (15.3%)	1.32 (0.97 , 1.77) ; 0.072
Mortality	251 (11.7%)	146 (33.0%)	3.92 (3.04 , 5.04) ; < 0.001

Age in years and BMI are presented as mean ± standard deviation (SD). Co-morbidities number of patients reported at admission and (%) over total number of subjects for each group is presented. For age, gender and BMI adjusted p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For comorbidities comparison between age groups - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for gender and BMI.

Supplemental Table 6: *IL1RN* rs419598, rs9005 and rs315952 variant genotypes and haplotypes frequency in COVID-19 patients.

RACE	Whites (N=1250)	Blacks (N=471)	East Asians (N=105)	South Asians (N=75)	Hispanic (N=419)
Total no. of subjects N=2589 (<i>IL1RN</i> - rs419598)					
CC	100	5	-	7	36
CT	504	65	16	21	174
TT	646	401	89	47	209
CC Genotype Frequency (%) NYU- COVID-19	8.0	1.1	0.0	9.3	8.6
1000 Genomes Project CC genotype Frequency (%)	8.2	0.2	0.6	9.2	11.8
(<i>IL1RN</i> – rs315952)					
CC	68	93	28	7	25
CT	460	230	52	19	180
TT	722	148	25	49	214
CC Genotype Frequency (%) NYU- COVID-19	5.4	19.7	26.7	9.3	6.0
1000 Genomes Project CC genotype Frequency (%)	9.9	21.2	30.6	2.9	4.9
(<i>IL1RN</i> - rs9005)					
AA	138	14	10	5	85
AG	542	152	50	37	198
GG	570	305	45	33	136
AA Genotype Frequency (%) NYU- COVID-19	11.0	2.96	9.5	6.7	20.3
1000 Genomes Project AA genotype Frequency (%)	8.7	2.9	10.9	15.7	26.8
<i>IL1RN</i> haplotype (rs419598, rs315952 and rs9005)					
CTA-1	56 (4.5%)	2 (0.4%)	5 (4.8%)	3	38 (9.1%)
CTA-2	80 (6.4%)	3 (0.6%)	0 (0.0%)	4	29 (6.9%)
TTG-1	523 (41.8%)	251 (53.3%)	61 (58.1%)	32	204 (48.7%)
TTG-2	246 (19.7%)	90 (19.2%)	2 (1.9%)	16	35 (8.4%)
CTA-1/TTG-1	257 (20.6%)	27 (5.7%)	3 (2.9%)	18	64 (15.3%)
CTA-0/TTG-0	88 (7.0%)	98 (20.8%)	33 (31.4%)	2	49 (11.7%)

Supplemental Table 7. Comparison of biomarkers - IL1RN - rs419598 CC vs. CT/TT – Whites only

	IL1RN rs419598 (N=1250)		
	CC (N=100)	CT/TT (N=1150)	
Demography	Mean (SD)	Mean (SD)	FDR – p value
Age	62.95 (20.73)	64.46 (19.50)	0.4830
BMI	30.66 (7.55)	30.07 (8.00)	0.4830
Male %	44 (44.0%)	599 (52.1%)	0.4440
Inflammation Markers			
LAB - IL-1Ra	3790 (4010)	3130 (2620)	0.4670
LAB - IL1 β	5.40 (0.89)	7.72 (13.16)	0.0153
LAB - IL2	4.42 (1.19)	6.32 (10.07)	0.0153
LAB - IL6	22.57 (24.88)	38.21 (153.85)	0.0579
IL1 β - MAX	5.47 (0.92)	8.00 (13.76)	0.0153
IL2 - MAX	4.28 (1.29)	6.45 (12.20)	0.0160
IL6 - MAX	37.32 (69.39)	96.27 (476.65)	0.0295
LAB - CRP	89.76 (70.45)	109.66 (89.34)	0.0416
CRP - MAX	121.70 (96.40)	149.32 (110.05)	0.0416
LAB - PROCALCITONIN	0.50 (1.12)	0.87 (4.74)	0.0826
LAB - FERRITIN	827.15 (1115.86)	1155.42 (2896.63)	0.0579
LAB - D-dimer	827.82 (1637.43)	1223.58 (3107.91)	0.0798
D-dimer - MAX	1629.51 (2703.57)	2908.58 (6138.77)	0.0013
Mortality	10 (10.0%)	201 (17.6%)	0.54 (0.25 - 1.03); 0.081
Mortality - Men only	5 (11.4%)	120 (20.0%)	0.44 (0.14 - 1.08); 0.099
Mortality - Women only	5 (8.9%)	81 (14.7%)	0.70 (0.23 - 1.74); 0.478

Summary of demographics and inflammation biomarkers by genotype. *IL1RN* – interleukin-1 receptor antagonist gene, CRP – C - reactive protein, IL1 β – interleukin-1 beta, IL-2 – interleukin-2, IL6 –interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. Biomarkers data are presented as mean $\pm$ standard deviation (SD). Adjusted p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For overall mortality comparison between genotype - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for age, gender and BMI ; and for age and BMI for gender specific mortality.

Supplemental Table 8: White patients with the *IL1RN* CTA haplotype carriers exhibit lower plasma inflammation cytokine levels and mortality.

	IL1RN Whites Only (N=1250)					
	CTA-1/2 (N=137)	TTG-(0/1/2) (N=856)	CTA-1/TTG-1 (N=257)	CTA-1/2 Vs. TTG-0/1/2	TTG-0/1/2 vs. CTA 1/TTG1	CTA-1/2 vs. CTA-1/TTG-1
Demography	Mean (SD)	Mean (SD)	Mean (SD)	FDR - p value	FDR - p value	FDR - p value
Age	61.75 (19.18)	64.60 (19.59)	64.86 (19.76)	0.3270	0.8550	0.3930
BMI	30.35 (6.64)	29.91 (7.40)	30.67 (10.10)	0.4860	0.8070	0.7830
Gender (Male %)	66 (48.2%)	448 (52.3%)	129 (50.2%)	0.4860	0.8550	0.7830
Inflammation Markers						
IL-1Ra	4440 (4090)	2840 (2120)	4120 (3990)	0.1011	0.4977	0.9380
LAB - IL1β	6.39 (4.05)	8.04 (14.87)	6.52 (5.19)	0.2069	0.4977	0.9380
LAB - IL2	4.93 (2.77)	6.43 (10.91)	6.03 (7.31)	0.1413	0.7450	0.6419
LAB - IL6	28.51 (51.58)	40.54 (170.09)	29.33 (85.70)	0.2640	0.5558	0.9380
IL1β - MAX	6.58 (4.01)	8.31 (15.52)	6.81 (5.69)	0.2069	0.4977	0.9380
IL2 - MAX	4.57 (2.93)	6.33 (11.22)	7.16 (15.95)	0.1011	0.7450	0.6419
IL6 - MAX	49.81 (83.69)	105.95 (546.06)	66.92 (134.65)	0.1011	0.4977	0.6419
LAB - CRP	94.68 (74.69)	111.59 (89.26)	103.44 (90.08)	0.1011	0.4977	0.6419
CRP - MAX	123.93 (94.84)	151.71 (108.40)	144.24 (117.38)	0.0338	0.6175	0.6419
LAB - PROCALCITONIN	0.59 (1.12)	0.92 (5.30)	0.72 (2.34)	0.2069	0.6175	0.7323
LAB - FERRITIN	854.16 (1076.04)	1192.58 (2918.45)	1060.03 (2995.67)	0.0978	0.6949	0.6419
LAB - D-dimer	759.87 (1434.46)	1309.51 (3170.48)	1023.45 (3077.92)	0.0228	0.4977	0.6419
D-dimer - MAX	1681.82 (3178.80)	3076.46 (6492.51)	2491.68 (4971.28)	0.0013	0.4977	0.6419
Mortality (ALL)	15 (10.9%)	155 (18.1%)	41 (15.9%)	0.66 (0.35 - 1.15); 0.163	1.18 (0.80 - 1.77); 0.415	0.78 (0.39 - 1.50); 0.469
Mortality (Men only)	7 (10.6%)	93 (20.7%)	25 (19.3%)	0.51 (0.20 - 1.12); 0.117	1.13 (0.68 - 1.93); 0.633	0.61 (0.21 - 1.58); 0.327
Mortality (women only)	8 (11.3%)	62 (15.2%)	16 (12.5%)	0.87 (0.36 - 1.92); 0.751	1.28 (0.70 - 2.46); 0.443	1.20 (0.43 - 3.17); 0.717

Summary of demographics and inflammation biomarkers by haplotypes. Biomarkers are presented as mean ± standard deviation (SD). CRP – denotes – C - reactive protein, IL1β – interleukin-1 beta, IL2- interleukin-2, IL6 – interleukin-6. The body mass index (BMI) is the weight in kilograms divided by the square of the height in meters. FDR p-values control for false discovery rate at 5% using Benjamin Hochberg criteria within each subgroup of biomarkers. For mortality comparison between haplotypes - odds ratio [aOR (95% Confidence intervals)] and p values derived from multivariable logistic regression adjusting for age, gender and BMI; and for age and BMI for gender specific mortality.