

Supplementary Material

Supplementary Tables

Table S1 Demographics and frailty indicators for the participants (N= 44,520), separated by gender. Values are reported as counts (N).

	Full cohort (N = 44,520)	Female (N = 23,013)	Male (N = 21,507)
Weight loss (N)	7,840	3,873	3,967
Exhaustion (N)	3,467	2,146	1,321
Physical activity (N)			
Low physical activity	150	98	52
Activity less than once a week	12,925	6,859	6,066
Walking pace (N)			
Slow	2,007	1,095	912
Steady	21,275	10,838	10,437
Brisk	19,346	10,049	9,297
Low grip strength* (N)	7,712	3,897	3,815

* Definition used by Fried et al.[1].

Table S2 Summary statistics for skeletal muscle and fat IDPs. Values are reported as mean and standard deviation. Significance refers to the p-value for a Wilcoxon rank-sums test, where the null hypothesis is the medians between the two groups (male and female participants) being equal.

	Full cohort (N = 44,520)	Female (N = 23,013)	Male (N = 21,507)	p-values
Muscle volume IDPs				
Left Thigh Muscle Volume (L)	4.22 ± 1.12	3.34 ± 0.51	5.16 ± 0.77	$p < 10^{-16}**$
Right Thigh Muscle Volume (L)	4.23 ± 1.13	3.35 ± 0.51	5.18 ± 0.78	$p < 10^{-16}**$
L-R Thigh Muscle Volume Difference (L)	-0.01 ± 0.22	-0.004 ± 0.17	-0.02 ± 0.27	$p < 10^{-16}**$
Left Mid-thigh Muscle Volume (L)	1.13 ± 0.27	0.93 ± 0.14	1.35 ± 0.21	$p < 10^{-16}**$
Right Mid-thigh Muscle Volume (L)	1.15 ± 0.28	0.94 ± 0.14	1.38 ± 0.21	$p < 10^{-16}**$
L-R Mid-thigh Muscle Volume Difference (L)	-0.02 ± 0.07	-0.01 ± 0.06	-0.03 ± 0.08	$p < 10^{-16}**$

Left Iliopsoas Muscle Volume (L)	0.31 ± 0.09	0.25 ± 0.04	0.39 ± 0.06	$p < 10^{-16}**$
Right Iliopsoas Muscle Volume (L)	0.33 ± 0.09	0.26 ± 0.04	0.40 ± 0.06	$p < 10^{-16}**$
L-R Iliopsoas Muscle Volume Difference (L)	-0.01 ± 0.02	-0.01 ± 0.02	-0.01 ± 0.02	0.00011*
Muscle quality IDPs				
Left Thigh IMAT Volume (L)	0.39 ± 0.17	0.36 ± 0.15	0.42 ± 0.19	$p < 10^{-16}**$
Right Thigh IMAT Volume (L)	0.39 ± 0.17	0.35 ± 0.15	0.42 ± 0.19	$p < 10^{-16}**$
L-R Thigh IMAT Volume Difference (L)	0.00031 ± 0.05	0.0041 ± 0.04	-0.0038 ± 0.05	$p < 10^{-16}**$
Left Mid-thigh IMAT Volume (ml)	25.69 ± 22.57	24.16 ± 18.95	27.32 ± 25.79	$p < 10^{-16}**$
Right Mid-thigh IMAT Volume (ml)	25.88 ± 24.11	24.03 ± 20.78	27.87 ± 27.10	$p < 10^{-16}**$
L-R Mid-thigh IMAT Volume Difference (ml)	-0.19 ± 13.64	0.13 ± 10.35	-0.54 ± 16.45	$p < 10^{-16}**$
Thigh SAT Volume (L)	7.48 ± 3.03	8.98 ± 2.98	5.87 ± 2.14	$p < 10^{-16}**$
Thigh SAT Volume Index (L/m²)	2.67 ± 1.20	3.39 ± 1.11	1.89 ± 0.68	$p < 10^{-16}**$
Left Thigh SAT Volume (L)	3.74 ± 1.52	4.48 ± 1.50	2.94 ± 1.08	$p < 10^{-16}**$
Right Thigh SAT Volume (L)	3.74 ± 1.55	4.51 ± 1.53	2.93 ± 1.08	$p < 10^{-16}**$
L-R Thigh SAT Volume Difference (L)	-0.0089 ± 0.44	-0.03 ± 0.53	0.01 ± 0.31	$p < 10^{-16}**$
Mid-thigh SAT Volume (L)	1.67 ± 0.75	2.07 ± 0.76	1.25 ± 0.46	$p < 10^{-16}**$
Left Mid-thigh SAT Volume (ml)	0.83 ± 0.37	1.02 ± 0.38	0.63 ± 0.23	$p < 10^{-16}**$
Right Mid-thigh SAT Volume (L)	0.84 ± 0.39	1.04 ± 0.39	0.62 ± 0.24	$p < 10^{-16}**$
L-R Mid-thigh SAT Volume Difference (L)	-0.0063 ± 0.10	-0.02 ± 0.12	0.007 ± 0.08	$p < 10^{-16}**$

IMAT: Intermuscular adipose tissue; SAT: Subcutaneous adipose tissue. * indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

Table S3 Demographics of the participants (N=44,520), separated by self-reported ethnicity and gender. Values are reported as mean and standard deviation for continuous variables and counts (N) for categorical variables.

	White Female (N = 22,227)	White Male (N = 20,782)	Asian Female (N = 175)	Asian Male (N = 301)	Black Female (N = 164)	Black Male (N = 133)	Chinese Female (N = 80)	Chinese Male (N = 53)	Others Female (N = 265)	Others Male (N = 173)
Age (yrs.)	63.64 ± 7.55	65.06 ± 7.78	60.93 ± 8.19	61.32 ± 8.61	59.51 ± 7.24	58.88 ± 7.46	59.55 ± 6.96	60.45 ± 6.57	60.80 ± 7.79	61.73 ± 8.41
Weight (kg)	68.88 ± 12.94	83.75 ± 13.33	64.63 ± 10.49	76.15 ± 11.64	77.70 ± 16.70	86.24 ± 14.67	57.23 ± 8.87	70.04 ± 9.42	68.95 ± 15.31	81.99 ± 13.75
Height (cm)	162.81 ± 6.21	176.18 ± 6.60	157.95 ± 6.51	170.90 ± 6.08	162.59 ± 7.27	175.67 ± 6.49	157.26 ± 5.10	168.79 ± 5.11	160.43 ± 7.02	173.23 ± 7.11
BMI (kg/m²)	25.99 ± 4.70	26.96 ± 3.90	25.94 ± 4.08	26.03 ± 3.50	29.43 ± 6.16	27.85 ± 3.91	23.12 ± 3.23	24.58 ± 3.15	26.73 ± 5.28	27.27 ± 4.06
Waist circumference (cm)	82.77 ± 11.80	94.45 ± 10.71	82.90 ± 10.94	90.93 ± 9.61	88.04 ± 13.85	91.98 ± 10.24	75.11 ± 9.15	85.14 ± 9.34	82.37 ± 12.69	92.68 ± 11.39
Hip circumference (cm)	100.83 ± 9.79	100.79 ± 7.31	98.73 ± 8.22	97.13 ± 7.69	105.23 ± 11.62	100.72 ± 7.50	91.98 ± 7.08	93.60 ± 6.28	100.21 ± 11.40	98.99 ± 7.69
Waist to Hip Ratio	0.82 ± 0.07	0.94 ± 0.06	0.84 ± 0.07	0.94 ± 0.06	0.83 ± 0.07	0.91 ± 0.06	0.82 ± 0.06	0.91 ± 0.06	0.82 ± 0.07	0.94 ± 0.07
Dominant HGS (kg)	23.98 ± 6.10	38.87 ± 8.82	22.05 ± 7.03	34.55 ± 8.89	26.84 ± 7.01	43.89 ± 10.43	21.83 ± 5.21	36.44 ± 7.93	24.57 ± 6.68	37.76 ± 8.43
Walking MET (hours/week)	12.57 ± 17.00	12.23 ± 16.17	10.21 ± 12.54	9.87 ± 12.25	8.61 ± 14.80	8.56 ± 12.57	10.04 ± 12.77	9.69 ± 9.99	13.37 ± 17.76	9.98 ± 13.63
Moderate MET (hours/week)	12.70 ± 18.97	12.82 ± 19.64	11.08 ± 14.92	8.65 ± 13.72	8.63 ± 13.98	8.52 ± 13.16	12.90 ± 21.03	6.59 ± 12.36	11.82 ± 16.29	10.17 ± 16.70
Vigorous MET (hours/week)	8.44 ± 16.11	11.04 ± 19.36	8.34 ± 13.55	10.14 ± 15.70	9.24 ± 14.76	11.89 ± 17.79	5.49 ± 10.75	8.22 ± 11.39	10.33 ± 16.59	10.39 ± 16.64

Total MET (hours/week)	33.71 ± 39.00	36.09 ± 42.08	29.63 ± 31.83	28.66 ± 33.95	26.48 ± 32.02	28.98 ± 36.15	28.43 ± 33.43	24.50 ± 23.17	35.53 ± 39.60	30.53 ± 35.17
Townsend deprivation score	-1.89 ± 2.69	-1.98 ± 2.69	-0.84 ± 2.97	-1.01 ± 3.13	1.33 ± 3.56	1.19 ± 3.69	-2.09 ± 2.50	-1.67 ± 2.85	0.32 ± 3.40	-0.06 ± 3.35
Alcohol frequency (N)										
Daily	3,081	4,387	11	27	7	13	2	4	14	23
1-4 times per week	11,583	11,948	36	93	41	51	11	12	103	76
1-3 times per month	2,969	1,953	18	33	30	18	12	14	46	23
Occasionally or never	4,484	2,350	105	145	83	48	55	23	99	51
Smoking status (N)										
Never	14,491	11,981	155	219	125	77	72	37	161	97
Previous	6,913	7,799	15	62	28	41	6	12	88	62
Current	632	810	1	15	8	13	1	4	12	13
VAT (L)	2.83 ± 1.57	5.19 ± 2.37	2.68 ± 1.20	4.50 ± 1.72	2.26 ± 1.37	3.41 ± 1.73	2.07 ± 1.07	3.29 ± 1.59	2.62 ± 1.49	4.88 ± 2.42
ASAT (L)	9.81 ± 4.36	7.02 ± 3.27	10.12 ± 3.45	7.32 ± 3.01	11.70 ± 5.24	6.65 ± 3.61	6.97 ± 2.73	5.18 ± 2.13	10.10 ± 4.71	7.32 ± 3.48

BMI: Body mass index; HGS: Hand grip strength; MET: Metabolic equivalent task; PA: Physical activity; VAT: Visceral adipose tissue; ASAT: Abdominal subcutaneous adipose tissue.

Table S4 Summary statistics for skeletal muscle fat IDPs for each self-reported ethnicity and each gender. Values are reported as mean and standard deviation.

	White Female (N = 22,227)	White Male (N = 20,782)	Asian Female (N = 175)	Asian Male (N = 301)	Black Female (N = 164)	Black Male (N = 133)	Chinese Female (N = 80)	Chinese Male (N = 53)	Others Female (N = 265)	Others Male (N = 173)
Muscle volume IDPs										
Total Muscle Volume (L)	14.0 ± 1.9	21.8 ± 3.0	12.5 ± 1.8	19.3 ± 2.9	16.0 ± 2.6	24.9 ± 4.0	12.5 ± 1.7	19.4 ± 2.1	14.1 ± 2.4	21.5 ± 3.4
Total Muscle Volume Index (L/m²)	5.3 ± 0.6	7.0 ± 0.8	5.0 ± 0.6	6.6 ± 0.8	6.1 ± 0.8	8.1 ± 1.1	5.1 ± 0.6	6.8 ± 0.7	5.5 ± 0.7	7.1 ± 0.9
Thigh Muscle Volume (L)	6.7 ± 1.0	10.3 ± 1.5	6.0 ± 1.0	9.2 ± 1.5	7.9 ± 1.4	12.3 ± 2.1	6.0 ± 0.9	9.2 ± 1.0	6.8 ± 1.2	10.3 ± 1.7
Thigh Muscle Volume Index (L/m²)	2.5 ± 0.3	3.3 ± 0.4	2.4 ± 0.3	3.2 ± 0.4	3.0 ± 0.4	4.0 ± 0.6	2.4 ± 0.3	3.2 ± 0.3	2.6 ± 0.4	3.4 ± 0.5
Mid-thigh Muscle Volume (L)	1.9 ± 0.3	2.7 ± 0.4	1.7 ± 0.3	2.5 ± 0.4	2.3 ± 0.4	3.3 ± 0.6	1.8 ± 0.2	2.5 ± 0.3	2.0 ± 0.3	2.8 ± 0.5
Iliopsoas Muscle Volume (L)	0.5 ± 0.1	0.8 ± 0.1	0.4 ± 0.1	0.7 ± 0.1	0.5 ± 0.1	0.9 ± 0.1	0.4 ± 0.1	0.7 ± 0.1	0.5 ± 0.1	0.8 ± 0.1
Iliopsoas Muscle Volume Index (L/m²)	0.2 ± 0.0	0.3 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.3 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.3 ± 0.0
Muscle quality IDPs										
Thigh IMAT Volume	0.7 ± 0.3	0.8 ± 0.4	0.7 ± 0.3	0.8 ± 0.3	0.8 ± 0.4	0.8 ± 0.4	0.5 ± 0.2	0.6 ± 0.2	0.7 ± 0.3	0.8 ± 0.3

(L)										
Thigh IMAT Volume Index (ml/m²)	266.2 ± 109.0	273.4 ± 120.8	287.1 ± 107.2	281.2 ± 105.6	323.3 ± 143.0	273.2 ± 113.3	208.5 ± 72.8	193.9 ± 80.5	264.7 ± 112.3	268.1 ± 112.1
Mid-thigh IMAT Volume (ml)	48.0 ± 38.2	55.4 ± 50.6	56.1 ± 36.3	51.8 ± 35.4	72.6 ± 62.7	55.6 ± 52.7	33.0 ± 21.4	27.8 ± 23.1	48.0 ± 35.8	51.4 ± 42.6
Mid-thigh IMAT/Muscle (%)	2.5 ± 1.9	2.0 ± 1.8	3.1 ± 2.0	2.1 ± 1.4	3.1 ± 2.4	1.7 ± 1.4	1.8 ± 1.1	1.1 ± 0.9	2.4 ± 1.6	1.8 ± 1.5
Paraspinal PDFF (%)	7.8 ± 4.0	6.9 ± 3.6	9.1 ± 4.6	7.0 ± 3.9	7.7 ± 3.9	5.3 ± 2.8	6.4 ± 2.8	5.0 ± 2.5	7.6 ± 3.9	6.8 ± 4.4
Paraspinal Iron concentration (mg/g)	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.1	1.2 ± 0.2	1.2 ± 0.2	1.2 ± 0.1

IMAT: Intermuscular adipose tissue; SAT: Subcutaneous adipose tissue; FTMR: Fat to muscle ratio.

Table S5 Summary of regression coefficients for linear regression models of the muscle and fat IDPs adjusted for baseline characteristics. Standard errors are in parentheses.

	Total Muscle Volume	Total Muscle Volume Index	Thigh Muscle Volume	Thigh Muscle Volume Index	Mid-thigh Muscle Volume	Iliopsoas Muscle Volume	Iliopsoas Muscle Volume Index	Thigh IMAT Volume	Thigh IMAT Volume Index	Mid-thigh IMAT Volume	Mid-thigh IMAT/Muscle	Paraspinal Muscle PDFF	Paraspinal Muscle Iron Concentration
Age	-0.137** (0.002)	-0.134** (0.003)	-0.135** (0.002)	-0.132** (0.003)	-0.185** (0.003)	-0.159** (0.003)	-0.160** (0.003)	0.141** (0.004)	0.171** (0.004)	0.099** (0.004)	0.145** (0.004)	0.220** (0.004)	-0.083** (0.005)
Male	1.371** (0.007)	1.322** (0.008)	1.306** (0.008)	1.238** (0.009)	1.246** (0.008)	1.386** (0.009)	1.330** (0.01)	0.120** (0.013)	-0.140** (0.012)	-0.048* (0.014)	-0.385** (0.014)	-0.536** (0.014)	-0.074** (0.017)
Asian	-0.395** (0.02)	-0.261** (0.023)	-0.337** (0.022)	-0.180** (0.025)	-0.269** (0.023)	-0.501** (0.024)	-0.416** (0.029)	0.148** (0.036)	0.287** (0.035)	0.152* (0.038)	0.261** (0.04)	0.268** (0.039)I	0.119* (0.048)
Black	0.293** (0.026)	0.506** (0.029)	0.462** (0.029)	0.764** (0.032)	0.588** (0.029)	-0.025 (0.031)	0.042 (0.037)	0.030 (0.045)	0.075 (0.044)	0.097* (0.049)	-0.019 (0.051)	-0.254** (0.049)	-0.191* (0.061)
Chinese	-0.259** (0.039)	0.031 (0.043)	-0.233** (0.043)	0.070 (0.047)	-0.054 (0.043)	-0.355** (0.046)	-0.121* (0.055)	-0.170* (0.068)	-0.046 (0.066)	0.0002 (0.073)	-0.025 (0.077)	-0.028 (0.074)	0.036 (0.09)
Others	-0.076* (0.021)	0.058* (0.023)	-0.031 (0.023)	0.120** (0.026)	0.054* (0.023)	-0.186** (0.025)	-0.108* (0.03)	-0.097* (0.037)	-0.042 (0.036)	-0.078* (0.039)	-0.087* (0.042)	-0.05 (0.04)	0.122* (0.049)
BMI	0.239** (0.002)	0.385** (0.003)	0.220** (0.003)	0.361** (0.003)	0.311** (0.003)	0.187** (0.003)	0.301** (0.003)	0.571** (0.004)	0.619** (0.004)	0.551** (0.005)	0.449** (0.005)	0.423** (0.005)	-0.085** (0.006)
Waist-to-Hip ratio	-0.028** (0.003)	-0.049** (0.004)	-0.017** (0.003)	-0.035** (0.004)	-0.018** (0.004)	-0.049** (0.004)	-0.078** (0.005)	0.106** (0.006)	0.095** (0.006)	0.102** (0.006)	0.100** (0.006)	0.195** (0.006)	0.077** (0.007)
	0.058**	0.063**	0.078**	0.091**	0.096**	0.037**	0.033*	0.226**	0.219**	0.169**	0.138**	0.068**	-0.093**

Alcohol frequency [Daily]	(0.007)	(0.008)	(0.008)	(0.009)	(0.008)	(0.009)	(0.01)	(0.013)	(0.012)	(0.014)	(0.014)	(0.014)	(0.017)
Alcohol frequency [1-4 times per week]	0.061** (0.006)	0.075** (0.007)	0.077** (0.006)	0.099** (0.007)	0.100** (0.007)	0.066** (0.007)	0.080** (0.008)	0.094** (0.01)	0.088** (0.01)	0.077** (0.011)	0.045* (0.012)	0.018 (0.011)	-0.046* (0.014)
Alcohol frequency [1-3 times per month]	0.022* (0.008)	0.027* (0.009)	0.030* (0.009)	0.040** (0.01)	0.041** (0.009)	0.033* (0.009)	0.042* (0.011)	0.020 (0.014)	0.014 (0.014)	0.021 (0.015)	0.008 (0.016)	-0.033* (0.015)	-0.002 (0.019)
Smoking status [Previous]	-0.002 (0.004)	-0.010* (0.005)	-0.014* (0.005)	-0.029** (0.006)	-0.018* (0.005)	-0.028** (0.005)	-0.046** (0.006)	0.084** (0.008)	0.077** (0.008)	0.054** (0.008)	0.055** (0.009)	0.014 (0.009)	-0.021* (0.011)
Smoking status [Current]	-0.023* (0.011)	-0.036* (0.013)	-0.080** (0.013)	-0.113** (0.014)	-0.097** (0.013)	-0.065** (0.014)	-0.095** (0.016)	0.161** (0.02)	0.160** (0.02)	0.092** (0.022)	0.116** (0.023)	0.058* (0.022)	0.057* (0.027)
Townsend deprivation index	-0.016** (0.002)	-0.012** (0.002)	-0.017** (0.002)	-0.014** (0.003)	-0.017** (0.002)	-0.020** (0.003)	-0.018** (0.003)	0.014* (0.004)	0.021** (0.004)	0.018** (0.004)	0.021** (0.004)	0.018** (0.004)	-0.001 (0.005)
Dominant HGS	0.225** (0.003)	0.155** (0.003)	0.233** (0.003)	0.171** (0.004)	0.206** (0.003)	0.198** (0.004)	0.119** (0.004)	-0.019* (0.005)	-0.082** (0.005)	-0.060** (0.006)	-0.109** (0.006)	-0.076** (0.006)	-0.018* (0.007)
MET	0.040** (0.002)	0.077** (0.002)	0.039** (0.002)	0.076** (0.003)	0.054** (0.002)	0.020** (0.002)	0.048** (0.003)	-0.050** (0.004)	-0.043** (0.004)	-0.028** (0.004)	-0.041** (0.004)	-0.015* (0.004)	-0.001 (0.005)
Constant	-0.703** (0.006)	-0.696** (0.007)	-0.681** (0.007)	-0.669** (0.008)	-0.670** (0.007)	-0.695** (0.008)	-0.677** (0.009)	-0.186** (0.011)	-0.057** (0.011)	-0.075** (0.012)	0.111** (0.012)	0.230** (0.012)	0.078** (0.015)
Observations	42,250	42,207	42,248	42,181	42,245	42,250	42,207	42,248	42,205	42,245	42,245	42,037	42,037
Adjusted R²	0.823	0.778	0.784	0.733	0.777	0.745	0.639	0.449	0.477	0.372	0.298	0.341	0.013
AIC	46984.6	56347.4	55281.3	64062.3	56589.5	62406.8	76838.5	94579.5	92403.5	100340.1	105079	101242.2	118262.7

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

Table S6 Regression coefficients between total muscle volume and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Total Muscle Volume			
	(1)	(2)	(3)	(4)
Age	-0.026** (0.0003)	-0.023** (0.0003)	-0.027** (0.0003)	-0.024** (0.0003)
	1.532** (0.005)	1.817** (0.005)	1.532** (0.006)	1.825** (0.005)
Male	0.079** (0.001)		0.081** (0.001)	
		0.037** (0.001)		0.038** (0.001)
Sarcopenia	-0.234** (0.009)	-0.238** (0.009)		
			-0.202** (0.023)	-0.217** (0.023)
Frailty [Frail]			-0.073** (0.005)	-0.081** (0.005)
Constant	-0.721** (0.004)	-0.859** (0.003)	-0.704** (0.004)	-0.842** (0.004)
Observations	42,785	42,785	41,417	41,417
Adjusted R²	0.770	0.766	0.770	0.766
AIC	58572.6	59379.5	56775.7	57548.9

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S7 Regression coefficients between total muscle volume index and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Total Muscle Volume Index			
	(1)	(2)	(3)	(4)
Age	-0.026**	-0.023**	-0.027**	-0.023**

	(0.0004)	(0.0004)	(0.0004)	(0.0004)
Male	1.352** (0.007)	1.706** (0.006)	1.360** (0.007)	1.709** (0.006)
Visceral fat	0.096** (0.001)		0.095** (0.001)	
Abdominal subcutaneous fat		0.047** (0.001)		0.047** (0.001)
Sarcopenia	-0.166** (0.011)	-0.173** (0.011)		
Frailty [Frail]			-0.096* (0.027)	-0.122** (0.028)
Frailty [Pre Frail]			-0.008 (0.006)	-0.021* (0.006)
Constant	-0.643** (0.004)	-0.813** (0.004)	-0.652** (0.005)	-0.815** (0.005)
Observations	42,742	42,742	41,376	41,376
Adjusted R²	0.666	0.664	0.667	0.664
AIC	74463.2	74768.5	71942.3	72289.1

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S8 Regression coefficients between thigh muscle volume and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Thigh Muscle Volume			
	(1)	(2)	(3)	(4)
Age	-0.026** (0.0003)	-0.023** (0.0003)	-0.027** (0.0003)	-0.024** (0.0003)
Male	1.506** (0.006)	1.769** (0.005)	1.504** (0.006)	1.779** (0.005)
Visceral fat	0.072** (0.001)		0.075** (0.001)	

		0.035**		0.037**
Abdominal subcutaneous fat		(0.001)		(0.001)
	-0.245**	-0.249**		
Sarcopenia	(0.01)	(0.01)		
			-0.250**	-0.270**
Frailty [Frail]			(0.025)	(0.025)
			-0.088**	-0.099**
Frailty [Pre Frail]			(0.005)	(0.005)
	-0.708**	-0.835**	-0.683**	-0.812**
Constant	(0.004)	(0.004)	(0.004)	(0.004)
Observations	42,783	42,783	41,415	41,415
Adjusted R²	0.732	0.730	0.733	0.731
AIC	65153.9	65488.2	62973.7	63261.7

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S9 Regression coefficients between thigh muscle volume index and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Thigh Muscle Volume Index			
	(1)	(2)	(3)	(4)
Age	-0.026** (0.0004)	-0.023** (0.0004)	-0.027** (0.0004)	-0.023** (0.0004)
Male	1.327** (0.007)	1.653** (0.006)	1.332** (0.007)	1.659** (0.006)
Visceral fat	0.086** (0.001)		0.087** (0.002)	
Abdominal subcutaneous fat		0.045** (0.001)		0.046** (0.001)
Sarcopenia	-0.186** (0.011)	-0.193** (0.011)		
Frailty [Frail]			-0.164** -0.198**	

		(0.029)	(0.029)
Frailty [Pre Frail]		-0.032**	-0.048**
		(0.006)	(0.006)
Constant	-0.630** (0.005)	-0.786** (0.004)	-0.627** (0.005)
Observations	42,715	42,715	41,353
Adjusted R²	0.625	0.626	0.626
AIC	79384.6	79227.4	76720.5
			76592.1

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S10 - Regression coefficients between mid-thigh muscle volume and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Mid-thigh Muscle Volume			
	(1)	(2)	(3)	(4)
Age	-0.033** (0.0003)	-0.029** (0.0003)	-0.034** (0.0004)	-0.030** (0.0004)
Male	1.360** (0.006)	1.728** (0.006)	1.359** (0.006)	1.736** (0.006)
Visceral fat	0.102** (0.001)		0.104** (0.001)	
Abdominal subcutaneous fat		0.048** (0.001)		0.050** (0.001)
Sarcopenia	-0.224** (0.01)	-0.230** (0.01)		
Frailty [Frail]			-0.227** (0.026)	-0.249** (0.026)
Frailty [Pre Frail]			-0.065** (0.006)	-0.077** (0.006)
Constant	-0.639** (0.004)	-0.817** (0.004)	-0.623** (0.005)	-0.800** (0.004)

Observations	42,779	42,779	41,412	41,412
Adjusted R²	0.704	0.698	0.705	0.699
AIC	69370.6	70271.2	67041.8	67888

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S11 - Regression coefficients between iliopsoas muscle and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Iliopsoas Muscle Volume			
	(1)	(2)	(3)	(4)
Age	-0.028** (0.0003)	-0.026** (0.0003)	-0.029** (0.0003)	-0.026** (0.0003)
	1.492** (0.006)	1.722** (0.006)	1.486** (0.006)	1.732** (0.006)
Male	0.068** (0.001)		0.072** (0.001)	
Visceral fat				
Abdominal subcutaneous fat		0.027** (0.001)		0.029** (0.001)
Sarcopenia	-0.208** (0.01)	-0.208** (0.01)		
Frailty [Frail]			-0.266** (0.025)	-0.263** (0.026)
Frailty [Pre Frail]			-0.094** (0.006)	-0.096** (0.006)
Constant	-0.704** (0.004)	-0.815** (0.004)	-0.673** (0.005)	-0.791** (0.004)
Observations	42,785	42,785	41,417	41,417
Adjusted R²	0.712	0.704	0.714	0.707
AIC	68322.3	69368.2	65807.5	66857.4

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,

- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S12 - Regression coefficients between iliopsoas muscle index and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Iliopsoas Muscle Volume Index			
	(1)	(2)	(3)	(4)
Age	-0.028** (0.0004)	-0.025** (0.0004)	-0.028** (0.0004)	-0.026** (0.0004)
	1.288** (0.007)	1.555** (0.007)	1.287** (0.007)	1.561** (0.007)
Male	0.077** (0.002)		0.079** (0.002)	
		0.032** (0.001)		0.033** (0.001)
Visceral fat	-0.129** (0.012)	-0.131** (0.012)		
			-0.191** (0.031)	-0.191** (0.031)
Abdominal subcutaneous fat			-0.041** (0.007)	-0.044** (0.007)
	-0.615** (0.005)	-0.744** (0.005)	-0.602** (0.006)	-0.734** (0.005)
Sarcopenia	42,742	42,742	41,376	41,376
Adjusted R²	0.577	0.569	0.580	0.572
AIC	84558.3	85352.9	81531.5	82333.8

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S13 - Regression coefficients between thigh IMAT volume and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Thigh IMAT Volume			
	(1)	(2)	(3)	(4)
Age	0.011** (0.0005)	0.022** (0.0005)	0.011** (0.0005)	0.021** (0.0005)
	-0.324** (0.008)	0.811** (0.008)	-0.300** (0.008)	0.797** (0.008)
Male	0.305** (0.002)		0.296** (0.002)	
		0.155** (0.001)		0.151** (0.001)
Sarcopenia	0.029* (0.014)	0.007 (0.014)		
			0.386** (0.035)	0.287** (0.035)
Frailty [Frail]			0.094** (0.008)	0.047** (0.008)
Constant	0.154** (0.006)	-0.393** (0.005)	0.095** (0.006)	-0.415** (0.006)
Observations	42,783	42,783	41,415	41,415
Adjusted R²	0.427	0.421	0.429	0.421
AIC	97495.6	97908.3	92363.8	92983

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S14 - Regression coefficients between thigh IMAT volume index and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Thigh IMAT Volume Index			
	(1)	(2)	(3)	(4)
Age	0.016** (0.0005)	0.027** (0.0005)	0.015** (0.0005)	0.026** (0.0005)
	-0.676**	0.467**	-0.642**	0.452**
Male				

	(0.009)	(0.008)	(0.009)	(0.008)
Visceral fat	0.303**		0.291**	
	(0.002)		(0.002)	
Abdominal subcutaneous fat		0.159**		0.154**
		(0.001)		(0.001)
Sarcopenia	0.106**	0.081**		
	(0.014)	(0.014)		
Frailty [Frail]			0.507**	0.392**
			(0.036)	(0.035)
Frailty [Pre Frail]			0.145**	0.093**
			(0.008)	(0.008)
Constant	0.318**	-0.233**	0.237**	-0.270**
	(0.006)	(0.005)	(0.007)	(0.006)
Observations	42,740	42,740	41,374	41,374
Adjusted R²	0.395	0.413	0.401	0.413
AIC	99801.4	98541.4	93926.6	93046.3

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S15 - Regression coefficients between mid-thigh IMAT volume and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Mid-thigh IMAT Volume			
	(1)	(2)	(3)	(4)
Age	0.007**	0.017**	0.006**	0.016**
	(0.001)	(0.001)	(0.001)	(0.001)
Male	-0.520**	0.538**	-0.492**	0.522**
	(0.009)	(0.008)	(0.009)	(0.008)
Visceral fat	0.285**		0.275**	
	(0.002)		(0.002)	
Abdominal subcutaneous fat		0.144**		0.139**
		(0.001)		(0.001)

	0.073**	0.052*		
Sarcopenia	(0.015)	(0.015)		
			0.399**	0.314**
Frailty [Frail]			(0.037)	(0.038)
			0.107**	0.066**
Frailty [Pre Frail]			(0.008)	(0.008)
	0.245**	-0.264**	0.183**	-0.290**
Constant	(0.006)	(0.006)	(0.007)	(0.006)
Observations	42,779	42,779	41,412	41,412
Adjusted R²	0.336	0.326	0.340	0.325
AIC	103905.4	104560.2	97838.6	98783.1

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S16 - Regression coefficients between mid-thigh IMAT/muscle ratio and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Mid-thigh IMAT/Muscle			
	(1)	(2)	(3)	(4)
Age	0.015** (0.001)	0.024** (0.001)	0.015** (0.001)	0.023** (0.001)
Male	-0.851** (-0.01)	0.053** (0.009)	-0.818** (0.009)	0.041** (0.008)
Visceral fat	0.239** (0.002)		0.229** (0.002)	
Abdominal subcutaneous fat		0.126** (0.001)		0.121** (0.001)
Sarcopenia	0.145** (0.016)	0.125** (0.016)		
Frailty [Frail]			0.507** (0.038)	0.415** (0.038)
Frailty [Pre Frail]			0.135**	0.093**

			(0.008)	(0.008)
Constant	0.400**	-0.036**	0.325**	-0.072**
	(0.006)	(0.006)	(0.007)	(0.006)
Observations	42,779	42,779	41,412	41,412
Adjusted R²	0.271	0.286	0.290	0.300
AIC	107990	107094.8	98812	98250.4

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S17 - Regression coefficients between paraspinal muscle PDFF and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Paraspinal Muscle PDFF			
	(1)	(2)	(3)	(4)
Age	0.026**	0.034**	0.025**	0.033**
	(0.001)	(0.001)	(0.001)	(0.001)
Male	-0.836**	0.070**	-0.807**	0.060**
	(0.009)	(0.008)	(0.009)	(0.008)
Visceral fat	0.236**		0.227**	
	(0.002)		(0.002)	
Abdominal subcutaneous fat		0.129**		0.126**
		(0.001)		(0.001)
Sarcopenia	0.105**	0.082**		
	(0.015)	(0.015)		
Frailty [Frail]			0.469**	0.362**
			(0.039)	(0.038)
Frailty [Pre Frail]			0.142**	0.094**
			(0.008)	(0.008)
Constant	0.394**	-0.043**	0.321**	-0.077**
	(0.006)	(0.006)	(0.007)	(0.006)
Observations	42,546	42,546	41,186	41,186
Adjusted R²	0.297	0.328	0.299	0.327

AIC	105354.6	103414	100151.8	98452.6
------------	----------	--------	----------	---------

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S18 - Regression coefficients between paraspinal muscle iron concentration and disease (Sarcopenia, Sarcopenic Obesity and Frailty). Standard errors are in parentheses.

	Paraspinal Muscle Iron Concentration			
	(1)	(2)	(3)	(4)
Age	-0.009** (0.001)	-0.010** (0.001)	-0.009** (0.001)	-0.010** (0.001)
	-0.015 (0.011)	-0.061** (0.01)	-0.018 (0.011)	-0.057** (0.01)
Male	-0.006* (0.002)		-0.003 (0.003)	
		-0.012** (0.001)		-0.011** (0.001)
Sarcopenia	0.061* (0.018)	0.066* (0.018)		
			-0.087 (0.047)	-0.051 (0.047)
Frailty [Frail]			0.006 (0.01)	0.018 (0.01)
Constant	-0.001 (0.007)	0.021* (0.007)	0.004 (0.009)	0.017* (0.008)
Observations	42,546	42,546	41,186	41,186
Adjusted R²	0.006	0.008	0.006	0.007
AIC	119932.9	119842.7	115937.3	115861.7

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex and visceral fat,
- (2) Model for sarcopenia adjusted for age, sex and abdominal subcutaneous fat,
- (3) Model for frailty adjusted for age, sex and visceral fat,
- (4) Model for frailty adjusted for age, sex and abdominal subcutaneous fat.

Table S19 - Regression coefficients between sarcopenia and the anthropometric covariate and muscle and fat IDPs using a logistic regression. Values are presented as odds ratios (OR) and 95% confidence intervals (CI) in parentheses.

	Model - Sarcopenia								
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Age	1.466** (1.404, 1.532)	1.496** (1.433, 1.562)	1.488** (1.424, 1.556)	1.542** (1.477, 1.611)	1.767** (1.696, 1.842)	1.766** (1.695, 1.841)	1.748** (1.678, 1.822)	1.743** (1.672, 1.817)	1.779** (1.707, 1.854)
Male	3.437** (2.955, 3.998)	2.953** (2.561, 3.404)	1.994** (1.752, 2.270)	2.129** (1.851, 2.448)	0.793** (0.719, 0.875)	0.811** (0.735, 0.895)	0.863* (0.780, 0.955)	0.846* (0.763, 0.937)	0.790** (0.717, 0.871)
Asian	1.301 (0.970, 1.720)	1.432* (1.069, 1.890)	1.614* (1.206, 2.129)	1.460* (1.091, 1.926)	2.190** (1.644, 2.875)	2.190** (1.643, 2.874)	2.147** (1.611, 2.819)	2.153** (1.616, 2.827)	2.168** (1.627, 2.846)
Black	0.613 (0.275, 1.176)	0.667 (0.298, 1.283)	0.691 (0.310, 1.325)	0.449* (0.202, 0.858)	0.440* (0.198, 0.838)	0.433* (0.195, 0.824)	0.433* (0.195, 0.826)	0.448* (0.202, 0.852)	0.452* (0.204, 0.860)
Chinese	1.087 (0.561, 1.922)	1.165 (0.601, 2.059)	1.374 (0.710, 2.426)	1.171 (0.604, 2.071)	1.677 (0.867, 2.958)	1.687 (0.872, 2.977)	1.704 (0.881, 3.006)	1.677 (0.866, 2.960)	1.638 (0.847, 2.889)
Others	0.868 (0.573, 1.265)	0.917 (0.606, 1.335)	1.006 (0.667, 1.463)	0.857 (0.567, 1.245)	1.007 (0.669, 1.458)	1.010 (0.671, 1.462)	1.015 (0.674, 1.470)	1.004 (0.667, 1.454)	0.996 (0.661, 1.441)
Waist-to-Hip ratio	1.304** (1.241, 1.370)	1.299** (1.237, 1.365)	1.319** (1.256, 1.386)	1.216** (1.158, 1.277)	1.153** (1.095, 1.214)	1.134** (1.078, 1.193)	1.120** (1.065, 1.177)	1.129** (1.072, 1.188)	1.171** (1.116, 1.229)
Alcohol frequency [Daily]	0.681** (0.602, 0.771)	0.696** (0.614, 0.787)	0.686** (0.606, 0.776)	0.672** (0.593, 0.760)	0.667** (0.590, 0.754)	0.668** (0.590, 0.755)	0.667** (0.590, 0.754)	0.671** (0.593, 0.759)	0.671** (0.593, 0.758)
	0.767**	0.777**	0.770**	0.763**	0.741**	0.743**	0.744**	0.745**	0.742**

Alcohol frequency [1-4 times per week]	(0.696, 0.846)	(0.705, 0.857)	(0.699, 0.849)	(0.693, 0.841)	(0.673, 0.817)	(0.674, 0.818)	(0.676, 0.820)	(0.676, 0.821)	(0.674, 0.818)
Alcohol frequency [1-3 times per month]	0.776*	0.780*	0.780*	0.780*	0.765*	0.765*	0.765*	0.769*	0.765*
	(0.675, 0.891)	(0.679, 0.895)	(0.679, 0.895)	(0.679, 0.895)	(0.667, 0.877)	(0.667, 0.877)	(0.666, 0.877)	(0.669, 0.881)	(0.666, 0.877)
	0.939	0.923	0.927	0.912*	0.916*	0.912*	0.907*	0.916*	0.922*
Smoking status [Previous]	(0.866, 1.017)	(0.851, 1.000)	(0.856, 1.004)	(0.841, 0.987)	(0.845, 0.991)	(0.842, 0.987)	(0.838, 0.982)	(0.846, 0.992)	(0.851, 0.998)
	0.977	0.925	0.931	0.952	1.004	1.004	0.998	1.005	1.004
Smoking status [Current]	(0.780, 1.212)	(0.738, 1.148)	(0.743, 1.154)	(0.760, 1.179)	(0.802, 1.241)	(0.803, 1.242)	(0.798, 1.234)	(0.804, 1.244)	(0.803, 1.242)
	1.045*	1.045*	1.052*	1.048*	1.061*	1.059*	1.058*	1.060*	1.062*
Townsend deprivation index	(1.007, 1.084)	(1.007, 1.085)	(1.013, 1.092)	(1.009, 1.087)	(1.022, 1.100)	(1.021, 1.099)	(1.019, 1.097)	(1.021, 1.099)	(1.023, 1.101)
	0.886**	0.887**	0.885**	0.869**	0.862**	0.865**	0.869**	0.863**	0.858**
MET	(0.847, 0.926)	(0.848, 0.926)	(0.846, 0.925)	(0.831, 0.908)	(0.824, 0.901)	(0.826, 0.903)	(0.830, 0.908)	(0.825, 0.902)	(0.820, 0.897)
	0.361**								
Total Muscle Volume	(0.332, 0.392)								
		0.387**							
Thigh Muscle Volume		(0.358, 0.418)							
			0.477**						
Mid-thigh Muscle Volume			(0.444, 0.512)						
				0.501**					
Iliopsoas Muscle Volume				(0.466, 0.539)					
					1.034				
Thigh IMAT Volume					(0.994, 1.074)				

						1.071**			
Mid-thigh IMAT Volume						(1.035, 1.107)			
							1.116**		
Mid-thigh IMAT/Muscle							(1.082, 1.152)		
								1.075**	
Paraspinal Muscle PDFF								(1.039, 1.113)	
Paraspinal Muscle Iron Concentration									1.054*
									(1.018, 1.091)
Constant	0.045** (0.040, 0.051)	0.049** (0.043, 0.054)	0.061** (0.054, 0.068)	0.061** (0.054, 0.068)	0.105** (0.095, 0.115)	0.103** (0.094, 0.113)	0.100** (0.091, 0.110)	0.101** (0.092, 0.111)	0.104** (0.095, 0.114)
Observations	42,010	42,010	42,010	42,010	42,010	42,010	42,010	42,010	42,010
AIC	20833.6	20839	21014.4	21085.9	21448.3	21436.2	21404.5	21434.6	21442.6

* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

- (1) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and total muscle volume,
- (2) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and thigh muscle volume,
- (3) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh muscle volume,
- (4) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and iliopsoas muscle volume,
- (5) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and thigh IMAT volume,
- (6) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh IMAT volume,

- (7) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh IMAT/Muscle,
- (8) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and paraspinal muscle PDFF,
- (9) Model for sarcopenia adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and paraspinal muscle iron concentration.

Table S20 - Regression coefficients between frailty and the anthropometric covariate and muscle and fat IDPs using an ordinal logistic regression. Values are presented as odds ratios (OR) and 95% confidence intervals (CI).

	Model - Frailty								
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Age	1.120** (1.097, 1.143)	1.110** (1.087, 1.132)	1.131** (1.108, 1.154)	1.112** (1.089, 1.134)	1.147** (1.126, 1.169)	1.155** (1.134, 1.176)	1.131** (1.109, 1.152)	1.115** (1.093, 1.136)	1.167** (1.146, 1.188)
	0.950 (0.866, 1.033)	1.010 (0.930, 1.089)	0.823** (0.750, 0.896)	0.986 (0.907, 1.065)	0.731** (0.676, 0.786)	0.772** (0.716, 0.827)	0.870** (0.812, 0.928)	0.870** (0.812, 0.928)	0.697** (0.643, 0.752)
Male	1.360* (1.163, 1.557)	1.346* (1.149, 1.542)	1.439* (1.242, 1.635)	1.321* (1.124, 1.518)	1.531** (1.335, 1.727)	1.518** (1.322, 1.714)	1.451* (1.254, 1.647)	1.460* (1.263, 1.657)	1.521** (1.326, 1.717)
	1.376* (1.118, 1.634)	1.467* (1.208, 1.726)	1.378* (1.119, 1.636)	1.285 (1.028, 1.542)	1.176 (0.918, 1.434)	1.166 (0.908, 1.425)	1.211 (0.952, 1.469)	1.280 (1.022, 1.537)	1.245 (0.989, 1.501)
Asian	0.998 (0.628, 1.368)	0.984 (0.615, 1.354)	1.057 (0.688, 1.426)	0.974 (0.603, 1.344)	1.248 (0.879, 1.618)	1.189 (0.820, 1.559)	1.186 (0.816, 1.555)	1.178 (0.807, 1.548)	1.101 (0.732, 1.470)
	1.157 (0.952, 1.362)	1.166 (0.961, 1.371)	1.184 (0.979, 1.389)	1.125 (0.920, 1.330)	1.195 (0.989, 1.400)	1.185 (0.980, 1.391)	1.194 (0.988, 1.399)	1.191 (0.986, 1.396)	1.175 (0.970, 1.380)
Others									

	1.387**	1.395**	1.384**	1.376**	1.218**	1.220**	1.220**	1.209**	1.355**
Waist-to-Hip ratio	(1.359, 1.415)	(1.367, 1.423)	(1.356, 1.412)	(1.348, 1.403)	(1.189, 1.247)	(1.191, 1.250)	(1.191, 1.249)	(1.179, 1.238)	(1.328, 1.383)
	0.541**	0.544**	0.542**	0.539**	0.524**	0.531**	0.530**	0.540**	0.540**
Alcohol frequency [Daily]	(0.470, 0.612)	(0.473, 0.615)	(0.471, 0.612)	(0.468, 0.610)	(0.453, 0.595)	(0.460, 0.602)	(0.459, 0.602)	(0.469, 0.611)	(0.469, 0.610)
Alcohol frequency [1-4 times per week]	0.622**	0.625**	0.622**	0.623**	0.615**	0.618**	0.620**	0.623**	0.618**
	(0.565, 0.678)	(0.568, 0.682)	(0.565, 0.679)	(0.566, 0.680)	(0.558, 0.672)	(0.561, 0.675)	(0.563, 0.677)	(0.566, 0.680)	(0.561, 0.675)
Alcohol frequency [1-3 times per month]	0.793**	0.795**	0.793**	0.795**	0.787**	0.789**	0.792**	0.794**	0.790**
	(0.716, 0.869)	(0.719, 0.872)	(0.717, 0.870)	(0.718, 0.871)	(0.711, 0.864)	(0.712, 0.866)	(0.715, 0.869)	(0.718, 0.871)	(0.713, 0.866)
	1.121**	1.119**	1.117**	1.114**	1.080*	1.088*	1.086*	1.100**	1.114**
Smoking status [Previous]	(1.077, 1.164)	(1.075, 1.162)	(1.073, 1.160)	(1.070, 1.158)	(1.036, 1.124)	(1.044, 1.132)	(1.042, 1.130)	(1.056, 1.144)	(1.070, 1.157)
	1.228*	1.212*	1.218*	1.217*	1.194*	1.211*	1.201*	1.228*	1.231*
Smoking status [Current]	(1.114, 1.342)	(1.097, 1.326)	(1.104, 1.333)	(1.103, 1.331)	(1.079, 1.309)	(1.096, 1.325)	(1.086, 1.316)	(1.113, 1.342)	(1.117, 1.345)
	1.044**	1.043*	1.045**	1.042*	1.042*	1.041*	1.039*	1.041*	1.047**
Townsend deprivation index	(1.023, 1.064)	(1.022, 1.064)	(1.024, 1.066)	(1.022, 1.063)	(1.021, 1.063)	(1.020, 1.061)	(1.018, 1.060)	(1.020, 1.062)	(1.026, 1.067)
	0.932**	0.933**	0.931**	0.929**	0.944**	0.940**	0.944**	0.936**	0.926**
MET	(0.911, 0.953)	(0.912, 0.954)	(0.910, 0.952)	(0.908, 0.950)	(0.923, 0.965)	(0.919, 0.961)	(0.923, 0.965)	(0.915, 0.957)	(0.905, 0.947)
	0.820**								
Total Muscle Volume	(0.779, 0.862)								
		0.783**							
Thigh Muscle Volume		(0.744, 0.822)							
Mid-thigh Muscle Volume			0.886**						

			(0.850, 0.922)						
Iliopsoas Muscle Volume				0.801**					
				(0.764, 0.838)					
Thigh IMAT Volume					1.268**				
					(1.244, 1.291)				
Mid-thigh IMAT Volume						1.273**			
						(1.249, 1.297)			
Mid-thigh IMAT/Muscle							1.328**		
							(1.304, 1.353)		
Paraspinal Muscle PDFF								1.290**	
								(1.266, 1.313)	
Paraspinal Muscle Iron Concentration								0.990	
								(0.969, 1.010)	
Not Frail Pre Frail	1.094 (1.093, 1.095)	1.131 (1.13, 1.132)	1.021 (1.021, 1.021)	1.111 (1.11, 1.112)	0.94 (0.941, 0.94)	0.971 (0.971, 0.971)	1.029 (1.029, 1.029)	1.041 (1.041, 1.042)	0.937 (0.938, 0.937)
	67.862 (65.144, 70.694)	70.375 (67.533, 73.337)	63.203 (60.714, 65.795)	69.1 (66.321, 71.996)	59.566 (57.253, 61.973)	61.889 (59.463, 64.413)	66.4 (63.754, 69.156)	66.773 (64.109, 69.548)	57.925 (55.691, 60.249)
Observations	40,897	40,897	40,897	40,897	40,897	40,897	40,897	40,897	
AIC	57232	57166.7	57277.9	57183.3	56915.6	56907	56794.8	56857	57320


* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

(1) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and total muscle volume,

- (2) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and thigh muscle volume,
- (3) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh muscle volume,
- (4) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and iliopsoas muscle volume,
- (5) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and thigh IMAT volume,
- (6) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh IMAT volume,
- (7) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and mid-thigh IMAT/Muscle,
- (8) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and paraspinal muscle PDFF,
- (9) Model for frailty adjusted for age, sex, ethnicity, waist-to-hip ratio, alcohol intake frequency, smoking status, Townsend deprivation index, MET, and paraspinal muscle iron concentration.

Supplementary Figures

Figure S1 Scatterplot showing the correlation coefficients between muscle volume IDPs and dominant hand grip strength, separated by presence of sarcopenia (red for TRUE, blue for FALSE) in both genders.



* indicate statistically significant for $p < 0.05$, ** indicate statistically significant after Bonferroni correction ($p = 5.7 \times 10^{-5}$).

References

1. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci.* 2001;56:M146–56.