Supplemental Information

Supplemental figures



Supplemental Figure 1: Cumulative Distribution Function plotted against consensus index. The Consensus index represents how often two samples are placed together in the same cluster. A consensus value of one indicates that the two samples are always grouped together while a value of 0 indicates that two samples are never grouped together. Percent of Ambiguous Clusters (PAC) is a measure of how much of the CDF falls between a consensus index between 0.1 and 0.9. The plot above shows that as we increase the number of clusters, it's clear that the PAC will always decrease. Hence choosing the minimal PAC will lead to a cluster size equal to the number of samples.



Intermediate vs Hypoinflammatory



Intermediate vs Hyperinflammatory

Supplemental Tables

		Standard				
Column1	Mean	Deviation	First	Last	Min	Max
CREACTIV	CREACTIVE_	CREACTIVE	CREACTIVE_	CREACTIVE_	CREACTIVE_	
E_PROTEI	PROTEIN_M	_PROTEIN_	PROTEIN_Fir	PROTEIN_La	PROTEIN_Mi	
Ν	ean	sd	st	st	n	
	DDIMER_Me		DDIMER_Fir	DDIMER_Las	DDIMER_Mi	
DDIMER	an		st	t	n	
	FERRITIN_M					
FERRITIN	ean					
LACTATE_	LACTATE_DE		LACTATE_D			
DEHYDRO	HYDROGENA		EHYDROGE			
GENASE	SE_Mean		NASE_First			
LYMPHOC	LYMPHOCYT	LYMPHOCY	LYMPHOCYT	LYMPHOCYT	LYMPHOCYT	LYMPHOCYT
YTES_perc	ES_percent_	TES_percen	ES_percent_	ES_percent_	ES_percent_	ES_percent_
ent	Mean	t_sd	First	Last	Min	Max
PLATELET_	PLATELET_C	PLATELET_C	PLATELET_C	PLATELET_C	PLATELET_C	PLATELET_C
COUNT	OUNT_Mean	OUNT_sd	OUNT_First	OUNT_Last	OUNT_Min	OUNT_Max
NEUTROP		NEUTROPHI		NEUTROPHI	NEUTROPHI	NEUTROPHIL
HIL_NUM		L_NUMBER		L_NUMBER_	L_NUMBER_	_NUMBER_
BER_ANC		_ANC_sd		ANC_Last	ANC_Min	ANC_Max
	EOSINOPHIL	EOSINOPHI	EOSINOPHIL	EOSINOPHIL	EOSINOPHIL	
EOSINOPH	_percent_M	L_percent_	_percent_Fi	_percent_La	_percent_M	
IL_percent	ean	sd	rst	st	in	
BUNCR	BUNCREA	BUNCRE	BUNCRE	BUNCREA	BUNCRE	
EATININE_	TININE_RATI	ATININE_R	ATININE_RA	TININE_RATI	ATININE_RA	
RATIO	O_Mean	ATIO_sd	TIO_First	O_Last	TIO_Min	
INRPROT			INRPROTI		INRPROTI	
IME			ME_First		ME_Min	
			WBC_COUN	WBC_COUN	WBC_COUN	WBC_COUN
WBC_COU					T_URINE_Mi	T_URINE_M
NT_URINE			st –	st – –	n – –	ax
WBC_ord_	WBC_ord_n	WBC_ord_n	WBC_ord_n	WBC_ord_n	WBC_ord_n	
num_valu	um_value_M	um_value_s	um_value_F	um_value_L	um_value_	
e	ean	d	irst –	ast	Min	
	INR_ord_nu			INR_ord_nu		
INR_ord_n	m_value_Me			m_value_La		
um value	an			st		

Conditio n	ICD-10 Code	Coun t	Total Count	Unique Total Count	
DVT	182	444	444	336	
	126	287			
	T81.7	3			
DE	T82.8 49		054	341	
PE	T79.0	T79.0 2 354			
	T79.1	1			
	T80.0	0	-		
	121	442			
	122	1		500	
MI	123 0		548	536	
	124	97			

Supplemental Table 2:ICD 10 codes used to explore outcomes

Stroke:	18 45 103 33 3 15		
---------	----------------------------------	--	--

Supplemental Table 3. Race Results

cluster	first_race	n
1	Am Indian	6
1	Asian	160
1	Black	1128
	NotDisclos	
1	е	1
1	Other	615
	Other	
1	Pacifi	1
	Рас	
1	Islander	1
1	Unknown	13
1	White	1148
1	NA	3
2	Am Indian	4
2	Asian	128
2	Black	598
2	Hispanic	1
	NotDisclos	
2	е	2
2	Other	335
	Other	
2	Pacifi	1
	Рас	
2	Islander	2
2	Unknown	15
2	White	1004
3	Am Indian	3
3	Asian	65
3	Black	859
3	Hispanic	1
	NotDisclos	
3	е	3
3	Other	370
3	Рас	1

Islander

 3
 Unknown
 7

 3
 White
 593

 3
 NA
 8

Supplemental Equation

Standardized Mean Difference

$$SMD = \frac{\overline{X_1} - \overline{X_2}}{\sqrt{(S_1^2 + S_2^2)/2}}$$

Where $\overline{X_1}$ and $\overline{X_2}$ represent the mean of a variable of interest in group 1 and two respectively while S_1 and S_2 represent the variance.