

Supplementary information for: Adolescent pregnancy is associated with child undernutrition: systematic review and meta-analysis

Supplementary Appendix 1. Search strategy and results

- 1 (Adolescen* or teen* or youth* or young person* or young people* or "10-19 year*").mp.
[mp=title, book title, abstract, original title, name of substance word, subject heading word, floating
sub-heading word, keyword heading word, organism supplementary concept word, protocol
supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms]
(2306108)
- 2 adolescent/ or young adult/ or child/ (3518101)
- 3 1 or 2 (3619539)
- 4 (Pregnan* or mother* or matern*).mp. [mp=title, book title, abstract, original title, name of
substance word, subject heading word, floating sub-heading word, keyword heading word, organism
supplementary concept word, protocol supplementary concept word, rare disease supplementary
concept word, unique identifier, synonyms] (1330001)
- 5 Pregnant Women/ (12699)
- 6 pregnancy/ or pregnancy in adolescence/ (957946)
- 7 maternal age/ (20750)
- 8 Adolescent Mothers/ (46)
- 9 4 or 5 or 6 or 7 or 8 (1330001)
- 10 Nutrition* status.mp. [mp=title, book title, abstract, original title, name of substance word,
subject heading word, floating sub-heading word, keyword heading word, organism supplementary
concept word, protocol supplementary concept word, rare disease supplementary concept word,
unique identifier, synonyms] (74538)
- 11 malnutrition.mp. or *nutrition/ or *weight/ or malnourished.mp. (61251)
- 12 Nutritional Status/ (51802)
- 13 malnutrition/ or deficiency diseases/ (25335)
- 14 severe acute malnutrition/ or kwashiorkor/ (3030)
- 15 Starvation/ (10139)
- 16 overnutrition/ or overweight/ or obesity/ or obesity, maternal/ (220192)
- 17 Wasting Syndrome/ (1432)
- 18 protein deficiency/ or protein-energy malnutrition/ (12271)
- 19 Thinness/ (7299)
- 20 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 (356466)
- 21 9 or 20 (1653395)
- 22 (Infant* or child* or baby or babies or toddler* or offspring or next generation or pre-school
child or "Under 5 years").mp. [mp=title, book title, abstract, original title, name of substance word,
subject heading word, floating sub-heading word, keyword heading word, organism supplementary
concept word, protocol supplementary concept word, rare disease supplementary concept word,
unique identifier, synonyms] (3404648)
- 23 child/ or child, preschool/ or infant/ (2354343)
- 24 22 or 23 (3404648)
- 25 (Wasting or wasted or kwashiorkor or marasmus or "weight for height" or WHZ or Growth
falter* or underweight or "weight for age" or WAZ or nutrition* depriv* or starvation or famine* or
acute malnutrition or moderate malnutrition).mp. [mp=title, book title, abstract, original title, name of
substance word, subject heading word, floating sub-heading word, keyword heading word, organism
supplementary concept word, protocol supplementary concept word, rare disease supplementary
concept word, unique identifier, synonyms] (107973)
- 26 exp malnutrition/ (132030)
- 27 (afghanistan or albania or algeria or american samoa or angola or "antigua and barbuda" or
antigua or barbuda or argentina or armenia or armenian or aruba or azerbaijan or bahrain or
bangladesh or barbados or republic of belarus or belarus or byelarus or belorussia or byelorussian or
belize or british honduras or benin or dahomey or bhutan or bolivia or "bosnia and herzegovina" or
bosnia or herzegovina or botswana or bechuanaland or brazil or brasil or bulgaria or burkina faso or

burkina fasso or upper volta or burundi or urundi or cabo verde or cape verde or cambodia or
 kampuchea or khmer republic or cameroon or cameron or cameroun or central african republic or
 ubangi shari or chad or chile or china or colombia or comoros or comoro islands or iles comores or
 mayotte or democratic republic of the congo or democratic republic congo or congo or zaire or costa
 rica or "cote d'ivoire" or "cote d' ivoire" or cote divoire or cote d ivoire or ivory coast or croatia or
 cuba or cyprus or czech republic or czechoslovakia or djibouti or french somaliland or dominica or
 dominican republic or ecuador or egypt or united arab republic or el salvador or equatorial guinea or
 spanish guinea or eritrea or estonia or eswatini or swaziland or ethiopia or fiji or gabon or gabonese
 republic or gambia or "georgia (republic)" or georgian or ghana or gold coast or gibraltar or greece or
 grenada or guam or guatemala or guinea or guinea bissau or guyana or british guiana or haiti or
 hispaniola or honduras or hungary or india or indonesia or timor or iran or iraq or isle of man or
 jamaica or jordan or kazakhstan or kazakh or kenya or "democratic people's republic of korea" or
 republic of korea or north korea or south korea or korea or kosovo or kyrgyzstan or kirghizia or
 kirgizstan or kyrgyz republic or kirghiz or laos or lao pdr or "lao people's democratic republic" or
 latvia or lebanon or lebanese republic or lesotho or basutoland or liberia or libya or libyan arab
 jamahiriya or lithuania or macau or macao or republic of north macedonia or macedonia or
 madagascar or malagasy republic or malawi or nyasaland or malaysia or malay federation or malaya
 federation or maldives or indian ocean islands or indian ocean or mali or malta or micronesia or
 federated states of micronesia or kiribati or marshall islands or nauru or northern mariana islands or
 palau or tuvalu or mauritania or mauritius or mexico or moldova or moldovian or mongolia or
 montenegro or morocco or ifni or mozambique or portuguese east africa or myanmar or burma or
 namibia or nepal or netherlands antilles or nicaragua or niger or nigeria or oman or muscat or pakistan
 or panama or papua new guinea or new guinea or paraguay or peru or philippines or philipines or
 philippines or phillippines or poland or "polish people's republic" or portugal or portuguese republic or
 puerto rico or romania or russia or russian federation or ussr or soviet union or union of soviet
 socialist republics or rwanda or ruanda or samoa or pacific islands or polynesia or samoan islands or
 navigator island or navigator islands or "sao tome and principe" or saudi arabia or senegal or serbia or
 seychelles or sierra leone or slovakia or slovak republic or slovenia or melanesia or solomon island or
 solomon islands or norfolk island or norfolk islands or somalia or south africa or south sudan or sri
 lanka or ceylon or "saint kitts and nevis" or "st. kitts and nevis" or saint lucia or "st. lucia" or "saint
 vincent and the grenadines" or saint vincent or "st. vincent" or grenadines or sudan or suriname or
 surinam or dutch guiana or netherlands guiana or syria or syrian arab republic or tajikistan or
 tadjikistan or tadjikistan or tadjhik or tanzania or tanganyika or thailand or siam or timor leste or
 east timor or togo or togolese republic or tonga or "trinidad and tobago" or trinidad or tobago or
 tunisia or turkey or turkmenistan or turkmen or uganda or ukraine or uruguay or uzbekistan or uzbek
 or vanuatu or new hebrides or venezuela or vietnam or viet nam or middle east or west bank or gaza
 or palestine or yemen or yugoslavia or zambia or zimbabwe or northern rhodesia or global south or
 africa south of the sahara or sub-saharan africa or subsaharan africa or africa, central or central africa
 or africa, northern or north africa or northern africa or magreb or maghrib or sahara or africa, southern
 or southern africa or africa, eastern or east africa or eastern africa or africa, western or west africa or
 western africa or west indies or indian ocean islands or caribbean or central america or latin america
 or "south and central america" or south america or asia, central or central asia or asia, northern or
 north asia or northern asia or asia, southeastern or southeastern asia or south eastern asia or southeast
 asia or south east asia or asia, western or western asia or europe, eastern or east europe or eastern
 europe or developing country or developing countries or developing nation? or developing
 population? or developing world or less developed countr* or less developed nation? or less
 developed population? or less developed world or lesser developed countr* or lesser developed
 nation? or lesser developed population? or lesser developed world or under developed countr* or
 under developed nation? or under developed population? or under developed world or
 underdeveloped countr* or underdeveloped nation? or underdeveloped population? or underdeveloped
 world or middle income countr* or middle income nation? or middle income population? or low
 income countr* or low income nation? or low income population? or lower income countr* or lower
 income nation? or lower income population? or underserved countr* or underserved nation? or
 underserved population? or underserved world or under served countr* or under served nation? or
 under served population? or under served world or deprived countr* or deprived nation? or deprived

population? or deprived world or poor countr* or poor nation? or poor population? or poor world or poorer countr* or poorer nation? or poorer population? or poorer world or developing econom* or less developed econom* or lesser developed econom* or under developed econom* or underdeveloped econom* or middle income econom* or low income econom* or lower income econom* or low gdp or low gnp or low gross domestic or low gross national or lower gdp or lower gnp or lower gross domestic or lower gross national or lmic or lmics or third world or lami countr* or transitional countr* or emerging economies or emerging nation?).mp. [mp=title, book title, abstract, original title, name of substance word, subject heading word, floating sub-heading word, keyword heading word, organism supplementary concept word, protocol supplementary concept word, rare disease supplementary concept word, unique identifier, synonyms] (2317513)

28 25 or 26 (222101)

29 limit 28 to yr="1990 -Current" (156748)

30 3 and 21 and 24 and 27 and 28 and 29 (6764)

Note: numbers in parenthesis show the number of results per line

Supplementary Appendix 2. Study characteristics

Ref.	Publication Reference	Country	Data source	Sample size	Child age category* (months)	Wasting outcome		Underweight outcome		Youngest maternal age <20 years	Data extractable for meta-analysis	Quality assessment
						Mod ^a	Sev ^b	Mod ^c	Sev ^d			
Cross-sectional studies												
29	(Adhikari et al., 2021)	India	NFHS-4 2015-16	126431	0-36	■	⊕	■	⊕	⊕	■	Good
28	(Aboagye et al., 2021)	32 Sub-Saharan Africa countries	DHS data 2010-2020	48968	6-23	■	⊕	■	⊕	■	■	Good
27	(Ahmed, Ahmed, Roy, Alam, & Hossain, 2012)	Bangladesh	Bangladesh National Nutrition Programme baseline survey 2004	8858	0-24	■	■	■	■	■	⊕	Fair
46	(Akseer et al., 2018)	Afghanistan	ANNS 2013	14000	0-59	■	⊕	■		■	⊕	Good
14	(Ali et al., 2019)	Bangladesh	Nobokoli programme baseline data	6468	0-59	■	⊕	■	⊕	⊕	⊕	Fair
19	(Alkaff, Flynn, Sukmajaya, & Salamah, 2020)	Indonesia	Cross-sectional survey and secondary data from child health records	218	0-59	⊕	⊕	■	⊕	■	■	Good
53	(Amadu et al., 2021)	31 Sub-Saharan countries	DHS 2010-2019	129646	0-59	■	⊕	■	⊕	■	■	Good
22	(Amare, Ahmed, & Mehari, 2019)	Ethiopia	DHS 2016	9419	0-59	■	⊕	⊕	⊕	■	■	Good
56	(Beiersmann et al., 2013)	Burkina Faso	Cross-sectional data	460	6-51	■	⊕	■	⊕		⊕	Good
86	(Bekele, Rawstorne, & Rahman, 2021)	Ethiopia	DHS 2000 and 2016	21514	0-59	■	⊕	■	⊕	⊕	⊕	Good
9	(Birhan & Belay, 2021)	Ethiopia	DHS 2016	8935	0-59	⊕	⊕	■	■	■	■	Fair
70	(Biswas, Townsend, Magalhaes, Hasan, & Mamun, 2021)	South and South-East Asian countries	DHS 2007-2016	798961	0-59	■	⊕	■	⊕	■	⊕	Good
36	(Boah, Azupogo, Amporfro, & Abada, 2019)	Ghana	DHS 2014	2636	0-59	■	⊕	■	⊕	■	⊕	Good
78	(S. Das et al., 2019)	Bangladesh	DHS 2014	5951	0-59	■	⊕	■	⊕	■	⊕	Good
20	(Fagbamigbe, Kandala, & Uthman, 2020)	51 LMIC	DHS data 2010-2018	532680	0-59	⊕	■	⊕	⊕		■	Good
47	(Faisal Ahmed et al., 2021)	Bangladesh	DHS 2014	6341	0-59	■	⊕	■	⊕	■	■	Good
16	(Finlay, Ozaltin, & Canning, 2011)	55 LMIC	118 DHS surveys 1990-2008	176583	0-59	■	⊕	■	⊕	■	■	Good

8	(Fuada, Latifah, Yunitawat, & Ashar, 2020)	Indonesia	Basic health research data 2013	978	0-59	■	⊕	■	⊕	■	⊕	Fair
81	(Gbadamosi, Goon, & Tugli, 2017)	South Africa	Cross-sectional data from the MAL-ED study	186	1-12	■	⊕	■	⊕	■	⊕	Good
60	(Geda et al., 2021)	Ethiopia	DHS 2016	9218	6-59	■	⊕	■	⊕	⊕	■	Good
88	(Gewa & Yandell, 2012)	Kenya	DHS survey 2003	1851	0-60	■	⊕	■	⊕	⊕	⊕	Good
84	(Ghimire, Aryal, Gupta, & Sapkota, 2020)	Nepal	Cross-sectional data from children admitted to the OTP	398	6-59	⊕	■	⊕	⊕	■	■	Good
62	(Haque et al., 2022)	South and South-East Asian Countries	DHS 2014-2018	213730	0-59	■	⊕	■	⊕	■	■	Good
64	(Hien & Hoa, 2009)	Vietnam	Cross-sectional data	383	6-36	■	⊕	■	⊕	⊕	⊕	Good
65	(Hien & Kam, 2008)	Vietnam	Cross-sectional data	650	0-59	■	⊕	■	⊕	⊕	⊕	Good
13	(Hiruy et al., 2021)	Ethiopia	DHS 2000-2016	8003	6-23	■	⊕	■	⊕	■	⊕	Good
66	(Horta et al., 2013)	Brazil	National Survey of Indigenous People's Health and Nutrition 2008-2009	6075	0-59	■	⊕	■	⊕	■	■	Good
31	(F. B. Hossain et al., 2020)	Bangladesh, India, Pakistan, Maldives and Nepal	DHS 2009 - 2016	146996	24-59	⊕	⊕	■	⊕	■	⊕	Good
83	(M. B. Hossain & Khan, 2018)	Bangladesh	DHS 2014	7173	0-59	■	⊕	■	⊕	■	■	Good
55	(Huynh, Huynh, Nguyen, Do, & Khanh Tran, 2019)	Vietnam	Cross-sectional data	225	6-59	⊕	⊕	■	⊕	⊕	■	Good
57	(Ickes, Hurst, & Flax, 2015)	Uganda	DHS 2006 and 2011	1009, 888	0-23	■	⊕	■	⊕	■	⊕	Good
76	(Islam & Biswas, 2020)	Bangladesh	DHS 2014	6965	0-59	■	⊕	■	⊕	■	■	Good
10	(Issah, Yeboah, Kpordoxah, Boah, & Mahama, 2022)	Nigeria	DHS 2018	2425	0-59	■	⊕	⊕	⊕	⊕	⊕	Good
82	(Janevic, Petrovic, Bjelic, & Kubera, 2010)	Serbia	MICS 2005	1192	0-59	■	⊕	■	⊕	■	■	Good
1	(Karim et al., 2021)	Bangladesh	Study data	540	6-59	■	⊕		⊕	■	■	Good
72	(Kasaye, Bobo, Yilma, & Woldie, 2019)	Ethiopia	DHS 2016	9494	0-59	■	⊕	■	⊕	⊕	■	Good
26	(S. Khan, Zaheer, & Safdar, 2019)	Pakistan	DHS 2012-2013	3071	0-59	■	⊕	■	⊕	■	■	Good
49	(M. S. Khan, Halder, Rashid, Afroja, & Islam, 2020)	Bangladesh	DHS 2014	8092	0-59	⊕	⊕	■	⊕	■	■	Good
7	(Kim et al., 2019)	India	INFHS-4 2015-16	140,444	0-59	■	⊕	■	⊕	■	■	Good

18	(R. Kumar & Paswan, 2021)	India	NFHS-3 2005-2006, and NFHS-4 2015-2016	11858 (NFHS-3), 92630 (NFHS-4)	0-59	⊕	⊕	■	⊕	■	⊕	Good
50	(A. Kumar & Ram, 2013)	India	NFHS 2005-2006	41306	0-59	⊕	⊕	■	⊕	⊕	■	Good
77	(R. Kumar, Abbas, Mahmood, & Somrongthong, 2019)	Pakistan	MICS 2014	24042	0-59	⊕	⊕	■	■	■	■	Good
21	(Linnemayr, Alderman, & Ka, 2008)	Senegal	Baseline data: nutrition intervention programme	4296	0-35	⊕	⊕	■	⊕	⊕	⊕	Good
40	(Mashal et al., 2008)	Afghanistan	Cross-sectional survey	2472	0-59	■	⊕	■	⊕	■	⊕	Good
32	(Masibo, Humwa, & Macharia, 2020)	Kenya	DHS 2014	7830	0-59	■	⊕	■	⊕	■	⊕	Good
37	(Mena-Melendez, 2020)	Latin America: Bolivia, Colombia, Guatemala and Peru	DHS 1986-2015	15827	0-59	■	⊕	⊕	⊕	■	⊕	Good
80	(Mokwena & Kachabe, 2022)	South Africa	Cross-sectional data	94	1-36	■	■	⊕	⊕	■	⊕	Fair
44	(Mutunga, Frison, Rava, & Bahwere, 2020)	Cambodia, Lao PDR, Myanmar, Thailand, Timor-Leste, Vietnam	DHS 2014, Lao PDR MICS 2017, Myanmar DHS 2015, Thailand MICS 2015/16, Timor-Leste NFNS 2013, and Vietnam MICS 2011	42308	0-59	■	⊕	⊕	⊕	■	■	Good
68	(Nakamori et al., 2010)	Vietnam	Cross-sectional data	188	6-18	⊕	⊕	■	⊕	⊕	⊕	Fair
2	(Nguyen et al., 2021)	Bangladesh	DHS 1996-2017	12006	0-59	■	⊕	⊕	⊕	■	⊕	Good
63	(Nguyen et al., 2017)	Bangladesh	Baseline data: scaled up MNCH programme	2000	0-5	■	⊕	■	⊕	■	⊕	Good
85	(Nguyen, Scott, Neupane, Tran, & Menon, 2019)	India	NFHS-4 2015-16	60096	0-59	■	⊕	■	⊕	■	■	Good
6	(Ntenda & Chuang, 2018)	Malawi	DHS 2004 and 2010	6384	0-59	■	⊕	■	⊕	⊕	■	Good
15	(Ntenda, 2019)	Malawi	DHS 2015-16	6033	0-59	■	⊕	■	⊕	⊕	■	Good
90	(Obayelu & Adeleye, 2021)	Nigeria	DHS 2013	17199	0-59	■	⊕	⊕	⊕	■	■	Good
67	(Olodu, Adeyemi, Olowookere, & Esimai, 2019)	Nigeria	Cross-sectional survey using	300	6-59	■	⊕	■	⊕	■	⊕	Good

primary health care centres records												
42	(Pramod Singh, Nair, Grubestic, & Connell, 2009)	Nepal	Cross-sectional survey	443	6-36	⊕	⊕	■	⊕	■	■	Good
48	(Paul, Chouhan, & Zaveri, 2019)	India	NFHS-4 2015-16	80539	0-59	■	⊕	■	⊕	■	■	Good
51	(Paul & Saha, 2022)	India	NFHS-4 2015-16	38685	0-59	■	⊕	■	⊕	⊕	■	Good
39	(Poda, Hsu, & Chao, 2017)	Burkina Faso	DHS 2010	6337	0-59	■	⊕	■	⊕	■	⊕	Good
17	(Porwal et al., 2021)	India	Comprehensive National Nutrition Survey	35452	0-59	■	⊕	■	⊕	■	■	Good
12	(Qu et al., 2017)	China	Cross sectional survey	12146	6-35	■	⊕	■	⊕		■	Good
91	(A. Rahman & Hossain, 2022)	Bangladesh	DHS 2014	17989	0-59	■	■	■	■	■	■	Good
52	(M. M. Rahman, 2015)	Bangladesh	BDHS 2011	6506	0-59	■	⊕	■	⊕	⊕	■	Good
73	(M. A. Rahman, Halder, Rahman, & Parvez, 2021)	Bangladesh	DHS 2017-18	7738	0-59	■	⊕	■	⊕	⊕	■	Good
79	(M. S. Rahman, Rahman, Maniruzzaman, & Howlader, 2020)	Bangladesh	DHS 2014	7131	0-59	■	■	⊕	⊕	■	■	Good
33	(Raj et al., 2010)	India	NFHS 2005-2006	19392	0-59	■	⊕	■	⊕	■	■	Good
5	(Schott, Aurino, Penny, & Behrman, 2017)	India: Kerala and Goa	NFHS-I 1995	2249	0-59	■	⊕	■	⊕	■	■	Good
38	(Rodgers, Kim, & Subramanian, 2020)	India	NFHS-4 2015-16	139116	6-59	■	⊕	■	⊕	■	⊕	Good
45	(Samuel et al., 2022)	Ethiopia,	Baseline data: effectiveness study	2036	6-11	■	⊕	⊕	⊕	⊕	⊕	Good
23	(Sanjay, Gaurav, Suman, & Ramchandra, 2020)	Rajasthan	Cross-sectional survey	396	0-72	⊕	⊕	■	■	■	■	Fair
30	(Sobkoviak, Yount, & Halim, 2012)	Liberia	DHS 2006-2007	2467	0-48	■	⊕	■	⊕	⊕	⊕	Good
69	(Subramanian, Ackerson, & Smith, 2010)	India	NFHS-3 2005-2006, and NFHS-4 2015-2016	19879	0-59	■	⊕	■	⊕	■	■	Good
87	(Subramanyam, Kawachi, Berkman, & Subramanian, 2010)	India	NFHS 1992, 1998 and 2005	78310	0-35	⊕	⊕	■		■	■	Good
35	(Sunil, 2009)	Yemen	DHS 1997	10414	0-59	■	⊕	■	⊕	■	■	Good
41	(Tariq, Sajjad, Zakar, Zakar, & Fischer, 2018)	Pakistan	PDHS 2012-2013	984	0-23	■	⊕	■	⊕	■	■	Good
59	(Tesfaw & Dessie, 2022)	Ethiopia	EMDHS 2019	5027	0-59	■	⊕	■	⊕	■	■	Good
61	(Tesfaw & Fenta, 2021)	Nigeria	DHS 2018	11314	0-59	■	⊕	■	⊕	■	■	Good

92	(Tibebu, Emiru, Tiruneh, Getu, & Azanaw, 2020)	Ethiopia	Cross-sectional data	634	6 - 59	⊕	⊕	■	⊕	⊕	■	Good
71	(Tiwari, Acharya, Paudel, Sapkota, & Kafle, 2020)	Nepal	DHS 2016	2355	0-59	⊕	⊕	■	⊕	■	■	Good
75	(Li et al., 2022)	China	Cross-sectional data	5529	0-71	■	⊕	■	⊕	■	■	Good
Cohort studies												
34	(Dadi, Miller, Woodman, Azale, & Mwanri, 2021)	Ethiopia	Study data	866	0-72	⊕	■	⊕	⊕	⊕	■	Fair
58	(R. Das et al., 2022)	Bangladesh	ABCD trial	1431	0-23	■	⊕	■	⊕	■	■	Good
11	(Fall et al., 2015)	Brazil, Guatemala, India, Philippines, and South Africa	COHORTs	19403	24**	■	⊕	⊕	⊕	■	⊕	Fair
54	(Le Roux et al., 2019)	South Africa	Cross-sectional data	470	0-24***	■	⊕	■	⊕	■	■	Fair
3	(Schott et al., 2017)	Ethiopia, India	Young lives study	343	0-78	⊕	⊕	■	⊕	■	⊕	Fair
Case control studies												
74	(Abitew, Yalew, Bezabih, & Bazzano, 2020)	Ethiopia	Baseline data: community-based programme records	1273	6-59	■	⊕	⊕	⊕	■	■	Fair
24	(Gebremaryam, Amare, Ayalew, Tigabu, & Menshaw, 2022)	Ethiopia	Bahir Dar City public hospitals	67 cases, 134 controls	6-23	⊕	■	⊕	⊕	⊕	■	Good
43	(Owoaje, Onifade, & Desmennu, 2014)	Nigeria	Oni Memorial Hospital data	100 cases, 200 controls	6-23	⊕	⊕	■	⊕	⊕	■	Fair
25	(Pravana et al., 2017)	Nepal	Community based random selection	146 cases, 146 controls	0-59	⊕	■	⊕	⊕	■	■	Fair
89	(Wemakor, Azongo, Garti, & Atosona, 2018)	Ghana	Study questionnaire	150 cases, 150 controls	6-59	■	⊕	■	⊕	■	■	Fair
Randomised controlled trials												
4	(Friebert, Callaghan-Gillespie, Papathakis, & Manary, 2017)	Malawi	Mamachiponde study	2284	6 weeks and 12 weeks**	■	⊕	■	⊕	■	⊕	Good

■ Yes
⊕ No
-- other studies cited

*disaggregated by maternal age

**cohort study: measurements taken at specific time point

***longitudinal cohort study, measurements taken at 3, 6, 9, 12 and 24 months

^a Moderate wasting: weight-for-height z-score (WHZ) <-2, weight-for-length z-score (WLZ) <-2, mid-upper arm circumference (MUAC) <125mm, and/or bilateral oedema

^b Severe wasting: weight-for-height z-score (WHZ) <-3, weight-for-length z-score (WLZ) <-3, mid-upper arm circumference (MUAC) ≤115mm, and/or bilateral oedema

^c Moderate underweight: weight-for-age z-score (WAZ) <-2

^d Severe underweight: weight-for-age z-score (WAZ) <-3

Supplementary Appendix 3. Reference list for all studies included in the systematic review and meta-analysis

- Abitew, D. B., Yalew, A. W., Bezabih, A. M., & Bazzano, A. N. (2020). Predictors of relapse of acute malnutrition following exit from community-based management program in Amhara region, Northwest Ethiopia: An unmatched case-control study. *PLoS ONE [Electronic Resource]*, 15(4), e0231524. doi:<https://dx.doi.org/10.1371/journal.pone.0231524>
- Aboagye, R. G., Seidu, A. A., Ahinkorah, B. O., Arthur-Holmes, F., Cadri, A., Dadzie, L. K., . . . Yaya, S. (2021). Dietary diversity and undernutrition in children aged 6-23 months in sub-Saharan Africa. *Nutrients*, 13(10). doi:<http://dx.doi.org/10.3390/nu13103431>
- Adhikari, T., Yadav, J., Tripathi, N., Tolani, H., Kaur, H., & Rao, M. V. V. (2021). Do tribal children experience elevated risk of poor nutritional status in India? A multilevel analysis. *Journal of Biosocial Science*, 53(5), 683-708. doi:<https://dx.doi.org/10.1017/S0021932020000474>
- Ahmed, A. S., Ahmed, T., Roy, S. K., Alam, N., & Hossain, M. I. (2012). Determinants of undernutrition in children under 2 years of age from Rural Bangladesh. *Indian Pediatrics*, 49(10), 821-824. doi:<http://dx.doi.org/10.1007/s13312-012-0187-2>
- Akseer, N., Bhatti, Z., Mashal, T., Soofi, S., Moineddin, R., Black, R. E., & Bhutta, Z. A. (2018). Geospatial inequalities and determinants of nutritional status among women and children in Afghanistan: An observational study. *The Lancet Global Health*, 6(4), e447-e459. doi:[https://dx.doi.org/10.1016/S2214-109X\(18\)30025-1](https://dx.doi.org/10.1016/S2214-109X(18)30025-1)
- Ali, N. B., Tahsina, T., Emdadul Hoque, D. M., Hasan, M. M., Iqbal, A., Huda, T. M., & El Arifeen, S. (2019). Association of food security and other socioeconomic factors with dietary diversity and nutritional statuses of children aged 6-59 months in rural Bangladesh. *PLoS ONE [Electronic Resource]*, 14(8) (no pagination). doi:<http://dx.doi.org/10.1371/journal.pone.0221929>
- Alkaff, F. F., Flynn, J., Sukmajaya, W. P., & Salamah, S. (2020). Comparison of WHO growth standard and national Indonesian growth reference in determining prevalence and determinants of stunting and underweight in children under five: A cross-sectional study from Musi sub-district. *F1000Research*, 9. doi:<https://dx.doi.org/10.12688/f1000research.23156.2>
- Amadu, I., Seidu, A. A., Duku, E., Okyere, J., Hagan, J. E., Hormenu, T., & Ahinkorah, B. O. (2021). The joint effect of maternal marital status and type of household cooking fuel on child nutritional status in sub-Saharan Africa: Analysis of cross-sectional surveys on children from 31 countries. *Nutrients*, 13(5). doi:<http://dx.doi.org/10.3390/nu13051541>
- Amare, Z. Y., Ahmed, M. E., & Mehari, A. B. (2019). Determinants of nutritional status among children under age 5 in Ethiopia: Further analysis of the 2016 Ethiopia Demographic and Health Survey. *Globalization and Health*, 15(1). doi:<http://dx.doi.org/10.1186/s12992-019-0505-7>
- Beiersmann, C., Lorenzo, J. B., Bountogo, M., Tiendrebeogo, J., Gabrysch, S., Ye, M., . . . Mu ller, O. (2013). Malnutrition determinants in young children from Burkina Faso. *Journal of Tropical Pediatrics*, 59(5), 372-379. doi:<http://dx.doi.org/10.1093/tropej/fmt037>
- Bekele, T., Rawstorne, P., & Rahman, B. (2021). Socioeconomic inequalities in child growth failure in Ethiopia: Findings from the 2000 and 2016 Demographic and Health Surveys. *BMJ Open*, 11(12). doi:<https://dx.doi.org/10.1136/bmjopen-2021-051304>

- Birhan, N. A., & Belay, D. B. (2021). Associated risk factors of underweight among under-five children in Ethiopia using multilevel ordinal logistic regression model. *African Health Sciences*, 21(1), 362-372. doi:<http://dx.doi.org/10.4314/ahs.v21i1.46>
- Biswas, T., Townsend, N., Magalhaes, R. J. S., Hasan, M., & Mamun, A. (2021). Patterns and determinants of the double burden of malnutrition at the household level in South and Southeast Asia. *European Journal of Clinical Nutrition*, 75(2), 385-391. doi:<https://dx.doi.org/10.1038/s41430-020-00726-z>
- Boah, M., Azupogo, F., Amporfro, D. A., & Abada, L. A. (2019). The epidemiology of undernutrition and its determinants in children under five years in Ghana. *PLoS ONE*, 14(7), e0219665. doi:10.1371/journal.pone.0219665
- Dadi, A. F., Miller, E. R., Woodman, R. J., Azale, T., & Mwanri, L. (2021). Effect of perinatal depression on risk of adverse infant health outcomes in mother-infant dyads in Gondar town: A causal analysis. *BMC Pregnancy & Childbirth*, 21(1), 255. doi:<https://dx.doi.org/10.1186/s12884-021-03733-5>
- Das, R., Kabir, M. F., Ashorn, P., Simon, J., Chisti, M. J., & Ahmed, T. (2022). Maternal underweight and its association with composite index of anthropometric failure among children under two years of age with diarrhea in Bangladesh. *Nutrients*, 14(9). doi:<https://dx.doi.org/10.3390/nu14091935>
- Das, S., Fahim, S. M., Islam, M. S., Biswas, T., Mahfuz, M., & Ahmed, T. (2019). Prevalence and sociodemographic determinants of household-level double burden of malnutrition in Bangladesh. *Public Health Nutrition*, 22(8), 1425-1432. doi:<https://dx.doi.org/10.1017/S1368980018003580>
- Fagbamigbe, A. F., Kandala, N. B., & Uthman, O. A. (2020). Decomposing the educational inequalities in the factors associated with severe acute malnutrition among under-five children in low- and middle-income countries. *BMC Public Health*, 20(1), 555. doi:<https://dx.doi.org/10.1186/s12889-020-08635-3>
- Faisal Ahmed, N. A. M., Sultana, M., Ali, M., Abedin, M. M., Ahammed, B., Yeasmin, M. A., & Maniruzzaman, M. (2021). Identifying the factors causing malnutrition and its impact on mortality among under-five bangladeshi children. *Family Medicine and Primary Care Review*, 23(3), 255-260. doi:<http://dx.doi.org/10.5114/fmpcr.2021.108185>
- Fall, C. H., Sachdev, H. S., Osmond, C., Restrepo-Mendez, M. C., Victora, C., Martorell, R., . . . investigators, C. (2015). Association between maternal age at childbirth and child and adult outcomes in the offspring: A prospective study in five low-income and middle-income countries (COHORTS collaboration). *The Lancet Global Health*, 3(7), e366-377. doi:[https://dx.doi.org/10.1016/S2214-109X\(15\)00038-8](https://dx.doi.org/10.1016/S2214-109X(15)00038-8)
- Finlay, J. E., Ozaltin, E., & Canning, D. (2011). The association of maternal age with infant mortality, child anthropometric failure, diarrhoea and anaemia for first births: Evidence from 55 low- and middle-income countries. *BMJ Open*, 1(2), e000226. doi:<https://dx.doi.org/10.1136/bmjopen-2011-000226>
- Friebert, A., Callaghan-Gillespie, M., Papathakis, P. C., & Manary, M. J. (2017). Adolescent pregnancy and nutrition: A subgroup analysis from the Mamachiponde study in Malawi. *Annals of the New York Academy of Sciences*, 16, 16. doi:<https://dx.doi.org/10.1111/nyas.13465>
- Fuada, N., Latifah, L., Yunitawat, D., & Ashar, H. (2020). Assessment of nutritional status of children under-five in families of adolescent mothers in Indonesia 2013. *Journal of Nutritional Science & Vitaminology*, 66(Supplement), S425-S431. doi:<https://dx.doi.org/10.3177/jns.v66.S425>

- Gbadamosi, M. A., Goon, D. T., & Tugli, A. (2017). Relationship between feeding practices and patterns of infant growth: A cross-sectional study. *Research Journal of Medical Sciences*, 11(4), 166-173. doi:<http://dx.doi.org/10.3923/rjmsci.2017.166.173>
- Gebremaryam, T., Amare, D., Ayalew, T., Tigabu, A., & Menshaw, T. (2022). Determinants of severe acute malnutrition among children aged 6-23 months in Bahir Dar City public hospitals, Northwest Ethiopia, 2020: A case control study. *BMC Pediatrics*, 22(1). doi:<https://dx.doi.org/10.1186/s12887-022-03327-w>
- Geda, N. R., Feng, C. X., Henry, C. J., Lepnurm, R., Janzen, B., & Whiting, S. J. (2021). Multiple anthropometric and nutritional deficiencies in young children in Ethiopia: a multi-level analysis based on a nationally representative data. *BMC Pediatrics*, 21(1), 11. doi:<https://dx.doi.org/10.1186/s12887-020-02467-1>
- Gewa, C. A., & Yandell, N. (2012). Undernutrition among Kenyan children: Contribution of child, maternal and household factors. *Public Health Nutrition*, 15(6), 1029-1038. doi:<https://dx.doi.org/10.1017/S136898001100245X>
- Ghimire, U., Aryal, B. K., Gupta, A. K., & Sapkota, S. (2020). Severe acute malnutrition and its associated factors among children under-five years: A facility-based cross-sectional study. *BMC Pediatrics*, 20(1), 249. doi:<https://dx.doi.org/10.1186/s12887-020-02154-1>
- Haque, R., Alam, K., Rahman, S. M., Mustafa, M. U. R., Ahammed, B., Ahmad, K., . . . Keramat, S. A. (2022). Nexus between maternal underweight and child anthropometric status in South and South-East Asian countries. *Nutrition*, 98. doi:<https://dx.doi.org/10.1016/j.nut.2022.111628>
- Hien, N. N., & Hoa, N. N. (2009). Nutritional status and determinants of malnutrition in children under three years of age in Nghean, Vietnam. *Pakistan Journal of Nutrition*, 8(7), 958-964. doi:<http://dx.doi.org/10.3923/pjn.2009.958.964>
- Hien, N. N., & Kam, S. (2008). Nutritional status and the characteristics related to malnutrition in children under five years of age in Nghean, Vietnam. *Journal of Preventive Medicine & Public Health / Yebang Uihakhoe Chi*, 41(4), 232-240. doi:<https://dx.doi.org/10.3961/jpmpmh.2008.41.4.232>
- Hiruy, A. F., Xiong, Q., Jin, Q., Zhao, J., Lin, X., He, S., . . . Ying, C. (2021). The association of feeding practices and sociodemographic factors on underweight and wasting in children in Ethiopia: A secondary analysis of four health surveys from 2000 to 2016. *Journal of Tropical Pediatrics*, 67(4), 27. doi:<https://dx.doi.org/10.1093/tropej/fmab047>
- Horta, B. L., Santos, R. V., Welch, J. R., Cardoso, A. M., Dos Santos, J. V., Assis, A. M. O., . . . Coimbra Jr, C. E. A. (2013). Nutritional status of indigenous children: Findings from the First National Survey of Indigenous People's Health and Nutrition in Brazil. *International Journal for Equity in Health*, 12(1). doi:<http://dx.doi.org/10.1186/1475-9276-12-23>
- Hossain, F. B., Shawon, M. S. R., Al-Abid, M. S. U., Mahmood, S., Adhikary, G., & Bulbul, M. M. I. (2020). Double burden of malnutrition in children aged 24 to 59 months by socioeconomic status in five South Asian countries: Evidence from demographic and health surveys. *BMJ Open*, 10(3) (no pagination). doi:<http://dx.doi.org/10.1136/bmjopen-2019-032866>
- Hossain, M. B., & Khan, M. H. R. (2018). Role of parental education in reduction of prevalence of childhood undernutrition in Bangladesh. *Public Health Nutrition*, 21(10), 1845-1854. doi:<https://dx.doi.org/10.1017/S1368980018000162>
- Huynh, G., Huynh, Q. H. N., Nguyen, N. H. T., Do, Q. T., & Khanh Tran, V. (2019). Malnutrition among 6-59-month-old children at District 2 Hospital, Ho Chi Minh

- City, Vietnam: Prevalence and associated factors. *BioMed Research International*, 2019, 6921312. doi:<http://dx.doi.org/10.1155/2019/6921312>
- Ickes, S. B., Hurst, T. E., & Flax, V. L. (2015). Maternal literacy, facility birth, and education are positively associated with better infant and young child feeding practices and nutritional status among Ugandan children. *Journal of Nutrition*, 145(11), 2578-2586. doi:<http://dx.doi.org/10.3945/jn.115.214346>
- Islam, M. S., & Biswas, T. (2020). Prevalence and correlates of the composite index of anthropometric failure among children under 5 years old in Bangladesh. *Maternal & Child Nutrition*, 16(2), e12930. doi:<https://dx.doi.org/10.1111/mcn.12930>
- Issah, A. N., Yeboah, D., Kpordoxah, M. R., Boah, M., & Mahama, A. B. (2022). Association between exposure to intimate partner violence and the nutritional status of women and children in Nigeria. *PLoS ONE [Electronic Resource]*, 17(5), e0268462. doi:<https://dx.doi.org/10.1371/journal.pone.0268462>
- Janevic, T., Petrovic, O., Bjelic, I., & Kubera, A. (2010). Risk factors for childhood malnutrition in Roma settlements in Serbia. *BMC Public Health*, 10, 509. doi:<https://dx.doi.org/10.1186/1471-2458-10-509>
- Karim, M. R., Al Mamun, A. S. M., Rana, M. M., Mahumud, R. A., Shoma, N. N., Dutt, D., . . . Hossain, M. G. (2021). Acute malnutrition and its determinants of preschool children in Bangladesh: Gender differentiation. *BMC Pediatrics*, 21(1), 573. doi:<https://dx.doi.org/10.1186/s12887-021-03033-z>
- Kasaye, H. K., Bobo, F. T., Yilma, M. T., & Woldie, M. (2019). Poor nutrition for under-five children from poor households in Ethiopia: Evidence from 2016 Demographic and Health Survey. *PLoS ONE*, 14(12), e0225996. doi:10.1371/journal.pone.0225996
- Khan, M. S., Halder, H. R., Rashid, M., Afroja, S., & Islam, M. (2020). Impact of socioeconomic and demographic factors for underweight and overweight children in Bangladesh: A polytomous logistic regression model. *Clinical Epidemiology and Global Health*, 8(4), 1348-1355. doi:<http://dx.doi.org/10.1016/j.cegh.2020.05.010>
- Khan, S., Zaheer, S., & Safdar, N. F. (2019). Determinants of stunting, underweight and wasting among children <5 years of age: Evidence from 2012-2013 Pakistan Demographic and Health Survey. *BMC Public Health*, 19(1), 358. doi:<https://dx.doi.org/10.1186/s12889-019-6688-2>
- Kim, R., Rajpal, S., Joe, W., Corsi, D. J., Sankar, R., Kumar, A., & Subramanian, S. V. (2019). Assessing associational strength of 23 correlates of child anthropometric failure: An econometric analysis of the 2015-2016 National Family Health Survey, India. *Social Science & Medicine*, 238, 112374. doi:<https://dx.doi.org/10.1016/j.socscimed.2019.112374>
- Kumar, A., & Ram, F. (2013). Influence of family structure on child health: Evidence from India. *Journal of Biosocial Science*, 45(5), 577-599. doi:<https://dx.doi.org/10.1017/S0021932012000764>
- Kumar, R., Abbas, F., Mahmood, T., & Somrongthong, R. (2019). Prevalence and factors associated with underweight children: A population-based subnational analysis from Pakistan. *BMJ Open*, 9(7). doi:<http://dx.doi.org/10.1136/bmjopen-2019-028972>
- Kumar, R., & Paswan, B. (2021). Changes in socio-economic inequality in nutritional status among children in EAG states, India. *Public Health Nutrition*, 24(6), 1304-1317. doi:<https://dx.doi.org/10.1017/S1368980021000343>
- Le Roux, K., Christodoulou, J., Stansert-Katzen, L., Dippenaar, E., Laurenzi, C., Le Roux, I. M., . . . Rotheram-Borus, M. J. (2019). A longitudinal cohort study of rural adolescent vs adult South African mothers and their children from birth to 24 months. *BMC Pregnancy and Childbirth*, 19(1). doi:<http://dx.doi.org/10.1186/s12884-018-2164-8>

- Li, H., Yuan, S., Fang, H., Huang, G., Huang, Q., Wang, H., & Wang, A. (2022). Prevalence and associated factors for stunting, underweight and wasting among children under 6 years of age in rural Hunan Province, China: A community-based cross-sectional study. *BMC Public Health*, 22(1), 483. doi:<https://dx.doi.org/10.1186/s12889-022-12875-w>
- Linnemayr, S., Alderman, H., & Ka, A. (2008). Determinants of malnutrition in Senegal: Individual, household, community variables, and their interaction. *Economics & Human Biology*, 6(2), 252-263. doi:<https://dx.doi.org/10.1016/j.ehb.2008.04.003>
- Mashal, T., Takano, T., Nakamura, K., Kizuki, M., Hemat, S., Watanabe, M., & Seino, K. (2008). Factors associated with the health and nutritional status of children under 5 years of age in Afghanistan: Family behaviour related to women and past experience of war-related hardships. *BMC Public Health*, 8. doi:<http://dx.doi.org/10.1186/1471-2458-8-301>
- Masibo, P. K., Humwa, F., & Macharia, T. N. (2020). The double burden of overnutrition and undernutrition in mother-child dyads in Kenya: Demographic and Health Survey data, 2014. *Journal of Nutritional Science*, 9, e5. doi:<https://dx.doi.org/10.1017/jns.2019.39>
- Mena-Melendez, L. (2020). Ethnoracial child health inequalities in Latin America: Multilevel evidence from Bolivia, Colombia, Guatemala, and Peru. *SSM - Population Health*, 12. doi:<http://dx.doi.org/10.1016/j.ssmph.2020.100673>
- Mokwena, K., & Kachabe, J. (2022). Profile of mothers whose children are treated for malnutrition at a rural district hospital in the North West province, South Africa. *South African Journal of Clinical Nutrition*, 35(1), 17-22. doi:<https://dx.doi.org/10.1080/16070658.2021.1921899>
- Mutunga, M., Frison, S., Rava, M., & Bahwere, P. (2020). The forgotten agenda of wasting in Southeast Asia: Burden, determinants and overlap with stunting: A review of nationally representative cross-sectional Demographic and Health Surveys in six countries. *Nutrients*, 12(2). doi:<http://dx.doi.org/10.3390/nu12020559>
- Nakamori, M., Ninh, N. X., Khan, N. C., Huong, C. T., Tuan, N. A., Mai, L. B., . . . Yamamoto, S. (2010). Nutritional status, feeding practice and incidence of infectious diseases among children aged 6 to 18 months in northern mountainous Vietnam. *Journal of Medical Investigation*, 57(1-2), 45-53. doi:<http://dx.doi.org/10.2152/jmi.57.45>
- Nguyen, P. H., Sanghvi, T., Tran, L. M., Afsana, K., Mahmud, Z., Aktar, B., . . . Menon, P. (2017). The nutrition and health risks faced by pregnant adolescents: Insights from a cross-sectional study in Bangladesh. *PLoS ONE*, 12(6), e0178878. doi:<https://dx.doi.org/10.1371/journal.pone.0178878>
- Nguyen, P. H., Scott, S., Khuong, L. Q., Pramanik, P., Ahmed, A., Rashid, S. F., . . . Menon, P. (2021). Adolescent birth and child undernutrition: An analysis of Demographic and Health Surveys in Bangladesh, 1996-2017. *Annals of the New York Academy of Sciences*, 1500(1), 69-81. doi:<https://dx.doi.org/10.1111/nyas.14608>
- Nguyen, P. H., Scott, S., Neupane, S., Tran, L. M., & Menon, P. (2019). Social, biological, and programmatic factors linking adolescent pregnancy and early childhood undernutrition: A path analysis of India's 2016 National Family and Health Survey. *The Lancet Child & Adolescent Health*, 3(7), 463-473. doi:[https://dx.doi.org/10.1016/S2352-4642\(19\)30110-5](https://dx.doi.org/10.1016/S2352-4642(19)30110-5)
- Ntenda, P. A. M. (2019). Association of low birth weight with undernutrition in preschool-aged children in Malawi. *Nutrition Journal*, 18(1). doi:<http://dx.doi.org/10.1186/s12937-019-0477-8>

- Ntenda, P. A. M., & Chuang, Y. C. (2018). Analysis of individual-level and community-level effects on childhood undernutrition in Malawi. *Pediatrics & Neonatology*, 59(4), 380-389. doi:<https://dx.doi.org/10.1016/j.pedneo.2017.11.019>
- Obayelu, O. A., & Adeleye, O. R. (2021). Explaining child nutritional status in rural Nigeria: Socioeconomic dimensions. *Journal of Hunger and Environmental Nutrition*, 16(6), 829-846. doi:<http://dx.doi.org/10.1080/19320248.2020.1781011>
- Olodu, M. D., Adeyemi, A. G., Olowookere, S. A., & Esimai, O. A. (2019). Nutritional status of under-five children born to teenage mothers in an urban setting, south-western Nigeria. *BMC Research Notes*, 12(1), 116. doi:<https://dx.doi.org/10.1186/s13104-019-4147-x>
- Owoaje, E., Onifade, O., & Desmennu, A. (2014). Family and socioeconomic risk factors for undernutrition among children aged 6 to 23 Months in Ibadan, Nigeria. *The Pan African medical journal*, 17, 161. doi:<https://dx.doi.org/10.11604/pamj.2014.17.161.2389>
- Paul, P., Chouhan, P., & Zaveri, A. (2019). Impact of child marriage on nutritional status and anaemia of children under 5 years of age: Empirical evidence from India. *Public Health*, 177, 95-101. doi:<https://dx.doi.org/10.1016/j.puhe.2019.08.008>
- Paul, P., & Saha, R. (2022). Is maternal autonomy associated with child nutritional status? Evidence from a cross-sectional study in India. *PLoS ONE*, 17(5), e0268126. doi:<https://dx.doi.org/10.1371/journal.pone.0268126>
- Poda, G. G., Hsu, C. Y., & Chao, J. C. J. (2017). Factors associated with malnutrition among children <5 years old in Burkina Faso: Evidence from the Demographic and Health Surveys IV 2010. *International Journal for Quality in Health Care*, 29(7), 901-908. doi:<http://dx.doi.org/10.1093/intqhc/mzx129>
- Porwal, A., Agarwal, P. K., Ashraf, S., Acharya, R., Ramesh, S., Khan, N., . . . Sarna, A. (2021). Association of maternal height and body mass index with nutrition of children under 5 years of age in India: Evidence from Comprehensive National Nutrition Survey 2016-18. *Asia Pacific Journal of Clinical Nutrition*, 30(4), 675-686. doi:[https://dx.doi.org/10.6133/apjcn.202112_30\(4\).0014](https://dx.doi.org/10.6133/apjcn.202112_30(4).0014)
- Pramod Singh, G. C., Nair, M., Grubestic, R. B., & Connell, F. A. (2009). Factors associated with underweight and stunting among children in rural terai of Eastern Nepal. *Asia-Pacific Journal of Public Health*, 21(2), 144-152. doi:<https://dx.doi.org/10.1177/1010539509332063>
- Pravana, N. K., Piryani, S., Chaurasiya, S. P., Kawan, R., Thapa, R. K., & Shrestha, S. (2017). Determinants of severe acute malnutrition among children under 5 years of age in Nepal: A community-based case-control study. *BMJ Open*, 7(8). doi:<http://dx.doi.org/10.1136/bmjopen-2017-017084>
- Qu, P., Mi, B., Wang, D., Zhang, R., Yang, J., Liu, D., . . . Yan, H. (2017). Association between the Infant and Child Feeding Index (ICFI) and nutritional status of 6- to 35-month-old children in rural western China. *PLoS ONE*, 12(2). doi:<http://dx.doi.org/10.1371/journal.pone.0171984>
- Rahman, A., & Hossain, M. M. (2022). Quantile regression approach to estimating prevalence and determinants of child malnutrition. *Journal of Public Health*, 30(2), 323-339. doi:<https://dx.doi.org/10.1007/s10389-020-01277-0>
- Rahman, M. A., Halder, H. R., Rahman, M. S., & Parvez, M. (2021). Poverty and childhood malnutrition: Evidence-based on a nationally representative survey of Bangladesh. *PLoS ONE*, 16(8), e0256235. doi:<https://dx.doi.org/10.1371/journal.pone.0256235>
- Rahman, M. M. (2015). Is unwanted birth associated with child malnutrition in Bangladesh? *International Perspectives on Sexual & Reproductive Health*, 41(2), 80-88. doi:<https://dx.doi.org/10.1363/4108015>

- Rahman, M. S., Rahman, M. A., Maniruzzaman, M., & Howlader, M. H. (2020). Prevalence of undernutrition in Bangladeshi children. *Journal of Biosocial Science*, 52(4), 596-609. doi:<https://dx.doi.org/10.1017/S0021932019000683>
- Raj, A., Saggurthi, N., Winter, M., Labonte, A., Decker, M. R., Balaiah, D., & Silverman, J. G. (2010). The effect of maternal child marriage on morbidity and mortality of children under 5 in India: Cross sectional study of a nationally representative sample. *BMJ*, 340, b4258. doi:<https://dx.doi.org/10.1136/bmj.b4258>
- Rodgers, J., Kim, R., & Subramanian, S. V. (2020). Explaining within- vs between-population variation in child anthropometry and hemoglobin measures in India: A multilevel analysis of the National Family Health Survey 2015-2016. *Journal of Epidemiology*, 30(11), 485-496. doi:<https://dx.doi.org/10.2188/jea.JE20190064>
- Samuel, A., Osendarp, S. J. M., Feskens, E. J. M., Lelisa, A., Adish, A., Kebede, A., & Brouwer, I. D. (2022). Gender differences in nutritional status and determinants among infants (6-11 m): A cross-sectional study in two regions in Ethiopia. *BMC public health*, 22(1), 401. doi:<https://dx.doi.org/10.1186/s12889-022-12772-2>
- Sanjay, C., Gaurav, M., Suman, B., & Ramchandra, C. (2020). Determinants of nutritional status in children aged 0 - 6 years in rural Rajasthan. *Journal of Evolution of Medical and Dental Sciences*, 9(28), 1977-1983. doi:<http://dx.doi.org/10.14260/jemds/2020/431>
- Schott, W., Aurino, E., Penny, M. E., & Behrman, J. R. (2017). Adolescent mothers' anthropometrics and grandmothers' schooling predict infant anthropometrics in Ethiopia, India, Peru, and Vietnam. *Annals of the New York Academy of Sciences*, 24, 24. doi:<https://dx.doi.org/10.1111/nyas.13455>
- Sobkoviak, R. M., Yount, K. M., & Halim, N. (2012). Domestic violence and child nutrition in Liberia. *Social Science & Medicine*, 74(2), 103-111. doi:<https://dx.doi.org/10.1016/j.socscimed.2011.10.024>
- Subramanian, S. V., Ackerson, L. K., & Smith, G. D. (2010). Parental BMI and childhood undernutrition in India: An assessment of intrauterine influence. *Pediatrics*, 126(3), e663-671. doi:<https://dx.doi.org/10.1542/peds.2010-0222>
- Subramanyam, M. A., Kawachi, I., Berkman, L. F., & Subramanian, S. V. (2010). Socioeconomic inequalities in childhood undernutrition in India: Analyzing trends between 1992 and 2005. *PLoS ONE*, 5(6). doi:<http://dx.doi.org/10.1371/journal.pone.0011392>
- Sunil, T. S. (2009). Effects of socio-economic and behavioural factors on childhood malnutrition in Yemen. *Maternal & Child Nutrition*, 5(3), 251-259. doi:<https://dx.doi.org/10.1111/j.1740-8709.2008.00174.x>
- Tariq, J., Sajjad, A., Zakar, R., Zakar, M. Z., & Fischer, F. (2018). Factors associated with undernutrition in children under the age of two years: Secondary data analysis based on the Pakistan Demographic and Health Survey 2012-2013. *Nutrients*, 10(6). doi:<http://dx.doi.org/10.3390/nu10060676>
- Tesfaw, L. M., & Dessie, Z. G. (2022). Multilevel multivariate analysis on the anthropometric indicators of under-five children in Ethiopia: EMDHS 2019. *BMC Pediatrics*, 22(1). doi:<https://dx.doi.org/10.1186/s12887-022-03172-x>
- Tesfaw, L. M., & Fenta, H. M. (2021). Multivariate logistic regression analysis on the association between anthropometric indicators of under-five children in Nigeria: NDHS 2018. *BMC Pediatrics*, 21(1), 193. doi:<https://dx.doi.org/10.1186/s12887-021-02657-5>
- Tibebu, N. S., Emiru, T. D., Tiruneh, C. M., Getu, B. D., & Azanaw, K. A. (2020). Underweight and its associated factors among children 6-59 months of age in Debre

- Tabor Town, Amhara Region of Ethiopia, 2019: A community-based cross-sectional study. *Pediatric Health Med Ther*, 11, 469-476. doi:10.2147/phmt.S288071
- Tiwari, I., Acharya, K., Paudel, Y. R., Sapkota, B. P., & Kafle, R. B. (2020). Planning of births and childhood undernutrition in Nepal: evidence from a 2016 national survey. *BMC Public Health*, 20(1), 1788. doi:<https://dx.doi.org/10.1186/s12889-020-09915-8>
- Wemakor, A., Azongo, T., Garti, H., & Atosona, A. (2018). Young maternal age is a risk factor for child undernutrition in Tamale Metropolis, Ghana. *BMC research notes*, 11(1), 877. doi:<http://dx.doi.org/10.1186/s13104-018-3980-7>

Supplementary Appendix 4. Risk of bias and quality assessments

a. Cross-sectional and cohort studies

ID	Year of Publication	Data Source	Questions														Overall Rating
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	
Cross-sectional studies																	
1	2021	Cross-sectional study data	■	■	■	■	■	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
2	2021	DHS Data 1996 - 2017	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
5	2003	India's NFHS-I data 1995	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
6	2017	Malawi DHS 2004 and 2010	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
7	2019	India NFHS - 4 2015 - 16	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
8	2020	Basic health research data 2013 (RISKESDAS 2013)	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	⊕	⊕	N/A	⊕	Fair
9	2021	Ethiopia DHS 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Fair
10	2022	Nigeria DHS 2018	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
12	2017	Cross sectional survey conducted as part of the Rural Primary Health Care project	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
13	2021	Ethiopia DHS 2000 - 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	N/A	■	⊕	N/A	■	Good
14	2019	Baseline data collected to assess the Nobokoli programme	■	■	■	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Fair
15	2019	Malawi DHS 2015 -16	■	■	■	■	⊕	■	■	■	■	⊕	■	⊕	N/A	■	Good
16	2011	118 DHS surveys conducted in 55 countries between 1990 - 2008	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
17	2021	Comprehensive National Nutrition Survey (CNNS) India	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
18	2021	India's NFHS-3 2005-2006, and NFHS-4 2015-2016	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
19	2020	Cross-sectional survey and secondary data from child health books and health record books	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
20	2020	DHS survey data from 51 countries 2010 - 2018	■	⊕	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
21	2008	Baseline data collected for a nutrition intervention programme	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good

22	2019	Ethiopia DHS 2016	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
23	2020	Cross-sectional survey	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	⊕	Fair
26	2019	Pakistan DHS 2012 - 2013	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
27	2012	Bangladesh National Nutrition Programme baseline survey 2004	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	⊕	Fair
28	2021	DHS survey data from 32 countries of Sub-Saharan Africa 2010 - 2020	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
29	2021	India NFHS - 4 2015 - 16	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
30	2012	Liberia DHS survey 2006-2007	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
31	2020	DHS surveys from Bangladesh, India, Pakistan, Maldives and Nepal between 2009 - 2016	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
32	2020	Kenya DHS survey 2014	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
33	2010	India's National Family Health Survey 2005 - 2006	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
35	2009	Yemen DHS survey 1997	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
36	2019	Ghana DHS 2014	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
37	2020	DHS data for Bolivia, Colombia, Guatemala and Peru 1986 - 2015	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
38	2020	India NFHS - 4 2015 - 16	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
39	2017	Burkina-Faso DHS 2010	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
40	2008	Cross-sectional survey data	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
41	2018	PDHS 2012 - 2013	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
42	2009	Cross-sectional survey data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
		Cambodia DHS 2014, Lao PDR Multiple Indicator Survey (MICS) 2017, Myanmar DHS 2015, Thailand MICS 2015/16, Timor-Leste National Food and Nutrition Survey (NFNS) 2013, and Vietnam MICS 2011															
44	2020	Baseline data collected for a large effectiveness study on the use of multi-micronutrient powders	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
45	2022	ANNS 2013 (Afghanistan National Nutrition Survey)	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
46	2018	Bangladesh DHS 2014	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
47	2021	India NFHS - 4 2015 - 16	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
48	2019	Bangladesh DHS 2014	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good
49	2020		■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	■	N/A	■	Good

50	2013	India's NFHS 2005 - 2006	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
51	2022	India NFHS - 4 2015 - 16	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
52	2015	BDHS 2011	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
53	2021	DHS Data for 31 countries 2010 - 2019	■	⊕	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
55	2019	Cross-sectional study data	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
56	2013	Cross-sectional study data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
57	2015	Uganda DHS 2006 and 2011	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
59	2022	EMDHS 2019 (Ethiopian mini DHS)	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
60	2021	Ethiopia DHS 2016	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
61	2021	Nigeria DHS 2018	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
62	2022	DHS Data 2014 - 2018	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
63	2017	Baseline data from a scaled up MNCH programme	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
64	2009	Cross-sectional study data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
65	2008	Cross-sectional study data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
66	2013	National Survey of Indigenous People's Health and Nutrition in Brazil 2008 - 2009	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
67	2019	Cross-sectional survey using primary health care centres records as the sampling frame	■	■	■	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
68	2010	Cross-sectional study data	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	⊕	Fair
69	2010	India's NFHS-3 2005-2006, and NFHS-4 2015-2016	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
70	2021	DHS Survey data for Bangladesh, India, Nepal, Pakistan, Myanmar, Timor, Maldives and Cambodia conducted between 2007 and 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
71	2020	Nepal DHS 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
72	2019	Ethiopia DHS 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
73	2021	Bangladesh DHS 2017-18	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
75	2022	Cross-sectional study data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
76	2020	Bangladesh DHS 2014	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
77	2019	MICS 2014	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
78	2019	Bangladesh DHS 2014	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
79	2020	Bangladesh DHS 2014	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good

80	2022	Cross-sectional study data	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	⊕	Fair
81	2017	Cross-sectional data from the MAL-ED Cohort study	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
82	2010	Serbia MICS 2005	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
83	2018	Bangladesh DHS 2014	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
84	2020	Cross-sectional study data from children admitted to the Outpatients Therapeutic Care Centers	■	■	■	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
85	2019	India's NFHS - 4 2015 - 16	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
86	2021	Ethiopia DHS 2000 and 2016	■	■	⊕	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
87	2010	India's NFHS 1992, 1998 and 2005	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
88	2012	Kenya DHS survey 2003	■	■	■	■	⊕	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
90	2021	Nigeria DHS 2013	■	■	⊕	■	⊕	■	⊕	■	■	⊕	■	⊕	N/A	■	Good
91	2022	Bangladesh DHS 2014	■	■	⊕	■	⊕	■	⊕	■	■	⊕	■	⊕	N/A	■	Good
92	2020	Cross-sectional study data	■	■	■	■	■	⊕	⊕	■	■	⊕	■	⊕	N/A	■	Good
Cohort Studies																	
3	2017	Secondary analysis of data from the Young lives study	■	■	■	■	⊕	■	■	■	■	■	■	⊕	N/A	■	Fair
11	2015	Data from COHORTs (Consortium for Health Orientated Research in Transitioning Societies)	■	■	■	■	⊕	■	■	■	■	■	■	⊕	■	■	Fair
34	2021	Study data	■	■	⊕	■	■	⊕	⊕	■	■	⊕	■	⊕	■	■	Fair
54	2019	Cross-sectional study data	■	■	■	■	⊕	■	■	■	■	■	■	⊕	■	■	Fair
58	2022	Data from ABCD trial Bangladesh Sit	■	■	⊕	■	■	■	■	■	■	⊕	■	⊕	■	■	Good

■ = Yes

⊕ = No / not reported

Questions:

1. Was the research question or objective in this paper clearly stated?
2. Was the study population clearly specified and defined?
3. Was the participation rate of eligible persons at least 50%?
4. Were all the subjected selected or recruited from the same or similar populations (including the same time period)? Were inclusions and exclusion criteria for being in the study prespecified and applied uniformly to all participants?

5. Was a sample size justification, power description, or variance and effect estimates provided?
6. For the analyses in this paper, were the exposure(s) of interest measured prior to the outcome(s) being measured?
7. Was the timeframe sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?
8. For the exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcomes e.g. categories of exposure or exposure measured as a continuous variable
9. Were the exposure measures (independent variables) clearly defined, valid, reliable and implemented consistently across all study participants?
10. Were the exposures measures assessed more than once over time?
11. Were the outcome measures clearly defined, valid, reliable and implemented consistently across all participants?
12. Were the outcome assessors blinded to the exposure status of the participants?
13. Was loss to follow-up after baseline 20% or less?
14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposure(s) and outcome(s)?

b. Case control studies

ID	Year of Publication	Data Source	1	2	3	4	5	6	7	8	9	10	11	12	Overall Rating
24	2022	Bahir Dar City public hospitals	■	■	■	■	■	■	■	■	⊖	■	⊕	■	Good
25	2017	Community based random selection	■	■	■	■	■	■	⊕	■	■	■	⊕	⊖	Fair
43	2014	Oni Memorial Hospital data	■	■	■	■	■	■	⊕	⊕	⊖	⊕	⊕	⊖	Fair
74	2020	Baseline data from community-based management programme records assessing factors associated with relapse	■	■	■	■	■	■	⊕	⊕	■	■	⊕	No	Fair
89	2018	Study questionnaire	■	■	■	■	■	■	⊕	⊕	■	■	⊕	■	Fair

■ = Yes

⊕ = No / not reported

⊖ = Partial

Questions:

1. Was the research question or objective in this paper clearly stated?
2. Was the study population clearly specified and defined?
3. Did the authors include a sample size justification?
4. Were the controls selected or recruited from the same or similar population that gave rise to the cases (including the same timeframe)?
5. Were the definitions, inclusion and exclusion criteria, algorithms or processes used to identify or select cases and controls valid, reliable and implemented consistently across all study participants?
6. Were the cases clearly defined and differentiated from controls?
7. If less than 100 percent of eligible cases and/or controls were selected from the study, were the cases and/or controls randomly selected from those eligible?
8. Was there use of concurrent controls?
9. Were the investigators able to confirm that the exposure / risk occurred prior to the development of the condition or event that defined a participant as a case?
10. Were the measures of exposure/risk clearly defined, valid, reliable, and implemented consistently (including the same time period) across all study participants?
11. Were the assessors of exposure / risk blinded to the case or control status of the participants?
12. Were key potential confounding variables measured and adjusted statistically in the analyses? If matching was used, did the investigators account for matching during study analysis?

c. Randomised controlled trials

ID	Year of Publication	Data Source	Questions														Overall Rating
			1	2	3	4	5	6	7	8	9	10	11	12	13	14	
4	2017	Subgroup analysis from the Mamachiponde study	■	■	■	⊕	■	■	■	■	■	■	■	■	⊙	■	Good

■ = Yes


⊕ = No / not reported

⊙ = Partial

Questions:


1. Was the study described as randomized, a randomized trial, a randomized clinical trial, or an RCT?
2. Was the method of randomization adequate (i.e., use of randomly generated assignment)?
3. Was the treatment allocation concealed (so that assignments could not be predicted)?
4. Were study participants and providers blinded to treatment group assignment?
5. Were the people assessing the outcomes blinded to the participants' group assignments?
6. Were the groups similar at baseline on important characteristics that could affect outcomes (e.g., demographics, risk factors, co-morbid conditions)?
7. Was the overall drop-out rate from the study at endpoint 20% or lower of the number allocated to treatment?
8. Was the differential drop-out rate (between treatment groups) at endpoint 15 percentage points or lower?
9. Was there high adherence to the intervention protocols for each treatment group?
10. Were other interventions avoided or similar in the groups (e.g., similar background treatments)?
11. Were outcomes assessed using valid and reliable measures, implemented consistently across all study participants?
12. Did the authors report that the sample size was sufficiently large to be able to detect a difference in the main outcome between groups with at least 80% power?
13. Were outcomes reported or subgroups analysed prespecified (i.e., identified before analyses were conducted)?
14. Were all randomized participants analysed in the group to which they were originally assigned, i.e., did they use an intention-to-treat analysis?

Supplementary Figure 1. Association between adolescent pregnancy (10-19 years) and the pooled odds of moderate childhood wasting (1-59 months), versus adult pregnancy




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) <-2 to -3 or mid-upper arm circumference (MUAC) 125mm to >115mm), no – number of children without the outcome

Supplementary Figure 2. Association between adolescent pregnancy (10-19 years) and the pooled odds of severe childhood wasting (1-59 months), versus adult pregnancy




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (severe wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) < -3 or mid-upper arm circumference (MUAC) ≤ 115 mm), no – number of children without the outcome

Supplementary Figure 3. Association between adolescent pregnancy (10-19 years) and the pooled odds of moderate childhood underweight (1-59 months), versus adult pregnancy




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate underweight, weight-for-age z-score (WAZ) <-2 to -3), no – number of children without the outcome

Supplementary Figure 4. Association between adolescent pregnancy (10-24 years) and the pooled odds of moderate childhood wasting (1-59 months), versus adult pregnancy, stratified by region (Asia)




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) <-2 to -3 or mid-upper arm circumference (MUAC) 125mm to >115mm), no – number of children without the outcome

Supplementary Figure 5. Association between adolescent pregnancy (10-24 years) and the pooled odds of moderate childhood underweight (1-59 months), versus adult pregnancy, stratified by region (Asia)




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate underweight, weight-for-age z-score (WAZ) <-2 to -3), no – number of children without the outcome

Supplementary Figure 6. Association between adolescent pregnancy (10-24 years) and the pooled odds of severe childhood wasting (1-59 months), versus adult pregnancy, stratified by region (Asia)




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (severe wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) <-3 or mid-upper arm circumference (MUAC) ≤ 115 mm), no – number of children without the outcome

Supplementary Figure 7. Association between adolescent pregnancy (10-24 years) and the pooled odds of severe childhood underweight (1-59 months), versus adult pregnancy, stratified by region (Asia)




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (severe underweight, weight-for-age z-score (WAZ) <-3), no – number of children without the outcome

Supplementary Figure 8. Association between adolescent pregnancy (10-24 years) and the pooled odds of moderate childhood wasting (1-59 months), versus adult pregnancy, stratified by region (sub-Saharan Africa)




Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) <-2 to -3 or mid-upper arm circumference (MUAC) 125mm to >115mm), no – number of children without the outcome

Supplementary Figure 9. Association between adolescent pregnancy (10-24 years) and the pooled odds of moderate childhood underweight (1-59 months), versus adult pregnancy, stratified by region (sub-Saharan Africa)



Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (moderate underweight, weight-for-age z-score (WAZ) <-2 to -3), no – number of children without the outcome

Supplementary Figure 10. Association between adolescent pregnancy (10-24 years) and the pooled odds of severe childhood wasting (1-59 months), versus adult pregnancy, stratified by region (sub-Saharan Africa)



Key: Treatment – number of children in adolescent mother group, control – number of children in adult mother group, yes – number of children with the outcome (severe wasting, weight-for-height z-score (WHZ) or weight-for-length z-score (WLZ) <-3 or mid-upper arm circumference (MUAC) ≤ 115 mm), no – number of children without the outcome

Supplementary Appendix 5. Summary of findings from studies included in the qualitative synthesis

a. Moderate and severe wasting

Ref	First Author	Year	Study Design	Infant Age categories (months)	Country	Study Size	Maternal age category <20 years	Key findings	Indication of childhood wasting outcomes*
Moderate wasting									
2	Nguyen	2021	Cross-sectional	0 - 59	Bangladesh	12,006	■	Adjusted regression model for age at first birth: 10 – 15 years: -0.03 (-0.12, 0.06) p = not significant 16 – 19 years: 0.01 (-0.05, 0.07) p = not significant	↔
4	Friebert	2017	RCT	Outcomes measured at 6 weeks & 12 weeks	Malawi	2284	■	Mean WLZ (SD) At 6 weeks: Mother <18 years: 0.3 (±1.1) Mother 18 - 20 years: 0.4 (±1.0) Mother >20 years: 0.3 (±1.1) P value = 0.33 At 12 weeks: Mother <18 years: 0.3 (±1.0) Mother 18 - 20 years: 0.3 (±1.0) Mother >20 years: 0.1 (±1.1) p = 0.01 NB: all mothers were malnourished	↔
8	Fuada	2020	Cross-sectional	0 - 59	Indonesia	978	■	OR for adolescent mothers ≤15 years compared to mothers aged 16-19 years: 1.810 (0.648 - 5.055) p = 0.402	↑
10	Issah	2022	Cross-sectional	0 - 59	Nigeria	2425	⊕	aOR 15 – 24 years: reference group 25 - 34: 0.66 (0.32 - 1.36) P = 0.260 35 - 49: 1.16 (0.38 - 3.56) p = 0.798	↑
11	Fall	2015	Cohort	Outcomes measured at 2 years	Brazil, Guatemala, India, the Philippines,	19403	■	aORs ≤19 years: 1.12 (0.86 – 1.46) 20 – 24 years: reference group 25 – 29 years: 0.91 (0.75 0 1.11) 30 – 34 years: 1.05 (0.84 – 1.31)	↑

					and South Africa			≥35 years: 0.97 (0.74 – 1.28) P = 0.64	
13	Hiruy	2021	Cross-sectional	6 - 23	Ethiopia	8003	■	The study shows the trend in aOR over the years: 15 – 19 years: Reference Group 2000 20 – 29 years: 1.03 (0.66-1.63) 30 – 39 years: 1.19 (0.74 – 1.90) 40 – 49 years: 1.06 (0.59 – 1.92) 2005 20 – 29 years: 0.97 (0.43 – 2.18) 30 – 39 years: 1.41 (0.61 – 3.25) 40 – 49 years: 1.79 (0.62 – 5.21) 2011 20 – 29 years: 1.02 (0.65 – 1.60) 30 – 39 years: 1.09 (0.69 – 1.75) 40 – 49 years: 1.36 (0.75 – 2.45) 2016 20 – 29 years: 0.75 (0.45 – 1.24) 30 – 39 years: 0.55* (0.34 – 0.90) 40 – 49 years: 0.84 (0.44 – 1.61) *p = <0.05	More recently: ↑↑
14	Ali	2019	Cross-sectional	0 - 59	Bangladesh	6468	⊕	aOR 15 – 24 years: reference group 25 - 49: 1.06 p = 0.419	↓↓
27	Ahmed	2012	Cross-sectional	0 - 24	Bangladesh	8858	■	aOR <20 years: 1.12 (0.81 – 1.57) 20 – 30 years: reference group >30 years: 1.38 (0.96 – 1.99)	↑↑
30	Sobkoviak	2012	Cross-sectional	0 - 48	Liberia	2467	⊕	aOR 15 – 24 years: reference group 25 - 34: 1.45 (SE: 0.44) 35 - 49: 0.68 (SE: 0.34)	↓↓
32	Masibo	2020	Cross-sectional	0 - 59	Kenya	7830	■	uOR <-2SD Overweight/obese mother – wasted child <20 years: 0.73 (0.11 – 4.87) p 0.749 20 – 30 years: 0.94 (0.29 – 3.03) p 0.918 30 – 40 years 1.49 (0.45 – 4.94) p 0.509 40+ years: reference group	↓↓

36	Boah	2019	Cross-sectional	0 - 59	Ghana	2636	■	15 - 19 years: reference group. 20 - 29: 1.34 (0.30 - 6.03) 30 - 39: 1.30 (0.27 - 6.38) 40 - 49: 2.51 (0.38 - 16.61) No significant p-values reported	↓
37	Mena-Melendez	2020	Cross-sectional	0 - 59	Latin America: Bolivia, Colombia, Guatemala and Peru	15827	■	Adjusted correlation co-efficient for age at first birth: 1.009 (0.821 – 1.240) SD: 0.106	↑
39	Poda	2017	Cross-sectional	0 - 59	Burkina Faso	6337	■	Univariate OR <20 years: reference group. 20 - 24: 0.61 (0.42 - 1.39) 25 - 29: 0.55 (0.38 - 1.29) 30 - 34: 0.54 (0.37 - 1.29) ≥35: 0.50 (0.34 - 0.72) P=<0.01 Multivariate OR <20 years: reference group. 20 - 24: 0.91 (0.61 – 1.36) 25 - 29: 0.88 (0.57 – 1.36) 30 - 34: 0.91 (0.56 – 1.46) ≥35: 0.88 (0.52 – 1.50)	↑
40	Mashal	2008	Cross-sectional	0 - 59	Afghanistan	2472	■	uOR Married ≥16: reference group Married <16: 1.18 (0.85 – 1.63)	↑
45	Samuel	2022	Cross-sectional	6 - 11	Ethiopia,	2036	⊕	uOR <25 years: 0.60 (0.42 – 0.85) p = 0.004 >25 years: reference group	↓
46	Akseer	2018	Cross-sectional	0 - 59	Afghanistan	22158	■	Crude coefficient: 40 – 49 years: reference group 30 – 39 years: -0.08 (-0.22 – 0.05) 20 – 29 years: -0.12 (-0.25 – 0) 15 – 19 years: -0.07 (-0.26 – 0.12)	↑ (compared to ref. grp only)
56	Beiersmann	2013	Cross-sectional	6 - 51	Burkina Faso	460	■	Mean (SD) WLZ <20 years: -1.68 (1.03) 20 – 24 years: -1.20 (1.15)	↑

								25 – 29 years: -1.04 (1.19) 30 – 34 years: -1.10 (1.25) >35 years: -1.34 (1.28) P 0.61	
57	Ickes	2015	Cross-sectional	0 - 23	Uganda	1897	■	aOR First birth >16 years: reference group 2006 First birth <16 years: 0.96 (0.62 – 1.49) 2011 First birth <16 years: 1.00 (0.57 – 1.75)	↔
63	Nguyen	2017	Cross-sectional	0 - 5	Bangladesh	2000	■	Adjusted mean WHZ ±SD ≤19 years: -0.68 ±1.60 >19 yrs) = -0.75 ±1.39	↔
64	Hien and Hoa	2009	Cross-sectional	6 - 36	Vietnam	383	⊕	aORs: >24 years: reference group ≤24 years: 2.16 (0.88 – 5.29)	↑↑
65	Hien and Kam	2008	Cross-sectional	0 - 59	Vietnam	650	⊕	aORs: 25 – 34 years: reference group ≤24 years: 1.20 (0.65 - 2.21)	↑↑
67	Olodu	2019	Cross-sectional	6 - 59	Nigeria	300	■	This study assessed the nutritional status of U5 children born to teenage mothers. 25.3% (72/285) of the children were wasted, but there is no comparison group provided.	N/A
70	Biswas	2021	Cross-sectional	0 - 59	South and South-East Asian countries	798961	■	aOR <-2SD Overweight/obese mother & wasted child: <20 years: reference group 20 – 29 years: 2.56 (2.11 - 3.1) p <0.005 30 – 39 years: 3.67 (3.03 - 4.46) p <0.005 40 – 49 years: 3.90 (3.14 - 4.85) p <0.005	↓↓
72	Kasaye	2019	Cross-sectional	0 – 59	Ethiopia	9494	⊕	Proportions were available, however this is a duplicated data source, so removed from the meta-analysis. Crude OR for mother's age: 15 – 24 years: 1.06 (0.90 – 1.24) >24 years: reference group P = 0.4872	↑↑
78	Das	2019	Cross-sectional	0 - 59	Bangladesh	5951	■	aOR <-2SD Overweight/obese mother & wasted child:	↔

								≤15 years: reference group 16 – 20 years: 1.17 (0.56 – 2.82) 21 – 25 years: 1.08 (0.39 – 3.08) ≥26 years: 0.93 (0.16 – 4.06)	
80	Mokwena	2022	Cross-sectional (hospital based study)	1 - 36	South Africa	94	■	All participants were malnourished. Disaggregated by maternal age the proportions are: 16 – 20 years: 19% 21 – 25 years: 35% 26 – 30 years: 26% 31 – 40 years: 14% NB: There are no 'healthy' controls included.	N/A
81	Gbadamosi	2017	Cross-sectional	1 - 12	South Africa	186	■	aOR: ≥19 years: reference group <19: 1.01 (0.232 - 4.403)	↔
86	Bekele	2021	Cross-sectional	0 - 59	Ethiopia	21514	⊕	Absolute contribution (AC) and percentage contribution (PC) show the adjusted contributions to inequalities for each predictor, for both 2000 and 2016: 2000 15 – 24 years: reference group 25 – 34 years: AC <0.0001 PC -0.71 35 – 44 years: AC <0.0001 PC -0.17 45 – 49 years: AC <0.0001 PC -1.71 2016 15 – 24 years: reference group 25 – 34 years: AC <0.0001 PC -0.03 35 – 44 years: AC <0.0001 PC -0.17 45 – 49 years: AC -0.001 PC 2.05	↔
88	Gewa and Yandell	2012	Cross-sectional	0 - 60	Kenya	1851	⊕	Mean maternal age of mothers first birth for children that were wasted versus children that were not wasted. 0 - 24-month-old children: Wasted: (n = 147) 19 (18.9 - 19.2) not wasted: (n = 1704) 18.2 (17.7 - 18.7) 25 - 60-month-old children: Wasted: (n = 95), 19.1 (18.3 - 19.9) not wasted: (n=1846) 19.1 (19.0 - 19.3)	N/A

Severe wasting									
27	Ahmed	2012	Cross-sectional	0 - 24	Bangladesh	8858	■	aOR <20 years: 0.96 (0.08 – 1.15) 20 – 30 years: reference group >30 years: 0.96 (0.77 – 1.19)	⇔
80	Mokwena	2022	Cross-sectional (hospital based study)	1 - 36 months	South Africa	94	■	All participants were malnourished. Proportion of those admitted with severe wasting in each maternal age group versus moderate wasting 16 – 20 years: 44.4% 21 – 25 years: 32.3% 26 – 30 years: 23.1% 31 – 40 years: 50% (NB n=2) NB: There are no 'healthy' controls included.	N/A
■ - Yes ⊕ - No aOR: adjusted odds ratio, uOR: unadjusted odds ratio p: p-value *Indication of outcome direction: ↑ = study indicates adolescent pregnancy is associated with increased childhood wasting, compared to adult pregnancy ⇔ = study reports no difference between the groups ↓ = study indicates adolescent pregnancy is associated with reduced childhood wasting compared to adult pregnancy									

b. Moderate and severe underweight

Ref	First Author	Year	Study Design	Infant Age categories (months)	Country	Study Size	Maternal age category <20 years	Key findings	Indication of childhood underweight outcomes*
Moderate underweight									
3	Schott	2017	Cohort	0 - 78	Ethiopia, India	253	■	This study was an intergenerational study that assessed the association between adolescent mothers' anthropometric measurements at aged 8, 12, 15 and 19 years, as well as maternal diet at the age of 15 years (and grandmother's characteristics) on their off-springs WAZ (HAZ and birthweight). Mother's age at birth predicted higher cWAZ between birth and 2013 ($\beta 0.46$ $p < 0.10$ and $\beta 0.53$ $p < 0.05$) There was no comparison with an adult cohort.	↑*
4	Friebert	2017	RCT (assessor blinded)	6 weeks and 12 weeks	Malawi	2284	■	Mean WAZ (SD) At 6 weeks: Mother <18 years: -1.1 +-1.2 Mother 18 - 20 years: -0.9 +-1.1 Mother >20 years: 0.8 +-1.0 P = <0.00 At 12 weeks: Mother <18 years: -0.8 +- 1.1 Mother 18 - 20 years: -0.8 +-1.0 Mother >20 years: -1.0 +-1.1 p <0.00 NB: All mothers were malnourished	↑* (At 6 weeks)
8	Fuada	2020	Cross-sectional	0 - 59	Indonesia	978	■	OR for Adolescent mother group ≤15 years: 1.311 (0.498 - 3.450) 15 – 19 years: reference group p 0.773 NB – no adult mother comparison	↑

13	Hiruy	2021	Cross-sectional	6 - 23	Ethiopia	8003	■	aOR provided for 2000, 2005, 2011, 2016. 15 – 19 years: reference group	↑*
								2000	
								20 – 29 years: 1.45 (0.98 – 2.14) 30 – 39 years: 1.79 (1.20 – 2.68)** 40 – 49 years: 1.75 (1.06 – 2.88)*	
								2005	
14	Ali	2019	Cross-sectional	0 - 59	Bangladesh	6468	⊕	20 – 29 years: 1.19 (0.59 – 2.38) 30 – 39 years: 1.32 (0.64 – 2.73) 40 – 49 years: 2.26 (0.91 – 5.61)	↓
								2011	
								20 – 29 years: 1.18 (0.81 – 1.73) 30 – 39 years: 1.28 (0.86 – 1.91) 40 – 49 years: 1.28 (0.76 – 2.15)	
								2016	
18	Kumar and Paswan	2021	Cross-sectional	0 - 59	India	11858	■	20 – 29 years: 0.54 (0.33 – 0.89)* 30 – 39 years: 0.67 (0.42 – 1.07) 40 – 49 years: 1.17 (0.65 – 2.21)	↔
								*p <0.05 **p <0.01	
								aOR: 15 – 24 years: reference group 25 - 49 years: 1.08 p 0.186	
								aOR provided for 2005-2006 and 2015 -2016 <20 years: reference group	
18	Kumar and Paswan	2021	Cross-sectional	0 - 59	India	11858	■	2005 - 2005	↔
								20 – 29 years: 0.93 (0.523 – 1.654) 30 – 49 years: 0.811 (0.443 – 1.485)	
								2015 - 2016	
								20 – 29 years: 1.021 (0.923 – 1.130) 30 – 49 years: 1.021 (0.916 – 1.139)	

								OLS (ordinary least squares) regression model (Standard error) for ‘teenage mother’ Adjusted for individual and household variables: -0.135** (0.053) Adjusted for individual, household and community variables: -0.122** (0.053) Adjusted for ‘instrumented variables’ (the presence of either an NGO or health post in the village): -0.113** (0.055)	
21	Linnemayr	2008	Cross-sectional	0 - 35	Senegal	4296	⊕		↑*
**significance at 5% level									
27	Ahmed	2012	Cross-sectional	0 - 24	Bangladesh	8858	■	aORs <-2SD: <20 years: 1.04 (0.86 - 1.26) 20 – 30 years: reference group >30 years: 1.31 (1.04 - 1.65)* *p <0.05	↑
30	Sobkoviak	2012	Cross-sectional	0 - 48	Liberia	2467	⊕	aOR: 15 – 24 years: reference group: 25 - 34: 0.65 (SE: 0.11) p <0.001 35 - 49: 0.41 (SE: 0.12) p <0.001	↑*
31	Hossain	2020	Cross-sectional	24 - 59	5 South Asian countries (India, Bangladesh, Pakistan, Maldives and Nepal)	146996	■	aOR: 20 - 24 years: reference group <20 years: Bangladesh: 1.1 (0.9 - 1.3) India: 1.1 (1.0 - 1.1) Maldives: 1.1 (0.8 - 1.5) Nepal: 0.9 (0.7 - 1.2) Pakistan: 1.0 (0.8 - 1.2) ≥25 years: Bangladesh: 0.8 (0.5 – 1.1) India: 0.8 (0.8 – 0.8) Maldives: 0.8 (0.5 - 1.3) Nepal: 0.8 (0.5 – 1.4) Pakistan: 0.9 (0.7 – 1.3)	↑
36	Boah	2019	Cross-sectional	0 - 59	Ghana	2636	■	aOR: 15 - 19 years: reference group 20 – 29 years: 0.50 (0.17 - 1.42)	↑

								30 – 39 years: 0.54 (0.16 - 1.76) 40 – 49 years: 0.76 (0.19 - 3.03)	
32	Masibo	2020	Cross-sectional	0 - 59	Kenya	7830	■	aOR Overweight/obese mother – underweight child (<- 2SD) <20 years: 0.43 (0.07 – 2.65) p 0.36 20 – 30 years: 0.55 (0.24 – 1.26) p 0.156 30 – 40 years: 1.08 (0.48 – 2.40) p 0.857 40+ years: reference group	⇓
39	Poda	2017	Cross-sectional	0 - 59	Burkina Faso	6337	■	Univariate and multivariate analysis: <20 years: reference group Univariate 20 – 24 years: 0.68 (0.49 - 1.25) 25 – 29 years: 0.57 (0.42 - 1.01) 30 – 34 years: 0.61 (0.45 - 0.83)* ≥35 years: 0.61 (0.45 - 0.84)* Multivariate 20 – 24 years: 0.77 (0.53 – 1.12) 25 – 29 years: 0.61 (0.41 – 1.18) 30 – 34 years: 0.61 (0.40 – 1.05) ≥35 years: 0.61 (0.38 – 0.97)* *p <0.05	⇑
40	Mashal	2008	Cross-sectional	0 - 59	Afghanistan	2472	■	uOR Married >16 years: reference group Married <16 years: 1.41 (1.12 - 1.78)	⇑*
46	Akseer	2018	Cross-sectional	0 - 59	Afghanistan	14000	■	Mother's age at delivery was disaggregated by: <20 years 20 – 29 years 30 – 39 years 40 – 49 years Using 30 – 39 years as the reference group it was found that mother's age was not significantly associated with underweight amongst the offspring. (non-significant ORs were not published)	⇔

								ORs for 2006 and 2011: First birth at >16 years old: reference group	
57	Ickes et al	2015	Cross-sectional	0 - 23	Uganda	1009	■	2006 First birth <16 years: 1.03 (0.72 - 1.47) 2011 First birth <16 years 1.39 (0.92 - 2.10)	↑
63	Nguyen	2017	Cross-sectional	0 - 5	Bangladesh	200	■	Adjusted mean WAZ $\pm$ SD: ≤ 19 years: -1.21 ± 1.10 >19 years: -1.08 ± 1.80 p <0.05	↑
64	Hien and Hoa	2009	Cross-sectional	6 - 36	Vietnam	383	⊕	aORs: ≤ 24 years: 1.12 (0.64 – 1.98) >24 years: reference group	↑
65	Hien and Kam	2008	Cross-sectional	0 - 59	Vietnam	650	⊕	aORs: ≤ 24 years: 0.93 (0.61 - 1.44) 25 – 34 years: reference group	↔
67	Olodu	2019	Cross-sectional	6 - 59	Nigeria	300	■	This study assessed the nutritional status of U5 children born to teenage mothers. 29.5% (84/285) of the children were underweight, but there is no comparison group provided.	N/A
68	Nakamori	2010	Cross-sectional	6 - 18	Vietnam	188	⊕	ORs: <25 years: reference group >25 years: 1.37 (0.60 - 3.14) p 0.451	↓
70	Biswas	2021	Cross-sectional	0 - 59	South and South-East Asian countries	798961	■	Overweight/obese mother & underweight child: <20 years: reference group 20 – 29 years: 2.56 (2.11- 3.09) p <0.005 30 – 39 years: 3.65 (3.01 - 4.43) p <0.005 40 – 49 years: 3.87 (3.12 - 4.80) p <0.005	↓
72	Kasaye	2019	Cross-sectional	0 – 59	Ethiopia	9494	⊕	Proportions were available, however this is a duplicated data source, so removed from the meta-analysis. Crude OR for mother's age: 15 – 24 years: 0.96 (0.85 – 1.08) >24 years: reference group P = 0.4798	↓
78	Das	2019	Cross-sectional	0 - 59	Bangladesh	5951	■	Overweight/obese mother & underweight child ≤ 15 years: reference group	↓

81	Gbadamosi	2017	Cross-sectional	1 - 12	South Africa	186	■	16 – 20 years: 1.47 (0.83 - 2.84) 21 – 25 years: 2.16 (1.08 - 4.48) p <0.05 ≥26 years: 2.17 (0.72 - 5.73) Not statistically significant aOR: <19 years:1.25 (0.062 - 25.271) ≥19 years: reference group	↑
86	Bekele	2021	Cross-sectional	0 - 59	Ethiopia	21514	⊕	Absolute contribution (AC) and percentage contribution (PC) show the adjusted contributions to inequalities for each predictor, for both 2000 and 2016: 2000 15 – 24 years: reference group 25 – 34 years: AC <0.0001 PC -0.10 35 – 44 years: AC <0.0001 PC -0.22 45 – 49 years: AC -0.011 PC 12.38 2016 15 – 24 years: reference group 25 – 34 years: AC <0.0001 PC -0.01 35 – 44 years: AC <0.0001 PC -0.03 45 – 49 years: AC -0.001 PC 0.77	↔
88	Gewa and Yandell	2012	Cross-sectional	0 - 60	Kenya	1851	⊕	Mean maternal age of mothers first birth for children that were wasted versus children that were not wasted. 0 - 24-month-old children: Underweight: (n = 230) 18.2 (17.7 0 18.6) not underweight: (n = 1621) 19.1 (18.9 – 19.3) 25 - 60-month-old children: Underweight: (n = 338), 18.5 (18.1 – 18.8) not wasted: (n=1604) 19.3 (19.1 – 19.4)	N/A
Severe underweight									
27	Ahmed et al	2012	Cross-sectional	0 – 24	Bangladesh	8858	■	aORs <-3SD: <20 years: 1.11 (0.95 – 1.29) 20 – 30 years: reference group >30 years: 1.21 (1.02 – 1.43)	↑

■ - Yes ⊕ - No

aOR: adjusted odds ratio, uOR: unadjusted odds ratio

p: p-value

*significance of <0.05

*Indication of outcome direction:

↑ = study indicates adolescent pregnancy is associated with **increased** childhood underweight, compared to adult pregnancy

↔ = study reports no difference between the groups

↓ = study indicates adolescent pregnancy is associated with **reduced** childhood underweight compared to adult pregnancy