

1 **Supplemental Information**

2

3 **Allelic strengths of encephalopathy-associated *UBA5* variants correlate between**
4 ***in vivo* and *in vitro* assays**

5

6 Xueyang Pan^{1,2}, Albert N. Alvarez³, Mengqi Ma^{1,2}, Shenzhao Lu^{1,2}, Michael W. Crawford³, Lauren C.
7 Briere⁴, Oguz Kanca^{1,2}, Shinya Yamamoto^{1,2,5}, David A. Sweetser^{4,6}, Jenny L. Wilson⁷, Ruth J.
8 Napier^{3,8,9}, Jonathan N. Pruneda^{3,#}, Hugo J. Bellen^{1,2,5,#}

9

10 1 Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX 77030,
11 USA

12 2 Jan & Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston, TX 77030,
13 USA

14 3 Department of Molecular Microbiology & Immunology, Oregon Health & Science University,
15 Portland, OR 97239, USA

16 4 Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA 02114, USA

17 5 Department of Neuroscience, Baylor College of Medicine, Houston, TX 77030, USA

18 6 Division of Medical Genetics & Metabolism, Massachusetts General Hospital for Children, Boston,
19 MA 02114, USA

20 7 Division of Pediatric Neurology, Department of Pediatrics, Oregon Health & Science University,
21 Portland, OR 97239, USA

22 8 VA Portland Health Care System, Portland, OR 97239, USA

23 9 Division of Arthritis & Rheumatic Diseases, Oregon Health & Science University, Portland, OR
24 97239, USA

25

26 # Correspondence: hbellen@bcm.edu, pruneda@ohsu.edu

27

1 **Case report**

2 The reported proband is a boy with axial hypotonia, generalized dystonia, lower extremity spasticity,
3 global developmental delay, esotropia and failure to thrive. An electroencephalogram (EEG) showed
4 multifocal epileptiform discharges in drowsiness and sleep. However, he has not had seizures. Two
5 magnetic resonance imaging (MRI) studies at different ages were read as normal, although on review
6 show a slightly thin corpus callosum, posterior periventricular white matter T2 hyperintensity resulting
7 in increased conspicuity of the subcortical U-fibers and widening of the Sylvian fissures (Figure S1). He
8 is not microcephalic. For more information about the clinical record of the proband please contact the
9 corresponding author.

10 On trio exome sequencing, the proband was found to have compound heterozygous variants in *UBA5*,
11 NM_024818.6:c.169A>G (p.Met57Val) and c.935A>T (p.Gln312Leu). The two variants are in *trans*
12 phase. The p.Met57Val variant has been reported in one individual with DEE44 (Colin *et al*, 2016). The
13 p.Gln312Leu variant has not been previously reported. Neither variant is reported in the gnomAD
14 database v.2.1.1 (<https://gnomad.broadinstitute.org>). Both variants are predicted to be damaging or
15 probably damaging by multiple pathogenicity prediction tools (Table S3).




Figure S1. Brain magnetic resonance imaging (MRI) images

- (A) Axial T2 image showing periventricular T2 hyperintensity (blue arrow) resulting in prominence of the subcortical U-fibers (white arrows).
- (B) Sagittal flair showing mild thinning of the corpus callosum (red arrow).
- (C) Axial flair image demonstrating widening of the sylvian fissures (white asterisks).

Table S1. Summary of genotypes of the reported cases

References	Family	Allele #1	Allele #2
Colin, et al., 2016 (Colin et al., 2016)	A	p.Ala371Thr (IA [†])	p.Gln302*
	B	p.Ala371Thr (IA)	p.Lys324Asnfs*14
	C	p.Asp389Tyr (IA)	p.Val260Met (II)
	D	p.Met57Val (II)	p.Gly168Glu (III)
Muona, et al., 2016 (Muona et al, 2016)	A	p.Ala371Thr (IA)	p.Arg55His (III)
	B	p.Ala371Thr (IA)	p.Tyr285*
	C, E	p.Ala371Thr (IA)	p.Arg188*
	D	p.Ala371Thr (IA)	p.Arg61*
Arnadottir, et al., 2017 (Arnadottir et al, 2017)		p.Ala371Thr (IA)	p.Ala288= (splicing variant)
Daida, et al., 2018 (Daida et al, 2018)		p.Tyr72Cys (IB)	Deletion
Mignon-Ravix, et al., 2018 (Mignon-Ravix et al, 2018)		p.Tyr53Phe (II) ^{††}	p.Tyr53Phe (II)
Low, et al., 2019 (Low et al, 2019)		p.Asp389Gly (IA)	Deletion
Briere, et al., 2021 (Briere et al, 2021)	A	p.Ala371Thr (IA)	p.Cys303Arg (III)
	B	p.Ala371Thr (IA)	p.Arg188*
	C	p.Ala371Thr (IA)	p.Leu254Pro (III)
	D	p.Ala371Thr (IA)	p.Cys303Arg (III)
This study		p.Met57Val (II)	p.Gln312Leu (IB)

[†] The variant classification using fly phenotypic assays (results shown in Figure 3)

1 †† Consanguineous family

2

3 **Table S2. Clinical features of individuals with *UBA5*-associated DEE44**

4 (See separate Excel table)

5

6 **Table S3. Bioinformatic predictions of the pathogenicity of reported *UBA5* variants**

	Variant 1	Variant 2
Genomic position (GRCh38)	3:132665830A>G	3:132394214A>T
Amino acid change	p.Met57Val	p.Gln312Leu
Allele frequency in gnomAD	Absent	Absent
CADD score	25.3	29.1
SIFT	Damaging	Damaging
PolyPhen2	1.000 (probably damaging)	0.999 (probably damaging)
MutationTaster	Disease causing	Disease causing
PROVEAN	-3.18 (deleterious)	-6.59 (deleterious)

7

1 **References**

- 2 Arnadottir GA, Jensson BO, Marelsson SE, Sulem G, Oddsson A, Kristjansson RP, Benonisdottir S,
3 Gudjonsson SA, Masson G, Thorisson GA *et al* (2017) Compound heterozygous mutations in UBA5
4 causing early-onset epileptic encephalopathy in two sisters. *BMC Med Genet* 18: 103
- 5 Briere LC, Walker MA, High FA, Cooper C, Rogers CA, Callahan CJ, Ishimura R, Ichimura Y, Caruso
6 PA, Sharma N *et al* (2021) A description of novel variants and review of phenotypic spectrum in
7 UBA5-related early epileptic encephalopathy. *Cold Spring Harb Mol Case Stud* 7
- 8 Colin E, Daniel J, Ziegler A, Wakim J, Scrivo A, Haack TB, Khiati S, Denomme AS, Amati-Bonneau
9 P, Charif M *et al* (2016) Biallelic Variants in UBA5 Reveal that Disruption of the UFM1 Cascade Can
10 Result in Early-Onset Encephalopathy. *Am J Hum Genet* 99: 695-703
- 11 Daida A, Hamano SI, Ikemoto S, Matsuura R, Nakashima M, Matsumoto N, Kato M (2018) Biallelic
12 loss-of-function UBA5 mutations in a patient with intractable West syndrome and profound failure to
13 thrive. *Epileptic Disord* 20: 313-318
- 14 Low KJ, Baptista J, Babiker M, Caswell R, King C, Ellard S, Scurr I (2019) Hemizygous UBA5
15 missense mutation unmasks recessive disorder in a patient with infantile-onset encephalopathy,
16 acquired microcephaly, small cerebellum, movement disorder and severe neurodevelopmental delay.
17 *Eur J Med Genet* 62: 97-102
- 18 Mignon-Ravix C, Milh M, Kaiser CS, Daniel J, Riccardi F, Cacciagli P, Nagara M, Busa T, Liebau E,
19 Villard L (2018) Abnormal function of the UBA5 protein in a case of early developmental and epileptic
20 encephalopathy with suppression-burst. *Hum Mutat* 39: 934-938
- 21 Muona M, Ishimura R, Laari A, Ichimura Y, Linnankivi T, Keski-Filppula R, Herva R, Rantala H,
22 Paetau A, Poyhonen M *et al* (2016) Biallelic Variants in UBA5 Link Dysfunctional UFM1 Ubiquitin-like
23 Modifier Pathway to Severe Infantile-Onset Encephalopathy. *Am J Hum Genet* 99: 683-694
- 24