

Supplementary Material

Supplementary table 1: Proportion of individuals presenting at the GP practice over the total number of brochures distributed by study arm

Supplementary Table 2. Sensitivity analysis of Odds ratios (95% CI) from exploratory logistic regression results comparing SOC, healthy lifestyle brochure and care recipient voucher brochure arms

Supplementary Table 3: Effect of intervention on odds of presenting at the GP practice with sociodemographics

Supplementary Table 1: Proportion of individuals presenting at the GP practice over the total number of brochures distributed by study arm

		Standard of Care		Healthy Lifestyle Brochure		Recipient Care Voucher	
		Brochures distributed	Individuals presenting	Brochures distributed	Individuals presenting	Brochures distributed	Individuals presenting
		n	n (%)	n	n (%)	n	n (%)
Total		3802	137 (3.6)	3840	153 (4.0)	3829	158 (4.1)
Gender	Male	2246	69 (3.1)	2293	76 (3.3)	2235	82 (3.7)
	Female	1532	68 (4.4)	1512	77 (5.1)	1557	76 (4.9)
Age group	18-24 years	654	24 (3.7)	612	17 (2.8)	624	24 (3.8)
	25-34 years	1901	67 (3.5)	1911	60 (3.1)	1962	88 (4.5)
	35-44 years	1015	37 (3.6)	1060	48 (4.5)	991	35 (3.5)
	≥ 45 years	216	9 (4.2)	228	28 (12.3)	225	11 (4.9)
HIV test outcomes	HIV negative	-	130 (3.4)	-	146 (3.8)	-	156 (4.1)
	PLHIV	-	4/137 (0.1)	-	5/153 (0.1)	-	1/158 (0.0)
	Unknown HIV outcomes	-	3 (0.1)	-	2 (0.1)	-	1 (0.0)

Supplementary Table 2: Sensitivity analysis of Odds ratios (95% CI) from exploratory logistic regression results comparing SOC, healthy lifestyle brochure and care recipient voucher brochure arms*

		No. of participants	Presenting at the GP practice (%)	UOR (95% CI)	P-value	AOR (95% CI)**	P-value
Study arm	SOC	3603	137 (3.8%)	1 [Ref]		1 [Ref]	
	HLS	3620	153 (4.2%)	1.11 (0.88-1.40)	.394	0.92 (0.70-1.21)	.554
	RCV	3608	158 (4.4%)	1.12 (0.89-1.41)	.347	0.99 (0.76-1.23)	.924
Gender	Male	6406	227 (3.5%)	1 [Ref]		1 [Ref]	
	Female	4298	221 (5.1%)	1.11 (0.89-1.38)	.371	0.97 (0.77-1.21)	.762
Age group	18-24	1744	65 (3.7%)	1 [Ref]		1 [Ref]	
	25-34	5438	215 (4.0%)	0.76 (0.56-1.01)	.061	0.95 (0.70-1.28)	.728
	35-44	2927	120 (4.1%)	0.76 (0.55-1.05)	.101	0.89 (0.64-1.63)	.509
	≥ 45	612	48 (7.8%)	1.31 (0.84-2.04)	.230	1.04 (0.66-1.63)	.865
GP group	GP group with limited visibility	6630	44 (0.7)	1 [Ref]		1 [Ref]	
	GP group with high visibility	4201	404 (9.6)	6.49 (5.15-8.18)	.000	4.92 (3.82-6.34)	.000
Language	English	7756	-	1 [Ref]		1 [Ref]	
	IsiZulu	2466	-	1.50 (1.21-1.85)	.000	1.06 (0.82-1.37)	.665
	SeSotho	609	-	1.57 (1.09-2.26)	.016	0.63 (0.33-1.20)	.163

* data excludes all duplicates

**Adjusted for study arm category, gender, age group, GP group and language

Supplementary Table 3. Effect of intervention on odds of presenting at the GP practice with sociodemographics*

		No. of participants	Presenting at the GP practice (%)	UOR (95% CI)	P-value	AOR 95% (CI)	P-value
Study arm	SOC	3802	137 (3.6%)	1 [Ref]		1 [Ref]	
	HLS	3840	153 (4.0%)	1.11 (0.88-1.40)	.384	1.02 (0.79-1.32)	.873
	RCV	3829	158 (4.1%)	1.15 (0.91-1.45)	.236	1.08 (0.84-1.40)	.559
Gender	Male	6774	227 (3.4%)	1 [Ref]		1 [Ref]	
	Female	4601	221 (4.8%)	1.11 (0.89-1.38)	.371	1.02 (0.82-1.26)	.852
Age group	18-24	1890	65 (3.4%)	1 [Ref]		1 [Ref]	
	25-34	5774	215 (3.7%)	0.82 (0.62-1.08)	.155	1.03 (0.77-1.37)	.856
	35-44	3066	120 (3.9%)	0.77 (0.56-1.06)	.108	0.90 (0.65-1.24)	.505
	≥ 45	669	48 (7.2%)	1.27 (0.83-1.94)	.274	0.98 (0.63-1.51)	.918
GP group	GP group with limited visibility	6885	44 (0.6%)	1 [Ref]		1 [Ref]	
	GP group with high visibility	4586	404 (8.8%)	6.13 (4.86-7.72)	.000	5.30 (4.14-6.79)	.000
Language**	English	8165	-	1 [Ref]		1 [Ref]	
	IsiZulu	2656	-	1.46 (1.18-1.80)	.000	0.98 (0.77-1.26)	.895
	SeSotho	650	-	1.49 (1.03-2.16)	.035	1.07 (0.67-1.72)	.762

*Adjusted for gender, age group, clinic group, language (using combined GP groups and combined intervention arms)

**Language was not recorded at presentation at the GP practice