

Supplementary Table 1. Factors associated with HIV test positivity. Adjusted odds ratios from logistic regression analyses by sex.

		Men (n=1471) ^a			Women (n=2422) ^a		
		aOR ^a	95% CI	P-value	aOR ^a	95% CI	P-value
Sociodemographic variables							
Age group (in years)^b							
	15-19	Ref.			Ref.		
	20-24	3.00	0.36-24.68	0.307	1.52	0.81-2.84	0.187
	25-49	13.61	1.72-107.66	0.013	1.59	0.85-2.98	0.147
	≥50	4.66	0.55-39.35	0.158	0.43	0.18-1.04	0.061
Health facility^b							
	Manhiça district hospital	Ref.			Ref.		
	Xinavane rural hospital	0.82	0.52-1.30	0.404	0.41	0.27-0.61	<0.001
	Palmeira health unit	0.64	0.33-1.25	0.191	0.52	0.32-0.83	0.007
	Maragra health unit	0.51	0.26-1.02	0.057	0.27	0.16-0.48	<0.001
Occupation^b							
	Farmer	Ref.			Ref.		
	Industry/Miner	3.81	1.55-9.34	0.003	7.66	0.62-94.05	0.112
	Own or employer's business	0.98	0.57-1.67	0.935	1.75	0.98-3.11	0.057
	Street vendor	0.42	0.52-3.47	0.424	1.35	0.64-2.84	0.425
	Student	0.29	0.04-2.37	0.247	0.69	0.34-1.39	0.298
	Construction-related work	1.03	0.57-1.84	0.928	5.10	0.91-28.47	0.063
	State worker	0.52	0.23-1.18	0.117	0.42	0.10-1.82	0.247
	Unemployed	0.91	0.54-1.55	0.729	1.29	0.86-1.92	0.217
	Other	0.85	0.34-2.10	0.720	1.45	0.61-3.46	0.402

Risk factors							
HIV-positive partner							
	No	Ref.			Ref.		
	Yes	6.48	3.75-11.18	<0.001	4.12	2.52-6.74	<0.001
Don't know (The partner never told him/her) ^b		2.69	1.65-4.38	<0.001	2.80	2.00-3.92	<0.001
More than one sexual partner in the past year							
	No	Ref.			Ref.		
	Yes	1.65	1.04-2.64	0.034	2.21	1.10-4.43	0.026
Condom use							
	Always	Ref.			Ref.		
	Frequently ^c	1.81	0.78-4.21	0.171	1.27	0.55-2.95	0.577
	Occasionally ^c	1.46	0.75-2.83	0.261	1.86	0.89-3.90	0.099
	Never	1.77	0.92-3.42	0.089	2.30	1.11-4.79	0.026
Visited a healer in the previous 6 months ^b							
	No	Ref.			Ref.		
	Yes	2.10	0.95-4.63	0.066	1.59	0.76-3.31	0.217
Other factors							
Health department ^b							
	Triage	Ref.			Ref.		
	Emergency	1.50	0.76-2.94	0.239	2.23	1.23-4.05	0.008
Pregnant or partner of pregnant woman ^b							
	No	Ref.			Ref.		

	Yes	1.50	0.82-2.73	0.187	1.52	0.46-5.05	0.492
	Don't know	2.82	0.50-15.90	0.239	2.23	1.01-4.94	0.048
Signs and symptoms							
Skin or oral mucosa lesions							
	No	Ref.			Ref.		
	Yes	3.54	1.50-8.35	0.004	4.11	1.78-9.51	0.001
Cough for over 3 weeks							
	No	Ref.			Ref.		
	Yes	1.29	0.44- 3.79	0.641	2.04	0.97-4.29	0.061
Night sweats^{b, d}							
	No	Ref.					
	Yes	8.25	1.15-59.47	0.036			
Constitutional syndrome							
(asthenia anorexia weight loss)							
	No	Ref.			Ref.		
	Yes	2.07	0.66-6.45	0.211	2.94	0.44-19.80	0.268

^a Client participants tested for HIV in both observation phases. 14 individuals with an undetermined HIV test result were excluded. For the multivariable analysis, 1471 men and 2422 were included, respectively.

^b Sociodemographic characteristics, risk factors or signs and symptoms not included in the national targeted PITC algorithm of Mozambique.

^c Occasionally: less than half of the times; Frequently: more than half of the times

^d The aOR among women could not be estimated because there was not any woman included in the multivariable analysis who presented with night sweats.

Abbreviations: aOR: adjusted odds ratio, CI: confidence interval, PITC: provider-initiated testing and counselling, Ref.: reference category