




Supplementary Figures




Supplementary figure 1. Number of multi-day cycle groups determined by k-means clustering. The optimal number of clusters (k) was identified by minimising the **a.** Sum of squared error at the elbow point (k = 3), and maximising the **b.** Silhouette coefficient.




Supplementary figure 2. Age subgroups. Distribution of the maximum SI values in each cycle group for child (<13 years), adolescent (13 - 17 years) and adult (≥ 18 years) cohorts, **a.** circadian (0.5 - 1.5 days) **b.** about-weekly (2 - 12 days) **c.** about-fortnightly (13 - 22 days) and **d.** about-monthly (23 - 32 days)



Supplementary figure 3. Gender subgroups. Distribution of the maximum SI values in each cycle group for paediatric female, paediatric male, adult female and adult male subgroups, **a.** circadian (0.5 - 1.5 days) **b.** about-weekly (2 - 12 days) **c.** about-fortnightly (13 - 22 days) and **d.** about-monthly (23 - 32 days)



Supplementary figure 4. Epilepsy type. Distribution of the maximum SI values in each cycle group for paediatric focal, paediatric generalised, adult focal and adult generalised subgroups, **a.** circadian (0.5 - 1.5 days) **b.** about-weekly (2 - 12 days) **c.** about-fortnightly (13 - 22 days) and **d.** about-monthly (23 - 32 days)



Supplementary figure 5. Medication subgroups. Distribution of the maximum SI values in each cycle group for paediatric with ASM, paediatric with no ASM, adult with ASM and adult with no ASM subgroups, **a.** circadian (0.5 - 1.5 days) **b.** about-weekly (2 - 12 days) **c.** about-fortnightly (13 - 22 days) and **d.** about-monthly (23 - 32 days)

Supplementary Tables

Cycle group	Comparison (n)		Wilcoxon Rank Sum		Kolmogorov-Smirnov	
			p-value	Adjusted p-value	p-value	Adjusted p-value
Circadian	Child (46)	Adolescent (25)	0.531	0.974	0.957	0.974
	Child (46)	Adult (254)	0.046	0.552	0.023	0.552
	Adolescent (25)	Adult (254)	0.533	0.974	0.510	0.974
About-weekly	Child (46)	Adolescent (25)	0.876	0.974	0.948	0.974
	Child (46)	Adult (254)	0.954	0.974	0.907	0.974
	Adolescent (25)	Adult (254)	0.789	0.974	0.931	0.974
About-fortnightly	Child (35)	Adolescent (21)	0.548	0.974	0.667	0.974
	Child (35)	Adult (203)	0.490	0.974	0.477	0.974
	Adolescent (21)	Adult (203)	0.853	0.974	0.974	0.974
About-monthly	Child (28)	Adolescent (19)	0.165	0.712	0.106	0.712
	Child (28)	Adult (173)	0.878	0.974	0.736	0.974
	Adolescent (19)	Adult (173)	0.178	0.712	0.139	0.712

Supplementary table 1. Age subgroups. Statistical tests with adjusted p-values (computed using the Benjamini-Hochberg method) comparing the distribution of the maximum SI values in each cycle group for child (<13 years), adolescent (13 - 17 years) and adult (≥18 years) cohorts.

Cycle group	Comparison (n)		Wilcoxon Rank Sum		Kolmogorov-Smirnov	
			<i>p</i> -value	Adjusted <i>p</i> -value	<i>p</i> -value	Adjusted <i>p</i> -value
Circadian	Paed female (37)	Paed male (25)	0.514	0.847	0.508	0.847
	Paed female (37)	Adult female (151)	0.061	0.448	0.034	0.416
	Paed male (25)	Adult male (74)	0.425	0.847	0.794	0.847
	Adult female (151)	Adult male (74)	0.790	0.847	0.772	0.847
About-weekly	Paed female (37)	Paed male (25)	0.305	0.751	0.475	0.847
	Paed female (37)	Adult female (151)	0.720	0.847	0.666	0.847
	Paed male (25)	Adult male (74)	0.070	0.448	0.120	0.549
	Adult female (151)	Adult male (74)	0.179	0.643	0.116	0.589
About-fortnightly	Paed female (28)	Paed male (21)	0.210	0.643	0.241	0.643
	Paed female (28)	Adult female (124)	0.628	0.847	0.863	0.863
	Paed male (21)	Adult male (58)	0.610	0.847	0.770	0.847
	Adult female (124)	Adult male (58)	0.039	0.416	0.015	0.416
About-monthly	Paed female (24)	Paed male (17)	0.427	0.847	0.576	0.847
	Paed female (24)	Adult female (106)	0.214	0.643	0.234	0.643
	Paed male (17)	Adult male (48)	0.743	0.847	0.766	0.847
	Adult female (106)	Adult male (48)	0.861	0.863	0.622	0.847

Supplementary table 2. Gender subgroups. Statistical tests with adjusted *p*-values (computed using the Benjamini-Hochberg method) comparing the distribution of maximum SI values for paediatric female, paediatric male, adult female and adult male subgroups.

Cycle group	Comparison (n)		Wilcoxon Rank Sum		Kolmogorov-Smirnov	
			<i>p</i> -value	Adjusted <i>p</i> -value	<i>p</i> -value	Adjusted <i>p</i> -value
Circadian	Paed focal (19)	Paed generalised (32)	0.360	0.683	0.496	0.794
	Paed focal (19)	Adult focal (141)	0.307	0.683	0.260	0.647
	Paed generalised (32)	Adult generalised (28)	0.025	0.371	0.033	0.371
	Adult focal (141)	Adult generalised (28)	0.675	0.864	0.822	0.974
About-weekly	Paed focal (19)	Paed generalised (32)	0.185	0.556	0.058	0.371
	Paed focal (19)	Adult focal (141)	0.325	0.682	0.384	0.682
	Paed generalised (32)	Adult generalised (28)	0.767	0.944	0.628	0.864
	Adult focal (141)	Adult generalised (28)	0.648	0.864	0.459	0.773
About-fortnightly	Paed focal (16)	Paed generalised (22)	0.098	0.523	0.132	0.528
	Paed focal (16)	Adult focal (119)	0.124	0.528	0.053	0.371
	Paed generalised (22)	Adult generalised (19)	0.583	0.864	0.857	0.979
	Adult focal (119)	Adult generalised (19)	0.988	0.988	0.942	0.980
About-monthly	Paed focal (12)	Paed generalised (18)	0.038	0.371	0.191	0.556
	Paed focal (12)	Adult focal (103)	0.188	0.556	0.263	0.647
	Paed generalised (18)	Adult generalised (17)	0.373	0.683	0.603	0.864
	Adult focal (103)	Adult generalised (17)	0.949	0.980	0.937	0.980

Supplementary table 3. Seizure subgroups. Statistical tests with adjusted *p*-values (computed using the Benjamini-Hochberg method) comparing the distribution of maximum SI values for paediatric focal, paediatric generalised, adult focal and adult generalised subgroups.

Cycle group	Comparison (n)		Wilcoxon Rank Sum		Kolmogorov-Smirnov	
			<i>p</i> -value	Adjusted <i>p</i> -value	<i>p</i> -value	Adjusted <i>p</i> -value
Circadian	Paed with ASM (44)	Paed with no ASM (27)	0.887	0.957	0.126	0.632
	Paed with ASM (44)	Adult with ASM (184)	0.075	0.632	0.183	0.651
	Paed with no ASM (27)	Adult with no ASM (70)	0.778	0.957	0.139	0.632
	Adult with ASM (184)	Adult with no ASM (70)	0.124	0.632	0.269	0.783
About-weekly	Paed with ASM (44)	Paed with no ASM (27)	0.776	0.957	0.922	0.957
	Paed with ASM (44)	Adult with ASM (184)	0.867	0.957	0.957	0.957
	Paed with no ASM (27)	Adult with no ASM (70)	0.760	0.957	0.851	0.957
	Adult with ASM (184)	Adult with no ASM (70)	0.361	0.825	0.298	0.795
About-fortnightly	Paed with ASM (37)	Paed with no ASM (19)	0.117	0.632	0.158	0.632
	Paed with ASM (37)	Adult with ASM (153)	0.479	0.957	0.616	0.957
	Paed with no ASM (19)	Adult with no ASM (50)	0.936	0.957	0.815	0.957
	Adult with ASM (153)	Adult with no ASM (50)	0.098	0.632	0.101	0.632
About-monthly	Paed with ASM (32)	Paed with no ASM (15)	0.819	0.957	0.872	0.957
	Paed with ASM (32)	Adult with ASM (132)	0.348	0.825	0.572	0.957
	Paed with no ASM (15)	Adult with no ASM (41)	0.677	0.957	0.610	0.957
	Adult with ASM (132)	Adult with no ASM (41)	0.264	0.783	0.482	0.957

Supplementary table 4. Medication subgroups. Statistical tests with adjusted *p*-values (computed using the Benjamini-Hochberg method) comparing the distribution of maximum SI values for paediatric with ASM, paediatric with no ASM, adult with asm and adult with no ASM subgroups.