

SUPPLEMENTARY MATERIALS

to:

Lamsma et al. (2024). Structural brain abnormalities and aggressive behaviour in schizophrenia: Mega-analysis of data from 2095 patients and 2861 healthy controls via the ENIGMA consortium

Funding information for participating ENIGMA sites	3
Table S1. Items used to measure aggressive behaviour and potential mediators	6
Table S2. Global imaging-derived phenotypes used in the normative models as predictors of local imaging-derived phenotypes	6
Table S3. Distribution of aggression scores in individuals with schizophrenia who underwent T1-weighted magnetic resonance imaging	7
Table S4. Distribution of aggression scores in individuals with schizophrenia who underwent diffusion-weighted magnetic resonance imaging.....	7
Table S5. Deviations from normative values of cortical thickness (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia (N = 2012)	8
Table S6. Deviations from normative values of surface area (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia (N = 2009).....	8
Table S7. Deviations from normative values of fractional anisotropy (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia (N = 733).....	9
Table S8. Deviations from normative values of mean diffusivity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia (N = 607).....	9
Table S9. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for current dosage of antipsychotic medication (N = 1499).....	10
Table S10. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for current dosage of antipsychotic medication (N = 549)	10
Table S11. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for duration of illness (N = 1680).....	11
Table S12. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for duration of illness (N = 454).....	11
Table S13. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for lifetime substance use disorder (N = 778).....	12

Table S14. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for lifetime substance use disorder (N = 217).....	12
Table S15. Deviations from normative values of grey matter volume (A) and their associations with scores on PANSS item P7 (B) in individuals with schizophrenia (N = 1569).....	13
Table S16. Deviations from normative values of white matter microstructural integrity (A) and their associations with scores on PANSS item P7 (B) in individuals with schizophrenia (N = 504).....	13

Funding information for participating ENIGMA sites

COBRE

The COBRE study was supported by the National Institutes of Health (NIH) (grants R01EB006841 and P20GM103472) and National Science Foundation (NSF) (grant 1539067). JAT and VDC are supported by NIH grants 5R01MH094524 and 5R01MH121246.

ESO

The ESO study was supported by the Ministry of Health of the Czech Republic (grants NU21-08-00432 and NU22-04-00143).

FIDMAG

The FIDMAG study was supported by a Miguel Servet Research Contract (MS14/00041), the Plan Nacional de I+D+i 2013–2016 (Research Project PI14/00292), a Juan de la Cierva-formación contract (FJCI-2015-25278), the Instituto de Salud Carlos III-Subdirección General de Evaluación y Fomento de la Investigación, the European Regional Development Fund and the Centro de Investigación Biomédica en Red de Salud Mental.

FOR2107

The FOR2107 study was supported by grants from the German Research Foundation to TK (KI 588/14-1 and KI 588/14-2), UD (DA 1151/5-1, DA 1151/5-2, DA1151/9-1, DA1151/10-1 and DA1151/11-1) and IN (NE2254/1-2, NE2254/2-1, NE2254/3-1 and NE2254/4-1). UD is also supported by the Interdisciplinary Centre for Clinical Research of the University of Münster (grant Dan3/022/22).

GAP

The GAP study was supported by the NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London.

IGP

The IGP study was supported by grants (APP630471 and APP1081603) and a Career Development Award (APP1061875) from the Australian National Health and Medical Research Council and a grant (CE110001021) from the Australian Research Council Centre of Excellence in Cognition and its Disorders to MJG.

IMH

This work was supported by grants from the National Healthcare Group (SIG/05004 and SIG/05028) and the Singapore Bioimaging Consortium (RP C009/2006) to KS.

JBNU

This work was supported by grants from the Korean Mental Health Technology R&D Project (HL19C0015) and the Korea Health Industry Development Institute (HR18C0016) to Y-CC.

MCIC

The MCIC study was supported by the NIH (grants P41RR14075 and R01EB005846 to VDC), Department of Energy (DE-FG02-99ER62764), Mind Research Network, Morphometry (1U24, RR021382A) and Function (U24RR021992-01, NIH.NCRR MO1 RR025758-01 and NIMH 1RC1MH089257 to VDC) testbeds of the Biomedical Informatics Research Network, German Research Foundation (research fellowship to SE), National Alliance for Research on Schizophrenia & Depression (Young Investigator Award to SE) and Else-Kröner Fresenius Foundation (grant 2019-A118 to DA).

Montreal

This work was supported through the Eli Lilly Chair of Schizophrenia at the University of Montreal, held by SP.

MPRC

The MPRC study was supported by NIH grants (U01MH108148, 2R01EB015611, R01MH112180, R01DA027680, R01MH085646, P50MH103222 and T32MH067533), a State of Maryland contract (M00B6400091) and an NSF grant (1620457).

NU

The NU study was supported by the NIH (grants P50 MH071616, R01 EB020062, R01 MH056584, R01 MH084803 and U01 MH097435 to LW and grant T32 NS047987 to DC) and NSF (grants 1636893 and NSF 1734853 to LW).

Olin

The Olin study was supported by NIH grant R01MH077945.

RomeSL

This work was supported by the Italian Ministry of Health (Ricerca Corrente 2023).

RSCZ

The RSCZ study was supported by the Russian Foundation for Basic Research (grant 20-013-00748).

SNUH

This work was supported by the National Research Foundation of Korea and Korea Brain Research Institute (grants 2019R1C1C1002457, 2020M3E5D9079910, RS-2023-00266120 and 21-BR-03-01).

UCISZ

The UCISZ study was supported by the National Institutes of Mental Health (grant R21MH097196 to TGMvE). Data were processed with the high-performance computing cluster at the University of California, Irvine, supported by Joseph Farran, Harry Mangalam, Adam Brenner, the National Center for Research Resources and National Center for Advancing Translational Sciences (grant UL1 TR000153).

UMCU

This work was supported by the Geestkracht programme of the Dutch Health Research Council (grant 10-000-1001), University Medical Center Utrecht and participating pharmaceutical companies (Lundbeck, AstraZeneca, Eli Lilly, Janssen Cilag) and mental healthcare institutions (Altrecht, GGZ Centraal and Delta).

UNINA

FI, AdB, MC and AB were supported by NextGenerationEU, the Italian Ministry of Education, University and Research, National Recovery and Resilience Plan and project MNESYS (PE0000006) – A Multiscale integrated approach to the study of the nervous system in health and disease (DN. 1553 11.10.2022).

Zürich

This work was supported by the Swiss National Science Foundation.

Table S1. Items used to measure aggressive behaviour and potential mediators.

Variable	Item
Aggressive behaviour	PANSS item P7 ‘hostility’
	SAPS item B3 ‘aggressive and agitated behaviour’
	BPRS item 10 ‘hostility’
Delusions	PANSS item P1 ‘delusions’
	SAPS item D13 ‘global rating of severity of delusions’
Hallucinations	PANSS item P3 ‘hallucinatory behaviour’
	SAPS item H7 ‘global rating of severity of hallucinations’
	BPRS item 12 ‘hallucinatory behaviour’
Disorganised thinking	PANSS item P2 ‘conceptual disorganisation’
	SAPS item P9 ‘global rating of positive formal thought disorder’
	BPRS item 4 ‘conceptual disorganisation’
Poor impulse control	PANSS item P14 ‘poor impulse control’
Lack of illness insight	PANSS item G12 ‘lack of judgment and insight’

PANSS, Positive and Negative Syndrome Scale; SAPS, Scale for the Assessment of Positive Symptoms; BPRS, Brief Psychiatric Rating Scale.

Table S2. Global imaging-derived phenotypes used in the normative models as predictors of local imaging-derived phenotypes.

Class of local IDPs	Global IDP
GM volume	Total cortical volume (cortical regions)
	Total GM volume (subcortical regions)
Cortical thickness	Average CT across the entire cortex
Surface area	Total SA of the cortex
WM microstructural integrity	Average of deviation scores for global FA and global MD
Fractional anisotropy	Average FA across the entire WM skeleton
Mean diffusivity	Average MD across the entire WM skeleton

IDP, imaging-derived phenotype; GM, grey matter; CT, cortical thickness; SA, surface area; WM, white matter; FA, fractional anisotropy; MD, mean diffusivity.

Table S3. Distribution of aggression scores in individuals with schizophrenia who underwent T₁-weighted magnetic resonance imaging.

Score	Item			
	PANSS P7 (<i>n</i> = 1572)	SAPS B3 (<i>n</i> = 336)	BPRS 10 (<i>n</i> = 104)	Total (<i>N</i> = 2012)
1 'none'	1165 (74)	280 (83)	63 (61)	1508 (75)
2 'minimal'	163 (10)	25 (7)	23 (22)	211 (10)
3 'mild'	150 (10)	22 (7)	13 (13)	185 (9)
4 'moderate'	65 (4)	6 (2)	2 (2)	73 (4)
5 'marked'	22 (1)	2 (<1)	1 (1)	25 (1)
6 'severe'	7 (<1)	1 (<1)	2 (2)	10 (<1)

Data are *n* (%). PANSS, Positive and Negative Syndrome Scale; SAPS, Scale for the Assessment of Positive Symptoms; BPRS, Brief Psychiatric Rating Scale.

Table S4. Distribution of aggression scores in individuals with schizophrenia who underwent diffusion-weighted magnetic resonance imaging.

Score	Item		
	PANSS P7 (<i>n</i> = 630)	SAPS B3 (<i>n</i> = 100)	Total (<i>N</i> = 734)
1 'none'	441 (70)	98 (94)	539 (73)
2 'minimal'	83 (13)	3 (3)	86 (12)
3 'mild'	56 (9)	3 (3)	59 (8)
4 'moderate'	31 (5)	-	31 (4)
5 'marked'	14 (2)	-	14 (2)
6 'severe'	5 (<1)	-	5 (<1)

Data are *n* (%). PANSS, Positive and Negative Syndrome Scale; SAPS, Scale for the Assessment of Positive Symptoms.

Table S5. Deviations from normative values of cortical thickness (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia ($N = 2012$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GCT	-0.63 (-0.70, -0.56)	0.024	4.24×10^{-147}	1.03 (0.91, 1.16)	0.047	.571
OFC	0.00 (-0.06, 0.06)	0.022	.905	1.22 (1.07, 1.39)	0.061	1.00×10^{-4}
DLPFC	-0.09 (-0.14, -0.03)	0.022	1.37×10^{-4}	0.95 (0.82, 1.08)	0.050	.295
ACC	0.15 (0.10, 0.21)	0.023	9.16×10^{-12}	1.13 (0.99, 1.28)	0.057	.016
PCC	0.05 (-0.01, 0.10)	0.022	.042	1.03 (0.90, 1.17)	0.052	.603
INS	-0.05 (-0.11, 0.01)	0.022	.023	1.02 (0.90, 1.16)	0.051	.684
STG	-0.09 (-0.15, -0.03)	0.023	1.41×10^{-4}	1.07 (0.94, 1.21)	0.052	.183
PREC	0.14 (0.08, 0.19)	0.022	1.04×10^{-9}	0.91 (0.80, 1.04)	0.046	.060
IPL	0.02 (-0.03, 0.08)	0.022	.303	0.85 (0.75, 0.97)	0.042	.001

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GCT, global cortical thickness; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule.

Table S6. Deviations from normative values of surface area (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia ($N = 2009$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GSA	-0.21 (-0.26, 0.15)	0.023	8.75×10^{-20}	0.85 (0.76, 0.95)	0.039	3.20×10^{-4}
OFC	0.06 (0.01, 0.12)	0.022	.004	0.97 (0.86, 1.09)	0.046	.499
DLPFC	-0.05 (-0.10, .01)	0.022	.042	0.88 (0.77, 1.00)	0.044	.008
ACC	-0.13 (-0.19, -0.08)	0.023	2.45×10^{-9}	0.97 (0.85, 1.10)	0.050	.493
PCC	-0.03 (-0.08, 0.03)	0.022	.240	0.94 (0.83, 1.06)	0.046	.198
INS	0.04 (-0.02, 0.09)	0.022	.108	1.01 (0.89, 1.16)	0.052	.795
STG	0.09 (0.03, 0.15)	0.023	7.40×10^{-5}	1.17 (1.05, 1.30)	0.050	2.30×10^{-4}
PREC	0.08 (0.03, 0.14)	0.022	1.64×10^{-4}	0.97 (0.85, 1.10)	0.050	.510
IPL	-0.04 (-0.10, 0.02)	0.023	.068	0.82 (0.72, 0.93)	0.042	9.00×10^{-5}

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GSA, global surface area; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule.

Table S7. Deviations from normative values of fractional anisotropy (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia ($N = 733$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GFA	-0.18 (-0.27, -0.08)	0.037	2.00×10^{-6}	0.82 (0.69, 0.97)	0.054	.003
CC	-0.17 (-0.27, -0.07)	0.037	5.00×10^{-6}	0.88 (0.73, 1.06)	0.065	.083
CG	0.21 (0.10, 0.32)	0.043	1.00×10^{-6}	0.97 (0.79, 1.19)	0.079	.691
FOF	0.02 (-0.08, 0.11)	0.037	.603	0.97 (0.79, 1.20)	0.080	.713
SLF	0.13 (0.03, 0.23)	0.037	4.47×10^{-4}	1.12 (0.92, 1.35)	0.083	.145
SS	-0.05 (-0.15, 0.04)	0.037	.152	0.90 (0.73, 1.10)	0.070	.171
UNC	0.05 (-0.04, 0.15)	0.037	.150	1.05 (0.84, 1.31)	0.090	.576
CR	0.07 (-0.03, 0.16)	0.037	.060	1.27 (1.06, 1.53)	0.092	.001
IC	0.36 (0.26, 0.46)	0.038	1.86×10^{-21}	0.81 (0.70, 0.94)	0.047	2.80×10^{-4}
EC	0.12 (0.02, 0.21)	0.037	.001	1.15 (0.94, 1.41)	0.090	.075
CST	-0.03 (-0.12, 0.07)	0.037	.434	0.99 (0.82, 1.20)	0.074	.903
PTR	-0.01 (-0.10, 0.09)	0.037	.828	1.01 (0.84, 1.23)	0.076	.845
FX	-0.12 (-0.22, -0.02)	0.037	.001	1.01 (0.85, 1.19)	0.065	.896
FXST	-0.05 (-0.15, 0.05)	0.037	.175	1.10 (0.90, 1.35)	0.086	.212

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GFA, global fractional anisotropy; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.

Table S8. Deviations from normative values of mean diffusivity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia ($N = 607$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GMD	0.27 (0.16, 0.37)	0.041	1.04×10^{-10}	1.41 (1.18, 1.68)	0.097	8.30×10^{-7}
CC	0.10 (0.00, 0.21)	0.041	.011	1.13 (0.94, 1.35)	0.079	.086
CG	-0.15 (-0.26, -0.05)	0.041	2.25×10^{-4}	0.78 (0.64, 0.93)	0.056	3.80×10^{-4}
FOF	-0.15 (-0.25, -0.04)	0.041	3.40×10^{-4}	1.00 (0.81, 1.23)	0.080	.958
SLF	-0.04 (-0.15, 0.06)	0.041	.294	0.94 (0.78, 1.13)	0.068	.408
SS	-0.08 (-0.18, 0.03)	0.041	.061	0.82 (0.67, 1.00)	0.063	.010
UNC	0.12 (0.02, 0.23)	0.041	.002	1.23 (1.01, 1.49)	0.093	.007
CR	0.04 (-0.07, 0.14)	0.041	.381	0.95 (0.76, 1.18)	0.081	.560
IC	-0.09 (-0.19, 0.02)	0.041	.029	1.10 (0.93, 1.31)	0.072	.133
EC	-0.09 (-0.19, 0.02)	0.041	.034	1.02 (0.83, 1.25)	0.081	.841
CST	-0.19 (-0.30, -0.09)	0.041	2.00×10^{-6}	0.91 (0.76, 1.10)	0.067	.212
PTR	0.05 (-0.06, 0.15)	0.041	.246	1.04 (0.86, 1.27)	0.079	.582
FX	0.24 (0.13, 0.35)	0.042	1.09×10^{-8}	0.94 (0.77, 1.13)	0.070	.378
FXST	0.01 (-0.09, 0.11)	0.041	.809	1.09 (0.92, 1.29)	0.072	.188

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GMD, global mean diffusivity; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.

Table S9. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for current dosage of antipsychotic medication ($N = 1499$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
TCV	-0.33 (-0.40, -0.26)	0.026	1.67×10^{-35}	0.84 (0.73, 0.96)	0.046	.001
OFC	0.06 (-0.01, 0.12)	0.026	.029	1.09 (0.94, 1.28)	0.066	.136
DLPFC	-0.03 (-0.09, 0.04)	0.026	.326	0.85 (0.73, 1.00)	0.052	.009
ACC	-0.02 (-0.08, 0.05)	0.026	.519	0.98 (0.84, 1.14)	0.058	.715
PCC	0.06 (-0.01, 0.12)	0.026	.028	0.89 (0.77, 1.03)	0.050	.038
INS	0.07 (0.00, 0.14)	0.026	.008	1.06 (0.91, 1.24)	0.063	.325
STG	-0.06 (-0.13, 0.01)	0.027	.033	1.18 (1.02, 1.38)	0.069	.004
PREC	0.12 (0.05, 0.18)	0.026	8.00×10^{-6}	0.92 (0.79, 1.08)	0.057	.204
IPL	-0.04 (-0.10, 0.03)	0.026	.173	0.73 (0.62, 0.85)	0.045	2.30×10^{-7}
HIPPO	-0.13 (-0.20, -0.06)	0.027	1.00×10^{-6}	1.05 (0.91, 1.22)	0.060	.369
AMYG	0.06 (0.00, 0.13)	0.027	.015	1.01 (0.86, 1.18)	0.061	.892
STRI	0.36 (0.29, 0.43)	0.027	2.59×10^{-39}	1.10 (0.97, 1.25)	0.054	.050
THAL	-0.10 (-0.17, -0.03)	0.027	2.18×10^{-4}	1.09 (0.94, 1.27)	0.064	.120

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; TCV, total cortical volume; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule; HIPPO, hippocampus; AMYG, amygdala; STRI, striatum; THAL, thalamus.

Table S10. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for current dosage of antipsychotic medication ($N = 549$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GWM	-0.21 (-0.32, -0.10)	0.043	7.63×10^{-7}	0.70 (0.57, 0.86)	0.057	1.20×10^{-5}
CC	-0.14 (-0.25, -0.03)	0.043	9.30×10^{-4}	0.82 (0.65, 1.04)	0.075	.034
CG	0.22 (0.09, 0.036)	0.051	1.80×10^{-5}	1.11 (0.85, 1.45)	0.114	.311
FOF	0.11 (0.00, 0.22)	0.043	.010	0.98 (0.74, 1.30)	0.108	.842
SLF	0.07 (-0.05, 0.18)	0.043	.128	1.15 (0.89, 1.47)	0.111	.161
SS	0.00 (-0.11, 0.11)	0.043	.975	1.08 (0.83, 1.42)	0.114	.438
UNC	-0.10 (-0.21, 0.01)	0.043	.022	0.80 (0.61, 1.05)	0.084	.037
CR	-0.01 (-0.12, 0.10)	0.043	.869	1.30 (0.99, 1.71)	0.138	.012
IC	0.27 (0.16, 0.38)	0.043	6.55×10^{-10}	0.77 (0.64, 0.94)	0.059	6.50×10^{-4}
EC	0.11 (0.00, 0.22)	0.043	.008	1.12 (0.86, 1.47)	0.118	.276
CST	0.04 (-0.07, 0.15)	0.043	.304	1.08 (0.84, 1.39)	0.105	.453
PTR	-0.03 (-0.15, 0.08)	0.043	.417	0.91 (0.69, 1.20)	0.098	.386
FX	-0.17 (-0.28, -0.05)	0.044	1.46×10^{-4}	1.08 (0.88, 1.34)	0.089	.330
FXST	-0.03 (-0.14, 0.08)	0.043	.422	1.01 (0.78, 1.31)	0.101	.928

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GWM, global white matter; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.

Table S11. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for duration of illness ($N = 1680$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
TCV	-0.33 (-0.39, -0.27)	0.024	8.89×10^{-40}	0.88 (0.77, 1.01)	0.047	.015
OFC	0.01 (-0.06, 0.07)	0.025	.839	1.08 (0.92, 1.25)	0.064	.217
DLPFC	-0.05 (-0.11, 0.02)	0.025	.061	0.84 (0.72, 0.98)	0.051	.004
ACC	-0.04 (-0.10, 0.02)	0.025	.108	0.96 (0.82, 1.12)	0.057	.470
PCC	0.08 (0.02, 0.14)	0.024	9.65×10^{-4}	0.95 (0.82, 1.09)	0.053	.317
INS	0.05 (-0.01, 0.11)	0.025	.040	1.06 (0.91, 1.24)	0.063	.320
STG	-0.08 (-0.15, -0.02)	0.025	.001	1.15 (0.99, 1.33)	0.067	.019
PREC	0.05 (-0.02, 0.11)	0.025	.061	1.00 (0.85, 1.17)	0.060	.964
IPL	-0.02 (-0.08, 0.04)	0.025	.441	0.81 (0.69, 0.94)	0.049	3.50×10^{-4}
HIPPO	-0.07 (-0.13, 0.00)	0.025	.006	1.09 (0.94, 1.26)	0.063	.138
AMYG	0.09 (0.03, 0.16)	0.025	1.73×10^{-4}	1.05 (0.90, 1.23)	0.064	.398
STRI	0.26 (0.11, 0.32)	0.025	1.27×10^{-23}	1.08 (0.95, 1.22)	0.052	.116
THAL	-0.04 (-0.10, 0.03)	0.026	.162	1.05 (0.90, 1.22)	0.062	.434

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; TCV, total cortical volume; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule; HIPPO, hippocampus; AMYG, amygdala; STRI, striatum; THAL, thalamus.

Table S12. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for duration of illness ($N = 454$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GWM	-0.13 (-0.25, 0.00)	0.047	.008	0.60 (0.47, 0.78)	0.060	3.9×10^{-7}
CC	0.02 (-0.10, 0.14)	0.047	.633	0.76 (0.55, 1.04)	0.094	.026
CG	0.24 (0.08, 0.39)	0.060	8.50×10^{-5}	1.08 (0.79, 1.49)	0.133	.518
FOF	0.17 (0.04, 0.29)	0.047	4.84×10^{-4}	1.15 (0.79, 1.67)	0.167	.329
SLF	0.03 (-0.09, 0.15)	0.047	.505	1.17 (0.88, 1.57)	0.133	.162
SS	0.10 (-0.02, 0.22)	0.047	.030	1.10 (0.79, 1.54)	0.144	.460
UNC	0.01 (-0.12, 0.13)	0.047	.894	0.67 (0.46, 0.97)	0.096	.005
CR	-0.01 (-0.13, 0.11)	0.047	.833	1.40 (0.99, 1.98)	0.188	.011
IC	0.20 (0.08, 0.33)	0.047	1.70×10^{-5}	0.72 (0.57, 0.91)	0.066	3.30×10^{-4}
EC	0.05 (-0.07, 0.17)	0.047	.307	1.30 (0.92, 1.85)	0.178	.053
CST	-0.06 (-0.18, 0.06)	0.047	.193	1.19 (0.85, 1.68)	0.158	.181
PTR	0.01 (-0.11, 0.13)	0.047	.879	0.91 (0.63, 1.30)	0.126	.483
FX	-0.16 (-0.28, -0.04)	0.048	9.05×10^{-4}	1.55 (1.15, 2.08)	0.178	1.50×10^{-4}
FXST	0.00 (-0.13, 0.12)	0.047	.938	0.97 (0.69, 1.35)	0.124	.788

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GWM, global white matter; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.

Table S13. Deviations from normative values of grey matter volume (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for lifetime substance use disorder ($N = 778$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
TCV	-0.23 (-0.32, -0.13)	0.036	5.00×10^{-10}	0.88 (0.75, 1.05)	0.058	.063
OFC	0.10 (0.00, 0.19)	0.036	.008	0.98 (0.81, 1.19)	0.073	.827
DLPFC	0.08 (-0.01, 0.18)	0.036	.019	0.82 (0.68, 0.99)	0.061	.008
ACC	-0.01 (-0.10, 0.09)	0.036	.878	0.98 (0.81, 1.18)	0.072	.777
PCC	0.11 (0.02, 0.21)	0.036	.002	0.99 (0.82, 1.20)	0.075	.901
INS	0.04 (-0.05, 0.14)	0.036	.242	0.97 (0.80, 1.17)	0.072	.660
STG	-0.15 (-0.24, -0.05)	0.038	1.28×10^{-4}	1.03 (0.86, 1.24)	0.073	.665
PREC	0.16 (0.07, 0.25)	0.036	8.00×10^{-6}	0.98 (0.81, 1.18)	0.071	.739
IPL	0.10 (0.01, 0.19)	0.036	.005	0.82 (0.68, 0.99)	0.060	.006
HIPPO	-0.20 (-0.30, -0.10)	0.038	1.05×10^{-7}	0.92 (0.76, 1.12)	0.069	.267
AMYG	-0.06 (-0.16, 0.04)	0.038	.112	0.95 (0.77, 1.17)	0.078	.533
STRI	0.25 (0.15, 0.35)	0.038	3.99×10^{-11}	1.19 (1.00, 1.42)	0.081	.011
THAL	-0.19 (-0.29, -0.08)	0.040	3.00×10^{-6}	0.92 (0.73, 1.15)	0.080	.321

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; TCV, total cortical volume; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule; HIPPO, hippocampus; AMYG, amygdala; STRI, striatum; THAL, thalamus.

Table S14. Deviations from normative values of white matter microstructural integrity (A) and their associations with aggressive behaviour (B) in individuals with schizophrenia, adjusting for lifetime substance use disorder ($N = 217$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GWM	-0.18 (-0.36, 0.00)	0.068	.009	0.73 (0.55, 0.97)	0.081	.005
CC	-0.15 (-0.33, 0.02)	0.068	.026	0.91 (0.70, 1.20)	0.095	.389
CG	0.26 (0.08, 0.43)	0.068	2.04×10^{-4}	1.11 (0.77, 1.59)	0.155	.476
FOF	0.17 (0.00, 0.35)	0.068	.012	0.85 (0.59, 1.23)	0.121	.257
SLF	0.11 (-0.07, 0.28)	0.068	.121	1.19 (0.83, 1.72)	0.171	.216
SS	0.01 (-0.17, 0.18)	0.068	.910	1.13 (0.75, 1.70)	0.179	.429
UNC	-0.11 (-0.29, 0.07)	0.068	.108	0.85 (0.63, 1.16)	0.102	.183
CR	-0.02 (-0.20, 0.15)	0.068	.744	1.31 (0.92, 1.86)	0.178	.045
IC	0.40 (0.23, 0.58)	0.068	1.20×10^{-8}	0.94 (0.67, 1.32)	0.124	.637
EC	0.17 (-0.01, 0.35)	0.068	.012	1.24 (0.86, 1.79)	0.177	.137
CST	0.08 (-0.09, 0.26)	0.068	.224	0.93 (0.65, 1.33)	0.129	.593
PTR	-0.03 (-0.21, 0.15)	0.068	.664	0.99 (0.69, 1.41)	0.136	.935
FX	-0.25 (-0.43, -0.08)	0.069	2.52×10^{-4}	0.93 (0.68, 1.26)	0.111	.518
FXST	-0.09 (-0.27, 0.09)	0.068	.190	1.12 (0.75, 1.67)	0.174	.472

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GWM, global white matter; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.

Table S15. Deviations from normative values of grey matter volume (A) and their associations with scores on PANSS item P7 (B) in individuals with schizophrenia ($N = 1569$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
TCV	-0.41 (-0.48, -0.34)	0.026	1.33×10^{-54}	0.84 (0.74, 0.96)	0.043	.001
OFC	0.07 (0.00, 0.13)	0.025	.011	1.04 (0.90, 1.21)	0.060	.447
DLPFC	-0.07 (-0.14, -0.01)	0.025	.004	0.83 (0.71, 0.97)	0.049	.002
ACC	-0.06 (-0.13, 0.01)	0.025	.019	0.99 (0.85, 1.14)	0.056	.813
PCC	0.11 (0.04, 0.18)	0.025	1.40×10^{-5}	0.92 (0.80, 1.06)	0.049	.118
INS	0.07 (0.00, 0.13)	0.025	.006	1.03 (0.89, 1.19)	0.058	.625
STG	-0.10 (-0.17, -0.03)	0.026	1.03×10^{-4}	1.14 (0.99, 1.32)	0.065	.017
PREC	0.16 (0.10, 0.23)	0.025	1.20×10^{-10}	0.93 (0.80, 1.09)	0.055	.233
IPL	-0.02 (-0.08, 0.05)	0.025	.545	0.75 (0.65, 0.87)	0.044	7.70×10^{-7}
HIPPO	-0.20 (-0.26, -0.13)	0.026	7.57×10^{-14}	1.04 (0.90, 1.20)	0.057	.474
AMYG	0.03 (-0.04, 0.10)	0.026	.262	1.03 (0.88, 1.19)	0.060	.639
STRI	0.33 (0.26, 0.39)	0.027	1.14×10^{-34}	1.08 (0.96, 1.23)	0.052	.098
THAL	-0.21 (-0.28, -0.14)	0.027	4.48×10^{-15}	1.08 (0.94, 1.26)	0.062	.159

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; TCV, total cortical volume; OFC, orbitofrontal cortex; DLPFC, dorsolateral prefrontal cortex; ACC, anterior cingulate cortex; PCC, posterior cingulate cortex; INS, insula; STG, superior temporal gyrus; PREC, precuneus; IPL, inferior parietal lobule; HIPPO, hippocampus; AMYG, amygdala; STRI, striatum; THAL, thalamus.

Table S16. Deviations from normative values of white matter microstructural integrity (A) and their associations with scores on PANSS item P7 (B) in individuals with schizophrenia ($N = 504$).

IDP	A			B		
	d (99% CI)	SE	p	OR (99% CI)	SE	p
GWM	-0.33 (-0.44, -0.21)	0.046	9.99×10^{-13}	0.77 (0.62, 0.94)	0.061	.001
CC	-0.18 (-0.29, -0.06)	0.045	8.20×10^{-5}	0.83 (0.66, 1.05)	0.076	.047
CG	0.33 (0.19, 0.48)	0.056	2.79×10^{-9}	1.14 (0.86, 1.52)	0.125	.217
FOF	0.16 (0.04, 0.27)	0.045	5.84×10^{-4}	0.91 (0.67, 1.23)	0.107	.412
SLF	0.12 (0.01, 0.24)	0.045	.005	1.11 (0.87, 1.40)	0.102	.277
SS	0.04 (-0.07, 0.16)	0.045	.348	1.05 (0.80, 1.36)	0.107	.663
UNC	-0.06 (-0.18, 0.05)	0.045	.147	0.81 (0.62, 1.05)	0.084	.039
CR	0.04 (-0.07, 0.16)	0.045	.357	1.26 (0.96, 1.65)	0.133	.028
IC	0.34 (0.22, 0.46)	0.046	1.31×10^{-13}	0.73 (0.60, 0.89)	0.055	3.40×10^{-5}
EC	0.16 (0.04, 0.27)	0.045	3.95×10^{-4}	1.03 (0.78, 1.35)	0.108	.797
CST	0.06 (-0.05, 0.18)	0.045	.171	1.12 (0.87, 1.45)	0.111	.251
PTR	0.00 (-0.12, 0.11)	0.045	.954	0.89 (0.67, 1.17)	0.096	.265
FX	-0.24 (-0.36, -0.12)	0.046	1.36×10^{-7}	1.13 (0.91, 1.39)	0.092	.147
FXST	-0.03 (-0.15, 0.08)	0.045	.482	1.01 (0.78, 1.30)	0.100	.941

IDP, imaging-derived phenotype; CI, confidence interval; SE, standard error; OR, odds ratio; GWM, global white matter; CC, corpus callosum; CG, cingulum; FOF, fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; UNC, uncinate fasciculus; CR, corona radiata; IC, internal capsule; EC, external capsule; CST, corticospinal tract; PTR, posterior thalamic radiations; FX, fornix; FXST, fornix stria terminalis.