Supplemental material

The Cost Outcome Pathway Framework: Integrating socio-economic impacts to Adverse Outcome Pathways

Thibaut Coustillet, Xavier Coumoul, Anne-Sophie Villégier, Michèle Bisson, Ellen Fritsche, Jean-Marc Brignon, Florence Zeman, Karine Audouze

Table of Contents

Supplementary Tables

Supplementary Table 1. Full distribution of stressor-event relationships from AOPhelpFinder results. For convenience, two sheets are supplied. Sheet 'stressor-event' provides an overview of stressor-based links, while Sheet 'event-stressor' provides an overview of event-based links. Data between the two sheets is strictly identical. https://github.com/systox1124/bioRxiv-COP-publication.git

Supplementary Table 2. Full distribution of event-event relationships from AOPhelpFinder results.

https://github.com/systox1124/bioRxiv-COP-publication.git

Supplementary Table 3. PubTator bioconcepts found for each article used in the development of the COP. Sheet 'all_bioconcepts' includes all the bio-concepts detected in each category for each article. Sheet 'top_bioconcepts' contains only the most frequently detected bio-concept for each category for each article.

https://github.com/systox1124/bioRxiv-COP-publication.git

Supplementary Table 4: List of the 24 pesticides. The selected chemicals were divided into two main categories: organophosphates (n=3) and their metabolites (n=4), and pyrethroids of type I (n=7), of type II (n=6) and their metabolites (n=3). Piperonyl-butoxide was considered as a separate category. a, Dereumeaux et al., 2016; b, Béranger et al., 2020; c, Tagne-Fosto et al., 2022; d, Anses, 2011; e, Cnam_IHIE, 2005; f, Anses, 2019; g, BNVD, 2011-2015. 3-BPA, 3-phenoxybenzoic acid; Cl2CA, cis-3-(2,2dichlorovinyl)-2,2-dimethylcyclopropane-

carboxylic acid; ClCF3CA, 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2dimethylcyclopropanecarboxylic acid; TCPy, 3,5,6-trichloro-2-pyridinol.

Supplementary Table 5. List of the 248 conventional events (MIE, KE, AO) and five CO-related terms used for searches with AOP-helpFinder v2.

https://github.com/systox1124/bioRxiv-COP-publication.git

Supplementary Figures

Supplementary Figure 1. Distribution of stressor-event links according to the 23 most common stressors, representing 100% of the total dataset (23 stressors and 2404 links). No link was found for the stressor CICF3CA.

Supplementary Figure 2. Distribution of stressor-event links according to the 20 most common events, representing 74.4% of the total dataset (141 distinct events and 2404 links).

Supplementary Figure 3. Distribution of event-event links according to the 20 most common events, representing 61.0% of the total dataset (225 distinct events and 73079 links).

KER Justification

Each KER's argumentation.

Supplementary references

Supplementary Table 4: List of the 24 pesticides. The selected chemicals were divided into two main categories: organophosphates (n=3) and their metabolites (n=4), and pyrethroids of type I (n=7), of type II (n=6) and their metabolites (n=3). Piperonyl-butoxide was considered as a separate category. a, Dereumeaux et al., 2016¹; b, Béranger et al., 2020²; c, Tagne-Fosto et al., 2022³; d, Anses, 2011⁴; e, Cnam_IHIE, 2005⁵; f, Anses, 2019⁶; g, BNVD, 2011-2015. 3-BPA, 3-phenoxybenzoic acid; Cl2CA , cis-3-(2,2dichlorovinyl)-2,2-dimethylcyclopropane-carboxylic acid; ClCF3CA, 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylic acid; TCPy, 3,5,6-trichloro-2-pyridinol.

Family	Substance name	CAS number	Туре	French HBM data			Food contami nation in France	Domestic use in France		Agricult ural use in France 2011- 2015
				Elfe cohort (detected in urine) ^a	Elfe cohort (detected in hair) _b	Esteban study (detected in urine)c	EAT2 (detected)d	Cnam- IHIE study ^e	Pesti'HO ME study ^f	BnVD database g
Pyrethroi d	Allethrin	584-79-2	type I pyreth roid					X		X
	Bifenthrin	82657- 04-3	type I pyreth roid				X	X	x	x
	Cyfluthrin	68359- 37-5	type II pyreth roid				x	X	x	x
	Cypermeth rin	52315- 07-8	tenfan ype II pyreth roid		X			X	x	X
	Deltameth rin	52918- 63-5	type II pyreth roid					X	x	X
	lambda- cyhalothri n	91465- 08-6	type II pyreth roid		X		X		x	X
	Esfenvaler ate	66230- 04-4	type II pyreth roid						x	X
	Permethri	52645-	type I		X		X	X	X	X

	n	53-1	pyreth roid							
	Resmethri n	10453- 86-8	type I pyreth roid						X	X
	Phenothrin	26002- 80-2	type I pyreth roid					X	X	
	tau- fluvalinate	102851- 06-9	type II pyreth roid						X	X
	Tefluthrin	79538- 32-2	type I pyreth roid							X
	Tetrameth rin	7696-12- 0	type I pyreth roid					X		X
	(3-PBA)	3739-38- 6	metab olite (vario us pyreth roids)	X	x					
	CI2CA	55701- 05-8	metab olite (cyflut hrin)	x	x					
	CICF3CA	72748- 35-7	metab olite (teflut hrin)		x					
Organo phosphat e	Chlorpyrif os	2921-88- 2	organo phosp hate			x	X	X	X	X
	Chlorpyrif os-methyl	5598-13- 0	organo phosp hate			X	X			X
	Ethoproph os	13194- 48-4	organo phosp hate				X			X
	Chlorpyrif os-oxon	5598-15- 2	metab olite (chlor pyrifos)			X				
	ТСРу	6515-38- 4	metab olite (chlor pyrifos)		x	x				
	diethyl-	598-02-7	metab	X	X	X				

	phosphate		olite (vario us organo phosp hates)						
	diethyl- thiophosph ate	2465-65- 8	metab olite (vario us organo phosp hates)	X	X	X			
Synergist	Piperonyl butoxide	51-03-6	pyreth roid synerg ist				X	X	X



Supplementary Figure 1. Distribution of the identified stressor-event linkage in the literature using AOP-helpFinder for the 23 retrieved stressors, representing 100% of the total dataset (23 stressors and 2404 links). No link was found for the pyrethroid metabolite of tefluthrin (CICF3CA).



Distribution of stressor-event links according to the 20 most common events, representing 74.4% of the total data set (141 distinct events and 2404 links)

Supplementary Figure 2. Distribution of stressor-event links according to the 20 most

common events, representing 74.4% of the total dataset (141 distinct events and 2404 links).



Distribution of event-event links according to the 20 most common events, representing 61.0% of the total data set (225 distinct events and 73079 links)

Supplementary Figure 3. Distribution of event-event links according to the 20 most

common events, representing 61.0% of the total dataset (225 distinct events and 73079 links).

KER justification:

1- KER 1-2: Activation, IP3R (KE ID 2037) and Activation, RyR (KE ID 2186) lead to Increased, glutamate (KE ID 1350).

IP3R and RyRs are expressed by many types of neurons⁷. Once they are activated (e.g., OPs such as CPF or TOCP (tri-o-cresyl phosphate)), a release of Ca²⁺ from intracellular stores occurs in the cytoplasm which triggers the exocytosis of glutamate. Separately, IP3Rs and RyRs agonists were capable of increasing glutamate levels in nerve tissue, while their antagonists did not reverse the effect, suggesting the participation of both receptors in the glutamate homeostasis^{8,9}.

2- KER 3-4-9-10: Increased, glutamate (KE ID 1350) leads to decreased, Bcl-2 expression (KE ID 2123) and increased, Bax expression (KE ID 2124), resulting in cell injury/death (KE ID 55)

Glutamate is a major excitatory neurotransmitter in the mammalian central nervous system (CNS). Neurons and glial cells, including microglia, express glutamate receptors. Bax activation leads to the mitochondrial release of cytochrome c leading to the activation of caspases (3 & 9) and subsequently to apoptosis¹⁰. In several CNS cell lines^{11,12}, glutamate down-regulates Bcl-2 with a concomitant up-regulation of Bax. Moreover, Lee *et al.* showed that CPF increases the activity of caspases 3 and 9 in human embryonic stem cell-derived neural precursor cells (hNPCs) and cytochrome c mitochondrial release and cell death¹³.

3- KER 5-6-7-8: Increased, glutamate (KE ID 1350) leads to Activation, microglia (KE ID 1998) and Activation, microglia leads to neuroinflammation (KE ID 188), resulting in cell injury/death (KE ID 55)

Neuroinflammation largely depends on microglia which are resident immune cells of the nerve tissue, constantly scavenging the CNS for damaged neurons, foreign material, or infectious agents (mainly through phagocytosis). In addition to releasing pro-inflammatory mediators such as IL-1 β or TNF- α , microglial cells could also deliver cytotoxic effectors such as proteases, reactive oxygen intermediates and nitrite oxide (NO)¹⁴ (immunoexcitotoxicity). Glutamate-induced overstimulation of microglia leads to microglia-induced

neuroinflammation, which in turn can strengthen inflammation by operating in an autocrine fashion¹⁵. Dichlorvos (an OP) exposure activates microglia and causes secretion of IL-1 β , TNF- α , and NO leading to neuronal death¹⁶ *via* p53, Bax, cytochrome c and caspase 3. In summary, a continuous neuroinflammation (due to the persistence of stressors) through the release of proinflammatory cytokines (IL-1 β , TNF- α) and NO, triggers apoptotic processes (involving p53, Bax, cytochrome c and caspase 3, for example) which ultimately leads to cell death (immunoexcitotoxicity).

4- KER 11: cell injury/death (KE ID 55) leads to Hippocampal Physiology, Altered (KE ID 758) resulting in Impairment, learning and memory (KE ID 341).

Stresses during childhood can lead to lowered hippocampal volume and neurogenesis in humans and animals¹⁷ due to impaired maturation of the brain. Indeed, in adolescent rats (PND 28-48, a window of vulnerability), alcohol exposure results in a reduced neurogenesis with lowered differentiating neurons that were due at least to an increase of cell death of immature neurons rather than a decrease in neural progenitor cell proliferation¹⁸. Interestingly, the loss of neurogenesis could be both period- and region- specific. In another study, cell death in the CA1 region (of the Ammon's corn) and in the dentate gyrus in PND-11 Sprague–Dawley rats leads to a decline in memory and learning efficiency later in life (PND-60, young adults); indeed, these rats showed poorer results in the Morris water maze test, both during the training and testing stages¹⁹. Difficulties in the training phase reflect learning disorders, while difficulties in the testing phase reflect memory disorders. Besides, authors showed that the hippocampal neurodegeneration could partly occur through Bcl-2 downregulation. On the other hand, other researchers observed longer times and distances to reach the platform in the Morris water maze test in rats with a 2.3% reduction in hippocampal volume²⁰.

Supplementary references

1. Dereumeaux, C. *et al.* Biomarkers of exposure to environmental contaminants in French pregnant women from the Elfe cohort in 2011. *Environ. Int.* **97**, 56–67 (2016).

2. Béranger, R. *et al.* Multiple pesticide analysis in hair samples of pregnant French women: Results from the ELFE national birth cohort. *Environ. Int.* **120**, 43–53 (2018).

3. Tagne-Fotso, R. et al. Exposure of the General French Population to Herbicides, Pyrethroids, Organophosphates, Organochlorines, and Carbamate Pesticides in 2014–2016: Results from the Esteban Study. https://www.ssrn.com/abstract=4452103 (2023) doi:10.2139/ssrn.4452103.

4. French Agency for Food, Environmental and Occupational Health & Safety (ANSES). *French Total Diet Study 2: Pesticide Residues, Additives, Acrylamide, Polycyclic Aromatic Hydrocarbons*. https://www.anses.fr/fr/system/files/PASER2006sa0361Ra2.pdf (2011).

5. Auburtin, G., Lecomte, J. & Moreau, J. *L'utilisation Des Biocides En Milieu Domestique et La Perception Des Risques Liés à Cette Utilisation Dans Une Population Française*. https://primequal.fr/sites/default/files/auburtint_rf.pdf (2005).

6. French Agency for Food, Environmental and Occupational Health & Safety (ANSES). *Pesti 'home Study: National Survey of Domestic Pesticide Use.*

https://www.anses.fr/fr/system/files/2019Pestihome.pdf (2019).

7. Ogawa Y. & Murayama T. Ryanodine receptors in the central nervous system. *Folia Pharmacol. Jpn.* **105**, 423–430 (1995).

8. Yamamura, S. *et al.* Effects of zonisamide on neurotransmitter release associated with inositol triphosphate receptors. *Neurosci. Lett.* **454**, 91–96 (2009).

9. Mori, F., Okada, M., Tomiyama, M., Kaneko, S. & Wakabayashi, K. Effects of ryanodine receptor activation on neurotransmitter release and neuronal cell death following kainic acid-induced status epilepticus. *Epilepsy Res.* **65**, 59–70 (2005).

10. Zou, H., Li, Y., Liu, X. & Wang, X. An APAF-1.cytochrome c multimeric complex is a functional apoptosome that activates procaspase-9. *J. Biol. Chem.* **274**, 11549–11556 (1999).

11. Nishi, T. *et al.* Participation of Bcl-2/Bax- α in Glutamate-induced Apoptosis of Human Glioblastoma Cells. *J. Neurooncol.* **44**, 109–117 (1999).

12. Gao, M. *et al.* Pinocembrin prevents glutamate-induced apoptosis in SH-SY5Y neuronal cells via decrease of bax/bcl-2 ratio. *Eur. J. Pharmacol.* **591**, 73–79 (2008).

13. Lee, J. E., Lim, M. S., Park, J. H., Park, C. H. & Koh, H. C. Nuclear NF-κB contributes to chlorpyrifos-induced apoptosis through p53 signaling in human neural precursor cells. *NeuroToxicology* **42**, 58–70 (2014).

14. Gehrmann, J., Matsumoto, Y. & Kreutzberg, G. W. Microglia: intrinsic immuneffector cell of the brain. *Brain Res. Brain Res. Rev.* **20**, 269–287 (1995).

15. Iovino, L., Tremblay, M. E. & Civiero, L. Glutamate-induced excitotoxicity in Parkinson's disease: The role of glial cells. *J. Pharmacol. Sci.* **144**, 151–164 (2020).

16. Sunkaria, A., Sharma, D. R., Wani, W. Y. & Gill, K. D. Attenuation of Dichlorvos-Induced Microglial Activation and Neuronal Apoptosis by 4-Hydroxy TEMPO. *Mol. Neurobiol.* **49**, 163–175 (2014).

17. Malave, L., Van Dijk, M. T. & Anacker, C. Early life adversity shapes neural circuit function during sensitive postnatal developmental periods. *Transl. Psychiatry* **12**, 306 (2022).

18. Broadwater, M. A., Liu, W., Crews, F. T. & Spear, L. P. Persistent Loss of Hippocampal Neurogenesis and Increased Cell Death following Adolescent, but Not Adult, Chronic Ethanol Exposure. *Dev. Neurosci.* **36**, 297–305 (2014).

19. Huang, L., Liu, Y., Jin, W., Ji, X. & Dong, Z. Ketamine potentiates hippocampal neurodegeneration and persistent learning and memory impairment through the PKCγ–ERK signaling pathway in the developing brain. *Brain Res.* **1476**, 164–171 (2012).

20. Su, J., Sripanidkulchai, K., Hu, Y., Wyss, J. M. & Sripanidkulchai, B. The Effect of Ovariectomy on Learning and Memory and Relationship to Changes in Brain Volume and Neuronal Density. *Int. J. Neurosci.* **122**, 549–559 (2012).