

**Table A1** Characteristics of the eleven hospitals from the Southwest region of the Netherlands<sup>1</sup>.

Characteristics	Hospital 1	Hospital 2	Hospital 3	Hospital 4	Hospital 5	Hospital 6	Hospital 7	Hospital 8	Hospital 9	Hospital 10	Hospital 11
	University hospital	Non-university teaching hospital	Non-university teaching hospital	Non-university teaching hospital	Non-university teaching hospital	Non-university teaching hospital	Non-university teaching hospital	Non-teaching hospital	Non-teaching hospital	Non-teaching hospital	Non-teaching hospital
No. of hospital beds	1233 (including Sophia children's hospital)	Not available	564	484	341	332	Not available	364	≈290	194	189
No. of single-occupancy rooms	576 (excluding Sophia children's hospital)	145	80	248	26	84 (including ICU and mother/child centre)	4	49	19	50	33
No. of isolation rooms with anteroom	54	62	33	66	12	7	0	18	7	10	5
Clinical hospital admissions	30,771	32,022	21,249	24,561	27,726	13,485	Not available	35,475	≈10,000	16,340	9,506
Outpatient clinic visits	659,317	402,758	29,5567	107,249 <sup>2</sup>	76,729 <sup>2</sup>	266,296 <sup>3</sup>	126,898	307,226 <sup>3</sup>	≈50,000	42,000	165,695

ICU intensive care unit.

<sup>1</sup>Data are from 2021.

<sup>2</sup> Only number of first outpatient clinic visits available.

<sup>3</sup> Only number of consultations available.

**Table A2** IPC measures recommended by the Dutch, ESCMID and ECDC guidelines.

	<b>Dutch guidelines</b>		<b>ESCMID guideline [1]</b>		<b>ECDC guideline on CPE [2]</b>
<b>IPC measures</b>	Strict isolation [3]	Contact isolation [4]	Contact precautions: endemic MDR- <i>P. aeruginosa</i>	Contact precautions: endemic ESBL-E (with the exception of <i>Escherichia coli</i> )	Contact precautions
<b>Type of room</b>	Isolation room <sup>1</sup>	Single-occupancy room, or under exceptional circumstances a multiple-occupancy room provided there is a 1.5 meter space around the patient's bed	Isolation room	Single room	Isolation room with en suite or bathroom facilities designated for use by known carriers, or commodes
<i>Door allowed to be open</i>	N/A	Yes	N/A	No recommendations	N/A
<i>Patient allowed to leave the room (for other reasons than e.g., operations, radiology)</i>	No	Yes, under specific conditions	No recommendations	No recommendations	No recommendations
<b>PPE for HCW</b>	Gloves before entering the room	Gloves during contact with the patient or contact with the patient's direct environment	Gloves before entering the room	Gloves before entering the room	Gloves during direct contact with the patient or their immediate surroundings and/or surfaces.
	Long-sleeved gown before entering the room		Gown before entering the room	Gown before entering the room	Gown/apron during direct contact with the patient or their immediate surroundings and/or surfaces.
	Mask (type of mask dependent on the type of				

	microorganism) before entering the room				
	Hair cap (only for MRSA)				
<b>Physiotherapy for inpatients</b>	No recommendations on whether and where physiotherapy can take place, and on recommended PPE use	No recommendations on whether and where physiotherapy can take place, and on recommended PPE use	No recommendations on whether and where physiotherapy can take place, and on recommended PPE use	No recommendations on whether and where physiotherapy can take place, and on recommended PPE use	No recommendations on whether and where physiotherapy can take place, and on recommended PPE use
<b>Visitors of inpatients</b>					
<i>PPE</i>	Mask (type of mask dependent on the type of microorganism)	No specific measures	No recommendations	No recommendations	No recommendations
	Gown				
<i>Other</i>	Visitor is not allowed to visit another patient	No specific measures	No recommendations	No recommendations	No recommendations
<i>Rooming-in</i>	No recommendations	No recommendations	No recommendations	No recommendations	No recommendations
<b>Cleaning and disinfection</b>	Daily cleaning of the patient room and anteroom and terminal cleaning and disinfection before admission of a new patient	Daily cleaning of the patient room and terminal cleaning and disinfection before admission of a new patient	Implement regular environmental cleaning procedures, which include detergents or disinfectants, depending on local practice to reduce the transmission rate. Ensure cleaning of patient	Implement regular environmental cleaning procedures, which include detergents or disinfectants, depending on local practice to reduce the transmission rate. Ensure cleaning of patient	Enhanced cleaning should be performed, especially for areas in close proximity to CRE carriers. Terminal disinfection of rooms should be performed upon transfer or discharge of patients.

---

care equipment  
and the  
environment.

care equipment  
and the  
environment.

---

ECDC European Centre for Disease Prevention and Control, *ESBL-E* extended-spectrum  $\beta$ -lactamase-producing Enterobacterales, *ESCMID* European Society of Clinical Microbiology and Infectious Diseases, *HCW* healthcare worker, *IPC* infection prevention and control, *MDR-P.* aeruginosa multidrug-resistant *Pseudomonas aeruginosa*, *MRSA* meticillin-resistant *Staphylococcus aureus*, *N/A* not applicable, *PPE* personal protective equipment.

<sup>1</sup> An isolation room is defined as a single-occupancy room with anteroom and pressure differences [5].

## References

- [1] Tacconelli E, Cataldo MA, Dancer SJ, De Angelis G, Falcone M, Frank U, et al. ESCMID guidelines for the management of the infection control measures to reduce transmission of multidrug-resistant Gram-negative bacteria in hospitalized patients. Clin Microbiol Infect 2014;20 Suppl 1:1-55.
- [2] Magiorakos AP, Burns K, Rodríguez Baño J, Borg M, Daikos G, Dumpis U, et al. Infection prevention and control measures and tools for the prevention of entry of carbapenem-resistant Enterobacteriaceae into healthcare settings: guidance from the European Centre for Disease Prevention and Control. Antimicrob Resist Infect Control 2017;6:113.
- [3] Dutch Working Party on Infection Prevention (WIP). Strikte isolatie. 2006.
- [4] Dutch Working Party on Infection Prevention (WIP). Contactisolatie. 2006.
- [5] Dutch Working Party on Infection Prevention (WIP). Bouw- en inrichtingseisen isolatie-afdeling Ventilatie isolatiekamers. 2004.

**Table A3** IPC measures for visitors of inpatients (n=11 hospitals).

<b>Transmission-based precautions</b>	<b>CPE</b>	<b>CPPA</b>	<b>ESBL-E</b>
<b>IPC measures for visitor(s)<sup>1</sup></b>	11 (100)	11 (100)	10 (90.9)
Hand disinfection	11 (100)	11 (100)	10 (100)
Disposable gown and non-sterile gloves	1 (9.1)	1 (9.1)	1 (10.0)
Disposable gown	1 (9.1)	1 (9.1)	0 (0.0)
Surgical mask type IIR, disposable gown, non-sterile gloves, and hair cap	1 (9.1)	1 (9.1)	0 (0.0)
<b>Temporary room leave by visitor(s) for a toilet visit is allowed</b>			
Yes	4 (36.4)	4 (36.4)	5 (45.5)
No	6 (54.5)	6 (54.5)	4 (36.4)
Not applicable <sup>2</sup>	1 (9.1)	1 (9.1)	2 (18.2)
<b>Temporary room leave by visitor(s) for use of the coffee area is allowed</b>			
Yes	4 (36.4)	4 (36.4)	6 (54.5)
No	6 (54.5)	6 (54.5)	3 (27.3)
Not applicable <sup>2</sup>	1 (9.1)	1 (9.1)	2 (18.2)
<b>Rooming-in is allowed</b>			
Yes	10 (90.9)	10 (90.9)	9 (81.8)
No	1 (9.1)	1 (9.1)	1 (9.1)
Not applicable	0 (0.0)	0 (0.0)	1 (9.1)
<b>Temporary leave of rooming-in visitor from patient room is allowed</b>			
Yes	5 (45.5)	5 (45.5)	4 (36.4)
No	5 (45.5)	5 (45.5)	5 (45.5)
Not applicable	1 (9.1)	1 (9.1)	2 (18.2)

Data are N (%). *CPPA* carbapenemase-producing *Pseudomonas aeruginosa*, *CPE* carbapenemase-producing Enterobacterales, *ESBL-E* extended-spectrum  $\beta$ -lactamase-producing Enterobacterales, *IPC* infection prevention and control, *N/A* not applicable.

<sup>1</sup> Multiple answers per hospital possible.

<sup>2</sup> One hospital has no specific policy on this topic in place.

**Table A4** Cleaning and disinfection after discharge of the patient (n=11 hospitals).

	CPE	CPPA	ESBL-E
<b>Cleaning product for final cleaning<sup>1,2</sup></b>			
Reusable microfiber cloth	2 (18.2)	2 (18.2)	2 (18.2)
Reusable mop	2 (18.2)	2 (18.2)	2 (18.2)
Disposable (microfiber) cloth	5 (45.5)	5 (45.5)	5 (45.5)
Disposable mop	5 (45.5)	5 (45.5)	5 (45.5)
No separate cleaning <sup>3</sup>	3 (27.3)	3 (27.3)	2 (18.2)
Not applicable	0 (0.0)	0 (0.0)	1 (9.1)
<b>Method or product for final disinfection<sup>1,2</sup></b>			
Chlorine 250 ppm	5 (45.5)	5 (45.5)	5 (45.5)
Chlorine 1000 ppm	2 (18.2)	2 (18.2)	1 (9.1)
Alcohol 70%	6 (54.5)	6 (54.5)	4 (36.4)
2-in-1 cleaning and disinfection product	3 (27.3)	3 (27.3)	2 (18.2)
Hydrogen peroxide wipes	1 (9.1)	1 (9.1)	1 (9.1)
No disinfection	0 (0.0)	0 (0.0)	1 (9.1)
Not applicable	0 (0.0)	0 (0.0)	1 (9.1)
<b>Changing separation curtains after isolation</b>			
Yes	9 (81.8)	9 (81.8)	7 (63.6)
No	2 (18.2) <sup>4</sup>	2 (18.2) <sup>4</sup>	3 (27.3) <sup>4</sup>
Not applicable	0 (0.0)	0 (0.0)	1 (9.1)
<b>Changing window curtains after isolation</b>			
Yes	1 (9.1)	1 (9.1)	0 (0.0)
No	10 (90.9) <sup>4</sup>	10 (90.9) <sup>4</sup>	10 (90.9) <sup>4</sup>
Not applicable	0 (0.0)	0 (0.0)	1 (9.1)

Data are N (%). CPPA carbapenemase-producing *Pseudomonas aeruginosa*, CPE carbapenemase-producing Enterobacterales, ESBL-E extended-spectrum  $\beta$ -lactamase-producing Enterobacterales.

<sup>1</sup> Multiple answers per hospital possible.

<sup>2</sup> Missing data from one hospital.

<sup>3</sup> No separate cleaning due to the use of a 2-in-1 cleaning and disinfection product.

<sup>4</sup> One hospital does not generally change the separation or window curtains after cessation of isolation measures, but only in case of visible contamination or if the patient provides a reason to do so (e.g., demented patient).