

Supplementary Figures and Tables

Supplementary Table 1. MBP-stimulated cytokines responses between binary age groups.

Cytokine Abbreviation	Cytokine Full Name	Age <50 (n=13)	Age ≥50 (n=15)	Unadjusted p-value	FDR
		mean (SD), pg/ml	mean (SD), pg/ml		
BCA-1	B cell-attracting chemokine-1	26.18 (16.65)	19.61 (12.87)	0.25	0.55
EGF	epidermal growth factor	7.04 (6.48)	4.62 (4.81)	0.268	0.57
ENA-78	epithelial-derived neutrophil-activating peptide 78	7858.54 (3051.66)	6029.08 (3872.69)	0.182	0.55
Eotaxin	eotaxin	0.20 (0.38)	0.86 (1.55)	0.149	0.53
Eotaxin.2	eotaxin-2	1058.22 (1562.06)	2167.34 (2144.98)	0.135	0.53
FGF-2	fibroblast growth factor 2	2.52 (2.13)	7.53 (9.85)	0.084	0.53
FLT-3L	fms-like tyrosine kinase receptor 3 ligand	0.39 (0.41)	0.28 (0.20)	0.398	0.6
Fractalkine	fractalkine	3.59 (3.27)	10.22 (11.02)	0.047	0.39
G.CSF	granulocyte colony stimulating factor	7.63 (13.39)	3.59 (5.61)	0.295	0.6
GM-CSF	granulocyte-macrophage colony stimulating factor	29.32 (62.32)	4.13 (8.08)	0.132	0.53
GRO-a	growth related oncogene-alpha	1433.48 (721.58)	1143.09 (797.93)	0.325	0.6
I-309	cytokine I-309	17.74 (38.15)	9.80 (8.77)	0.44	0.63
IFN-a2	interferon alpha 2	2.33 (1.82)	3.42 (5.16)	0.477	0.66
IFN.γ	interferon gamma	8.23 (16.50)	1.27 (1.70)	0.115	0.53
IL-1a	interleukin-1 alpha	2.52 (7.19)	2.90 (4.15)	0.86	0.98
IL-1b	interleukin-1 beta	13.09 (17.89)	17.92 (23.61)	0.552	0.72
IL-1RA	interleukin-1 receptor antagonist	37.84 (89.09)	40.13 (47.67)	0.932	0.98
IL-2	interleukin-2	12.41 (24.20)	6.72 (8.14)	0.398	0.6
IL-3	interleukin-3	2.09 (4.69)	1.21 (1.46)	0.5	0.66
IL-4	interleukin-4	0.28 (0.18)	0.16 (0.12)	0.048	0.39
IL-5	interleukin-5	7.38 (15.77)	1.81 (3.48)	0.194	0.55
IL-6	interleukin-6	686.48 (593.29)	462.24 (549.82)	0.309	0.6
IL-8	interleukin-8	6508.76 (3969.31)	5944.49 (3105.28)	0.677	0.84
IL-9	interleukin-9	2.62 (4.11)	1.11 (1.08)	0.18	0.55
IL-10	interleukin-10	51.96 (46.65)	38.82 (30.34)	0.379	0.6
IL-12p40	interleukin-12 p40 subunit	9.81 (12.40)	9.85 (9.28)	0.992	0.99
IL-12p70	interleukin-12 p70 subunit	0.08 (0.13)	0.30 (0.50)	0.14	0.53
IL-13	interleukin-13	16.96 (29.08)	7.14 (7.61)	0.218	0.55
IL-15	interleukin-15	0.42 (0.50)	0.61 (0.66)	0.413	0.6
IL-16	interleukin-16	33.97 (73.49)	21.02 (52.42)	0.592	0.75
IL-17A	interleukin-17A	9.09 (16.55)	2.60 (2.82)	0.146	0.53
IL-17E (IL-25)	interleukin-17A (also known as interleukin-25)	0.54 (0.44)	1.03 (1.28)	0.205	0.55

IL-17F	interleukin-17F	34.43 (86.32)	3.55 (5.16)	0.177	0.55
IL-18	interleukin-18	0.61 (1.09)	0.66 (1.44)	0.92	0.98
IL-23	interleukin-23	7.60 (9.28)	6.77 (7.98)	0.803	0.97
IL-27	interleukin-27	14.62 (10.14)	19.18 (17.02)	0.407	0.6
IP-10	Interferon gamma-induced protein 10	1725.41 (1963.95)	1836.38 (2045.87)	0.885	0.98
MCP-1	monocyte chemoattractant protein 1	1170.04 (780.66)	1416.12 (730.53)	0.397	0.6
MCP-2	monocyte chemoattractant protein 2	455.48 (490.72)	97.93 (135.06)	0.012	0.23
MCP-3	monocyte chemoattractant protein 3	906.47 (883.40)	300.90 (220.80)	0.016	0.23
MCP-4	monocyte chemoattractant protein 4	140.84 (253.65)	19.65 (46.88)	0.08	0.53
M-CSF	macrophage colony-stimulating factor	4.22 (11.13)	6.31 (3.23)	0.491	0.66
MDC	macrophage-derived chemokine	48.22 (85.43)	286.27 (232.03)	0.002	0.11
MIG (CXCL9)	monokine induced by gamma (also known as C-X-C motif chemokine ligand 9)	1741.65 (2011.68)	1170.79 (1014.53)	0.342	0.6
MIP-1a	macrophage inflammatory protein 1 alpha	34.89 (49.45)	34.37 (49.23)	0.978	0.99
MIP-1b	macrophage inflammatory protein 1 beta	181.07 (172.08)	190.56 (152.31)	0.878	0.98
PDGF-AA	platelet-derived growth factor AA	41.74 (30.94)	20.72 (14.36)	0.026	0.3
PDGF-AB/BB	platelet-derived growth factor AB/BB	357.92 (194.32)	198.19 (109.85)	0.011	0.23
RANTES	regulated on activation, normal T cell expressed and secreted (also known as C-C motif chemokine ligand 5)	18.57 (49.27)	57.93 (107.24)	0.235	0.55
sCD40L	soluble CD40 ligand	2.02 (3.17)	3.22 (4.17)	0.408	0.6
SDF-1a+b	stromal cell-derived factor 1alpha + beta	17.75 (25.39)	17.60 (18.65)	0.986	0.99
TARC	thymus and activation-regulated chemokine	25.29 (34.49)	10.10 (13.04)	0.125	0.53
TGFa	transforming growth factor alpha	1.40 (1.60)	1.33 (0.80)	0.886	0.98
TNFa	tumor necrosis factor alpha	59.83 (109.28)	63.14 (62.58)	0.921	0.98
TNFb	tumor necrosis factor beta	5.36 (7.87)	2.86 (1.77)	0.24	0.55
TRAIL	tumor necrosis factor-related apoptosis-inducing ligand	0.88 (1.60)	0.35 (0.47)	0.229	0.55
VEGF-A	vascular endothelial growth factor A	49.04 (48.17)	35.23 (36.30)	0.396	0.6

Abbreviation: *FDR*, false discovery rate accounting for multiple testing.

Supplementary Table 2. MBP-driven cytokine response as mediators of the association between age and annualized relapse rate.

n=28 (women=20, men=8)		
Fractalkine	Coefficient (95%CI)	p-value
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.270 (-0.428, -0.080)	0.004
Indirect effect	0.005 (-0.136, -0.0446)	0.956
MDC		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.254(-0.457, -0.059)	0.016
Indirect effect	-0.012 (-0.113, -0.090)	0.804
IL-4		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.310 (-0.460, -0.153)	<0.001
Indirect effect	0.045 (-0.037, 0.134)	0.276
MCP-2		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.255 (-0.388, -0.091)	0.004
Indirect effect	-0.011(-0.126, -0.098)	0.956
MCP-3		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.278 (-0.450, -0.108)	<0.001
Indirect effect	0.013 (-0.076, 0.078)	0.604
PDGF-AA		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.286 (-0.462, -0.114)	<0.001
Indirect effect	0.020 (-0.051, 0.113)	0.616
PDGF-AB/BB		
Total effect	-0.265 (-0.410, -0.125)	<0.001
Direct effect	-0.267 (-0.448, -0.099)	0.008
Indirect effect	0.002 (-0.075, 0.103)	0.984

Note: We conducted mediation analyses to assess the extent to which MBP-driven cytokine response explains the association between age and annualized relapse rate in people with multiple sclerosis. We compared the group ≥ 50 years of age to the group < 50 years of age as the reference group. In each mediation analysis, independent variables of the total effect model are age and race/ethnicity. Independent variables of the direct effect model are age, race/ethnicity and MBP-driven cytokine level. Indirect effect was calculated by subtracting direct effect from total effect. We consider a negation when the indirect effect is in the opposite direction as the direct or total effect.

Abbreviations: CI, confidence interval

Supplementary Figure 1. MBP-driven IFN- γ response was not associated with age as a continuous variable (A-C) or binary variable (D-F).

