

Supplementary Material

Supplementary Methods

Definition of confounding variables

We included multiple socioeconomic factors reported by mothers during pregnancy, including occupation (dichotomised into ‘manual’ or ‘non-manual’ based on the 1991 British Office of Population and Census Statistics classification), maternal education (‘Certificate of Secondary Education/vocational’, ‘O level’, ‘A level’, or ‘degree’), home ownership (‘owner or private renter’ or ‘renter or non-homeowner’), financial difficulties score (dichotomised to ‘any difficulties (score of 1 or above)’ or ‘no difficulties (score of 0)’), and smoking pre-pregnancy (‘any smoking’ or ‘no smoking’) as confounders. Other confounders include child ethnicity (‘white’ or ‘non-white’), age at menarche, body mass index (BMI; kg/m²) measured at a clinic visit at 12.8 years, maternal-reported depression in the past 2 years (‘yes’ or ‘no’) collected when the participant was aged 12.1 years, internalising and externalising problems at 9.6 years (each with a score ranging from 0 to 20 and measured with the Strengths and Difficulties Questionnaire),¹ and intelligence quotient at age 8 (measured with the Weschler Intelligence Scale for Children).² Adverse childhood experiences (ACEs), including parental separation, sexual abuse, and physical abuse before age 11, were either reported prospectively by their mother or retrospectively self-reported in adulthood. Multiple questionnaire items corresponded to each ACE (see Supplementary Table 1 for details). Binary variables for each ACE construct (‘any’ or ‘none’) were derived for participants who responded to at least 50% of the relevant questions for a given ACE; for participants with fewer than 50% of the relevant questions the ACE was coded as missing.³

Details of multiple imputation

We included all observed variables used in analyses, including exposures and outcomes, in imputation equations. Additional auxiliary variables, including maternal depression during pregnancy (measured with the Edinburgh Postnatal Depression Scale),⁴ BMI at age 7, and SDQ at age 11, were used to impute maternal depression, BMI, and internalising and externalising problems, respectively. Results were calculated across 60 imputed datasets, guided by results of mcerror tests, which assess statistical reproducibility of the imputation,⁵ and associations were generated by pooling across these datasets using Rubin’s rules.⁶ The prevalence/means of each confounder before and after multiple imputation are presented in Supplementary Table

Supplementary Table 1. Variables used to derive adverse childhood experiences (ACEs), including parental separation, physical abuse, and sexual abuse experienced up to age 11.

Variable ID	ACE	Description	Reported	Retrospective	Age reported	Start time period	End time period
f228	Parental separation	Divorce >CH born	Parent	No	8m	0	0.67
f237	Parental separation	Separation from PTNR >CH born	Parent	No	8m	0	0.67
g308	Parental separation	Mum divorced >CH8MTHs	Parent	No	1yrs9m	0.67	2
g317	Parental separation	Mum and partner separated >CH8MTHs	Parent	No	1yrs9m	0.67	2
h218	Parental separation	Whether mum got divorced since study child was 18 months old and effect this had	Parent	No	2yrs9m	1.5	3
h227	Parental separation	Whether mum and partner separated since study child was 18 months old and effect this had	Parent	No	2yrs9m	1.5	3
j308	Parental separation	MUM Divorced> CH 30 MTHs	Parent	No	3yrs11m	2.5	4
j317	Parental separation	MUM & PTR Separated> CH 30 MTHs	Parent	No	3yrs11m	2.5	4
k4008	Parental separation	Mother was divorced in past year	Parent	No	5yrs1m	4	5
k4017	Parental separation	Mother and partner separated in past year	Parent	No	5yrs1m	4	5
l4008	Parental separation	Respondent was divorced since study child's 5th birthday	Parent	No	6yrs1m	5	6
l4017	Parental separation	Respondent separated from partner since study child's 5th birthday	Parent	No	6yrs1m	5	6
p2008	Parental separation	Mother was divorced since the study child's 6th birthday	Parent	Yes	9yrs2m	6	7
p2017	Parental separation	Mother and husband/partner separated since the study child's 6th birthday	Parent	Yes	9yrs2m	6	7
r5008	Parental separation	Respondent has been divorced since child's 9th birthday	Parent	Yes	11yrs2m	9	10
r5017	Parental separation	Respondent separated from husband/partner since the study child's 9th birthday	Parent	Yes	11yrs2m	9	10
f246	Physical abuse	PTNR physically cruel to CHDR >CH born	Parent	No	8m	0	0.67
f247	Physical abuse	MUM physically cruel to CHDR >CH born	Parent	No	8m	0	0.67
g326	Physical abuse	Partner physically cruel to children >CH8MTHs	Parent	No	1yrs9m	0.67	2
g327	Physical abuse	Mum physically cruel to children >CH8MTHs	Parent	No	1yrs9m	0.67	2
h236	Physical abuse	Whether partner was physically cruel to children since study child was 18 months old and effect this had	Parent	No	2yrs9m	1.5	3
h237	Physical abuse	Whether mum was physically cruel to children since study child was 18 months old and effect this had	Parent	No	2yrs9m	1.5	3
j326	Physical abuse	PTR PHYS Cruel to CDRN> CH 30 MTHs	Parent	No	3yrs11m	2.5	4
j327	Physical abuse	MUM PHYS Cruel to CDRN> CH 30 MTHs	Parent	No	3yrs11m	2.5	4
k4026	Physical abuse	Mothers partner was physically cruel to children in past year	Parent	No	5yrs1m	4	5
k4027	Physical abuse	Mother was physically cruel to children in past year	Parent	No	5yrs1m	4	5
l4026	Physical abuse	Respondent's partner physically cruel to respondent's children since study child's 5th birthday	Parent	No	6yrs1m	5	6
l4027	Physical abuse	Respondent physically cruel to own children since study child's 5th birthday	Parent	No	6yrs1m	5	6
p2026	Physical abuse	Mother's husband/partner was physically cruel to her children since the study child's 6th birthday	Parent	Yes	9yrs2m	6	7
p2027	Physical abuse	Mother was physically cruel to her children since the study child's 6th birthday	Parent	Yes	9yrs2m	6	7
r5026	Physical abuse	Respondent's husband/partner was physically cruel to their children since study child's 9th birthday	Parent	Yes	11yrs2m	9	10
r5027	Physical abuse	Respondent was physically cruel to their children since the study child's 9th birthday	Parent	Yes	11yrs2m	9	10
ypb8002	Physical abuse	Frequency adult in family pushed, grabbed or shoved respondent before age of 11	Child	Yes	22yrs	0	11
ypb8003	Physical abuse	Frequency adult in family smacked respondent for discipline before age of 11	Child	Yes	22yrs	0	11
ypb8007	Physical abuse	Frequency adult in family hit respondent so hard it left bruises or marks before age of 11	Child	Yes	22yrs	0	11
kf455a	Sexual abuse	Child sexually abused > 18 months, Y/N	Parent	No	30m	1.5	2.5
kj465	Sexual abuse	CH was Sexually Abused Past 12 MTHs	Parent	No	42m	2.5	3.5
kl475	Sexual abuse	Child was sexually abused since age 3	Parent	No	57m	3	5

kn4005	Sexual abuse	Child sexually abused in past 15 months	Parent	No	69m	4.5	6
kq365	Sexual abuse	Child was sexually abused since his/her 5th birthday	Parent	No	81m	5	7
kt5005	Sexual abuse	Since 7th birthday child has been sexually abused	Parent	No	105m	7	9
ypb8030	Sexual abuse	Respondent was touched in a sexual way by adult or older child, or was forced to touch adult or older child in a sexual way, before age of 11	Child	Yes	22yrs	0	11
ypb8040	Sexual abuse	Adult or older child forced, or attempted to force, respondent into any sexual activity by threatening or holding respondent down or hurting respondent in some way, before age of 11	Child	Yes	22yrs	0	11

Information provided by Houtepen et al. (2018).³

Supplementary Table 2. Proportion of missing data in the sample of individuals who provided data on exposures and outcomes (N= 2698) and comparison of prevalence of confounders in the incomplete and imputed samples, and the full female cohort.

Variable	Prevalence/mean (SD) in sample restricted to those reporting on exposures and outcomes	Proportion missing data in sample restricted to those reporting on exposures and outcomes	Prevalence/mean in imputed dataset	Prevalence/mean (SD) in the full female cohort
Non-white ethnicity	3.72%	9.23%	3.87%	5.05%
Maternal education	21.10% CSE 38.16% O level 26.16% A level 14.57% Degree	12.01%	22.37% CSE 38.09% O level 25.51% A level 14.02% Degree	25.14% CSE 36.89% O level 24.01% A level 13.96% Degree
Manual parental social class	46.58%	12.16%	47.76%	49.23%
Any financial difficulties	60.55%	10.27%	61.06%	63.92%
Non-homeowner or renter	13.55%	8.08%	13.79%	19.90%
Any maternal smoking pre-pregnancy	26.93%	6.82%	27.11%	33.02%
Any maternal depression	22.78%	21.91%	23.62%	23.29%
Any parental separation	22.82%	12.12%	23.53%	24.95%
Any sexual abuse	3.31%	13.83%	3.53%	3.15%
Any physical abuse	38.76%	12.97%	38.39%	32.78%
Age at menarche	12.66 (1.11)	2.89%	12.66	12.62 (1.17)
BMI	20.21 (3.58)	21.42%	20.40	20.10 (3.59)
SDQ internal	2.59 (2.60)	17.12%	2.63	2.71 (2.66)
SDQ external	3.68 (2.91)	17.09%	3.74	3.77 (2.89)
IQ	104.59 (15.57)	22.46%	103.50	103.86 (15.98)

Abbreviations: SD, standard deviation; CSE, certificate of secondary education; BMI, body mass index; SDQ, strengths and difficulties questionnaire; IQ, intelligence quotient.

Supplementary Results

Supplementary Table 3. Linear regression analysis of the association between menstrual symptoms and school absences and GCSE score and logistic regression analysis of the association between menstrual symptoms and achieving five A*-C GCSEs including Maths and English and persistent absence (10% or more) (N=2698).

Continuous Outcomes						
Exposure	School absence in Year 11			GCSE points score		
	% change	95% CI	P value	Beta	95% CI	P Value
Heavy or prolonged bleeding						
Crude	22.71	15.20, 30.71	<0.0001	-16.97	-22.70, -11.24	<0.0001
Adjusted	16.58	9.43, 24.20	<0.0001	-5.65	-10.10, -1.20	0.0129
Menstrual pain						
Crude	16.28	9.37, 23.62	<0.0001	-3.42	-8.99, 2.14	0.2279
Adjusted	12.83	6.14, 19.95	0.0001	-3.14	-7.46, 1.17	0.1532
Binary Outcomes						
Exposure	Persistent absence in Year 11			Five A*-C GCSEs incl. Maths & English		
	OR	95% CI	P value	OR	95% CI	P Value
Heavy or prolonged bleeding						
Crude	1.69	1.65, 1.73	<0.0001	0.60	0.59, 0.62	<0.0001
Adjusted	1.48	1.45, 1.52	<0.0001	0.73	0.71, 0.75	<0.0001
Menstrual pain						
Crude	1.51	1.47, 1.54	<0.0001	0.91	0.89, 0.92	<0.0001
Adjusted	1.42	1.39, 1.46	<0.0001	0.84	0.81, 0.86	<0.0001

Abbreviations: GCSE, general certificate of secondary education; OR, odds ratio. Adjusted for ethnicity; maternal education, parental social class, financial difficulties, and home ownership during pregnancy; maternal smoking pre-pregnancy; parental separation, physical abuse, and sexual abuse before 11; maternal depression at 12.1; age at menarche; body mass index at 12.8; internalising and externalising problems at 9.6; and intelligence quotient at 8.

Supplementary Table 4. Linear regression analysis of the association between different categories of menstrual symptoms and GCSE score and school absences (N=2698).

		School absence in Year 11			GCSE points score		
		% increase	95% CI	P value	Beta	95% CI	P Value
Heavy or prolonged bleeding and visiting a doctor							
Crude analysis	No symptom (ref) (N=2353)	0	-	-	0	-	-
	Symptom, no doctor (N=967)	15.21	7.68, 23.27	<0.0001	-14.08	-20.23, -7.93	<0.0001
	Symptom and doctor (N=246)	57.36	39.53, 77.47	<0.0001	-28.29	-39.22, -17.35	<0.0001
Adjusted analysis	No symptom (ref) (N=2353)	0	-	-	0	-	-
	Symptom, no doctor (N=967)	10.93	3.70, 18.67	0.0026	-5.45	-10.23, -0.68	0.0252
	Symptom and doctor (N=246)	42.61	26.41, 60.89	<0.0001	-5.92	-14.56, 2.72	0.1791
Menstrual pain and visiting a doctor							
Crude analysis	No symptom (ref) (N=1625)	0	-	-	0	-	-
	Symptom, no doctor (N=1616)	10.65	3.88, 17.86	0.0017	-0.54	-6.30, 5.23	0.8552
	Symptom and doctor (N=292)	55.64	38.70, 74.64	<0.0001	-20.08	-30.6, -9.57	0.0002
Adjusted analysis	No symptom (ref) (N=1625)	0	-	-	0	-	-
	Symptom, no doctor (N=1616)	8.76	2.13, 15.83	0.0089	-2.53	-6.99, 1.94	0.2667
	Symptom and doctor (N=292)	42.30	26.74, 59.77	<0.0001	-6.77	-15.00, 1.46	0.1068
Heavy and prolonged, separately and combined							
Crude analysis	Neither (ref) (N=2013)	0	-	-	0	-	-
	Prolonged only (N=286)	-8.48	-18.37, 2.61	0.1290	4.78	-5.59, 15.15	0.3663
	Heavy only (N=882)	20.74	12.46, 29.62	<0.0001	-17.24	-23.69, -10.8	<0.0001
	Both (N=307)	23.22	10.49, 37.42	0.0002	-13.73	-23.63, -3.84	0.0065
Adjusted analysis	Neither (ref) (N=2013)	0	-	-	0	-	-
	Prolonged only (N=286)	-7.54	-17.39, 3.48	0.1723	5.15	-2.84, 13.15	0.2064
	Heavy only (N=882)	14.69	6.84, 23.13	0.0002	-4.82	-9.81, 0.16	0.0580
	Both (N=307)	17.71	5.65, 31.14	0.0031	-5.55	-13.19, 2.10	0.1549
Heavy and pain, separately and combined							
Crude analysis	Neither (ref) (N=1280)	0	-	-	0	-	-
	Pain only (N=952)	10.44	2.31, 19.21	0.0109	1.18	-5.76, 8.12	0.7388
	Heavy only (N=276)	19.40	6.28, 34.13	0.0028	-19.78	-30.35, -9.21	0.0002
	Both (N=920)	30.86	21.22, 41.26	<0.0001	-15.46	-22.4, -8.51	<0.0001
Adjusted analysis	Neither (ref) (N=1280)	0	-	-	0	-	-
	Pain only (N=952)	8.84	0.88, 17.42	0.0288	-2.88	-8.24, 2.48	0.2925
	Heavy only (N=276)	14.21	1.78, 28.14	0.0238	-8.18	-16.34, -0.01	0.0498
	Both (N=920)	23.13	14.02, 32.96	<0.0001	-6.53	-11.94, -1.11	0.0182

Abbreviations: GCSE, general certificate of secondary education. Adjusted for ethnicity; maternal education, parental social class, financial difficulties, and home ownership during pregnancy; maternal smoking pre-pregnancy; parental separation, physical abuse, and sexual abuse before 11; maternal depression at 12.1; age at menarche; body mass index at 12.8; internalising and externalising problems at 9.6; and intelligence quotient at 8.

The following results are from analyses conducted in the complete case samples only.

Supplementary Table 5. Distribution of sociodemographic factors, adverse childhood experiences, and child factors according to adolescent menstrual symptoms in complete case sample only.

	Heavy or Prolonged Bleeding		Menstrual Pain	
	Yes ≤ 972 (36.03%)	No ≤ 1726 (63.97%)	Yes ≤ 1496 (55.45%)	No ≤ 1202 (44.55%)
SOCIODEMOGRAPHIC FACTORS				
Ethnicity				
White	830 (96.51)	1528 (96.16)	1301 (96.44)	1057 (96.09)
Non-white	30 (3.49)	61 (3.84)	48 (3.56)	43 (3.91)
Maternal education				
CSE/Vocational	195 (23.75)	306 (19.70)	276 (21.15)	225 (21.05)
O level	327 (39.83)	579 (37.28)	486 (37.24)	420 (39.29)
A level	214 (26.07)	407 (26.21)	354 (27.13)	267 (24.98)
Degree	85 (10.35)	261 (16.81)	189 (14.48)	157 (14.69)
Parental social class				
Manual	423 (50.78)	681 (44.31)	609 (46.56)	495 (46.61)
Non-manual	410 (49.22)	856 (55.69)	699 (53.44)	567 (53.39)
Financial difficulties				
Any	556 (64.95)	910 (58.15)	826 (61.60)	640 (59.26)
None	300 (35.05)	655 (41.85)	515 (38.40)	440 (40.74)
Home ownership				
Renter or non-homeowner	144 (16.36)	192 (12.00)	194 (14.10)	142 (12.86)
Owner or private renter	736 (83.64)	1408 (88.00)	1182 (85.90)	962 (87.14)
Maternal smoking pre-pregnancy				
Yes	277 (30.88)	400 (24.74)	409 (29.34)	268 (23.93)
No	620 (69.12)	1217 (75.26)	985 (70.66)	852 (76.07)
ADVERSE CHILDHOOD EXPERIENCES				
Parental separation before age 11				
Yes	212 (25.27)	329 (21.48)	300 (22.76)	241 (22.89)
No	627 (74.73)	1203 (78.52)	1018 (77.24)	812 (77.11)
Physical abuse before age 11				
Any	317 (38.33)	593 (38.99)	533 (40.69)	377 (36.32)
None	510 (61.67)	928 (61.01)	777 (59.31)	661 (63.68)
Sexual abuse before age 11				
Any	38 (4.67)	39 (2.58)	42 (3.28)	35 (3.36)
None	775 (95.33)	1473 (97.42)	1240 (96.72)	1008 (96.64)
Maternal depression at 12.1 years				
Yes	213 (29.42)	267 (19.31)	291 (25.15)	189 (19.89)
No	511 (70.58)	1116 (80.69)	866 (74.85)	761 (80.11)
CHILD FACTORS				
Age at menarche				
Mean (SE)	12.60 (1.12)	12.69 (1.10)	12.53 (1.08)	12.83 (1.31)
BMI at 12.8 years				
Mean (SE)	20.40 (3.54)	20.11 (3.59)	20.39 (3.16)	19.99 (3.64)
Internalising SDQ at 9.6 years				
Mean (SE)	2.74 (2.72)	2.51 (2.53)	2.76 (2.69)	2.38 (2.47)
Externalising SDQ at 9.6 years				
Mean (SE)	4.05 (3.08)	3.49 (2.79)	3.82 (2.91)	3.52 (2.91)
IQ at 8 years				
Mean (SE)	103.31 (15.86)	105.26 (15.38)	104.87 (15.36)	104.23 (15.83)
Past year oral contraception at exposure timepoint				
Yes	239 (24.95)	168 (9.85)	290 (19.58)	117 (9.89)
No	719 (75.05)	1538 (90.15)	1191 (80.42)	1066 (90.11)

Abbreviations: SE, standard error; CSE, certificate of secondary education; BMI, body mass index; SDQ, strengths and difficulties questionnaire; IQ, intelligence quotient.

Supplementary Table 6. Linear regression analysis of the association between menstrual symptoms and school absences and GCSE score and logistic regression analysis of the association between menstrual symptoms and achieving five A*-C GCSEs including Maths and English and persistent absence (10% or more) in the complete case sample only (N=1258).

Exposure	Continuous Outcomes					
	School absence in Year 11			GCSE points score		
	% change	95% CI	P value	Beta	95% CI	P Value
Heavy or prolonged bleeding						
Crude	14.43	4.63, 25.14	0.0032	-13.17	-20.36, -5.98	0.0003
Adjusted	11.24	1.78, 21.57	0.0188	-6.12	-11.60, -0.64	0.0286
Menstrual pain						

Crude	15.40	6.12, 25.48	0.0008	-2.79	-9.56, 3.97	0.4181
Adjusted	11.06	2.11, 20.79	0.0145	-2.47	-7.67, 2.72	0.3503
Binary Outcomes						
Exposure	Persistent absence in Year 11			Five A*-C GCSEs incl. Maths & English		
	OR	95% CI	P value	OR	95% CI	P Value
Heavy or prolonged bleeding						
Crude	1.41	1.05, 1.90	0.0233	0.67	0.51, 0.88	0.0039
Adjusted	1.27	0.93, 1.73	0.1290	0.81	0.57, 1.14	0.2184
Menstrual pain						
Crude	1.48	1.10, 1.99	0.0087	0.94	0.72, 1.23	0.6671
Adjusted	1.40	1.03, 1.90	0.0324	0.92	0.66, 1.29	0.6387

Abbreviations: GCSE, general certificate of secondary education; OR, odds ratio. Adjusted for ethnicity; maternal education, parental social class, financial difficulties, and home ownership during pregnancy; maternal smoking pre-pregnancy; parental separation, physical abuse, and sexual abuse before 11; maternal depression at 12.1; age at menarche; body mass index at 12.8; internalising and externalising problems at 9.6; and intelligence quotient at 8.

Supplementary Table 7. Linear regression analysis of the associations between different categories of menstrual symptoms and school absences and GCSE score in the complete case sample only (N=1258).

		School absence in Year 11			GCSE points score		
		% increase	95% CI	P value	Beta	95% CI	P Value
Heavy doctor							
Crude analysis	No symptom (ref)	0	-	-	0	-	-
	Symptom, no doctor	8.03	-1.69, 18.71	0.1084	-10.82	-18.42, -3.22	0.0053
	Symptom and doctor	52.99	25.72, 86.17	<0.0001	-24.99	-40.81, -9.17	0.0020
Adjusted analysis	No symptom (ref)	0	-	-	0	-	-
	Symptom, no doctor	5.92	-3.54, 16.31	0.2277	-5.84	-11.63, -0.05	0.0482
	Symptom and doctor	42.22	17.03, 72.83	0.0004	-6.34	-18.41, 5.72	0.3023
Pain doctor							
Crude analysis	No symptom (ref)	0	-	-	0	-	-
	Symptom, no doctor	11.38	2.25, 21.32	0.0136	-0.45	-7.38, 6.47	0.8976
	Symptom and doctor	55.47	29.64, 86.44	<0.0001	-22.48	-37.18, -7.77	0.0028
Adjusted analysis	No symptom (ref)	0	-	-	0	-	-
	Symptom, no doctor	7.97	-0.89, 17.62	0.0793	-0.97	-6.27, 4.33	0.7194
	Symptom and doctor	43.28	19.38, 71.95	0.0001	-16.05	-27.34, -4.77	0.0053
Heavy and prolonged, separately and combined							
Crude analysis	Neither (ref)	0	-	-	0	-	-
	Prolonged only	-9.63	-22.36, 5.18	0.1906	1.77	-10.43, 13.96	0.7761
	Heavy only	12.58	1.87, 24.43	0.0203	-14.41	-22.45, -6.37	0.0005
	Both	14.28	-2.95, 34.57	0.1093	-8.04	-21.17, 5.10	0.2301
Adjusted analysis	Neither (ref)	0	-	-	0	-	-
	Prolonged only	-8.82	-21.54, 5.96	0.2282	5.40	-3.87, 14.67	0.2530
	Heavy only	8.79	-1.49, 20.14	0.0962	-6.38	-12.50, -0.26	0.0412
	Both	13.79	-3.21, 33.78	0.1174	-2.24	-12.22, 7.74	0.6603
Heavy and pain, separately and combined							
Crude analysis	Neither (ref)	0	-	-	0	-	-
	Pain only	10.99	0.32, 22.79	0.0433	3.59	-4.55, 11.72	0.3875
	Heavy only	6.80	-9.56, 26.11	0.4378	-5.92	-19.30, 7.46	0.3854
	Both	24.44	11.63, 38.72	0.0001	-13.28	-22.02, -4.53	0.0030
Adjusted analysis	Neither (ref)	0	-	-	0	-	-
	Pain only	7.10	-3.20, 18.49	0.1835	0.74	-5.51, 6.98	0.8172
	Heavy only	4.38	-11.42, 23.00	0.6083	-1.44	-11.58, 8.70	0.7807
	Both	18.32	6.14, 31.90	0.0024	-7.16	-13.87, -0.46	0.0364

Abbreviations: GCSE, general certificate of secondary education. Adjusted for ethnicity; maternal education, parental social class, financial difficulties, and home ownership during pregnancy; maternal smoking pre-pregnancy; parental separation, physical abuse, and sexual abuse before 11; maternal depression at 12.1; age at menarche; body mass index at 12.8; internalising and externalising problems at 9.6; and intelligence quotient at 8.

Supplementary References

1. Goodman R. The strengths and difficulties questionnaire: A research note. *J Child Psychol Psychiatry*. 1997 Jul;38(5):581–6.
2. Wechsler D, Golombok S, Rust J. WISC-III Wechsler Intelligence Scale for Children - Third Edition UK Manual. Sidcup, UK: The Psychological Corporation; 1992.
3. Houtepen LC, Heron J, Suderman MJ, Tilling K, Howe LD. Adverse childhood experiences in the children of the avon longitudinal study of parents and children (ALSPAC). *Wellcome Open Res*. 2018;3.

4. Cox JL, Holden JM, Sagovsky R. Detection of Postnatal Depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* [Internet]. 1987 Jun 2;150(6):782–6. Available from: https://www.cambridge.org/core/product/identifier/S0007125000214712/type/journal_article
5. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med*. 2011 Feb 20;30(4):377–99.
6. Rubin D. Multiple Imputation for Nonresponse in Surveys [Internet]. Hoboken, NJ, USA: John Wiley & Sons, Inc; 1987 [cited 2022 Aug 6]. Available from: <https://onlinelibrary.wiley.com/doi/book/10.1002/9780470316696>